

## Sections of Convex Bodies in John's and Minimal Surface Area Position

David Alonso-Gutiérrez<sup>1</sup> and Silouanos Brazitikos<sup>2,\*</sup>

<sup>1</sup>Área de Análisis Matemático, Departamento de Matemáticas, Facultad de Ciencias, Instituto Universitario de Matemáticas y Aplicaciones, Universidad de Zaragoza, Pedro Cerbuna 12, 50009 Zaragoza, Spain and

<sup>2</sup>Department of Mathematics, National and Kapodistrian University of Athens, Panepistimioupolis 157-84, Athens, Greece

\*Correspondence to be sent to: e-mail: [silouanb@gmail.com](mailto:silouanb@gmail.com)

We prove several estimates for the volume, the mean width, and the value of the Will's functional of sections of convex bodies in John's position, as well as for their polar bodies. These estimates extend some well-known results for convex bodies in John's position to the case of lower-dimensional sections, which had mainly been studied for the cube and the regular simplex. Some estimates for centrally symmetric convex bodies in minimal surface area position are also obtained.

### 1 Introduction and Notation

For any convex body  $K \subseteq \mathbb{R}^n$  (i.e., a compact convex set with non-empty interior), a position of  $K$  will be any affine image of  $K$ . Every position of the Euclidean ball,  $B_2^n$ , is called an ellipsoid. A well-known theorem by John (see [29]) states that every convex body  $K \subseteq \mathbb{R}^n$  has a unique maximal volume ellipsoid,  $\mathcal{E}(K)$ , contained in it. The volume ratio of  $K$  is defined as

$$\text{v.rat}(K) := \left( \frac{|K|}{|\mathcal{E}(K)|} \right)^{1/n}.$$

Received December 18, 2020; Revised July 8, 2021; Accepted August 16, 2021  
Communicated by Prof. Shiri Artstein-Avidan

Here and in what follows  $|\cdot|$  denotes the volume of a convex body in the appropriate dimension. Notice that the volume ratio does not depend on the position of the convex body  $K$ .

A convex body  $K \subseteq \mathbb{R}^n$  is said to be in John's position if the maximal volume ellipsoid contained in  $K$  is the Euclidean unit ball  $B_2^n$ . In other words,  $K$  is in John's position if  $B_2^n$  is contained in  $K$  and for every non-degenerate linear map  $T \in GL(n)$  and every  $a \in \mathbb{R}^n$  such that  $a + T(B_2^n) \subseteq K$  we have that  $|a + T(B_2^n)| = |T(B_2^n)| \leq |B_2^n|$ . By the uniqueness of  $\mathcal{E}(K)$ , this position is uniquely determined up to orthogonal transformations. Denoting by  $B_\infty^n$  the  $n$ -dimensional cube and by  $S_n$  the centered regular simplex with inradius  $r(S_n) = 1$  in  $\mathbb{R}^n$ , it is well known that both  $B_\infty^n$  and  $S_n$  are in John's position.

Ball proved in [6] that the simplex maximizes the volume ratio among all convex bodies in  $\mathbb{R}^n$  and the cube maximizes the volume ratio among all the centrally symmetric convex bodies in  $\mathbb{R}^n$ . The proof consists of the following three steps: first, since the volume ratio of a convex body does not depend on its position, it can be assumed that the convex body is John's position; second, substitute the convex body by a polytope that contains the convex body and it is also in John's position; finally, obtain an upper bound on the volume of such a polytope by using Brascamp–Lieb inequality (see Theorem 2.1 below).

Dual to John's position is the so-called Löwner's position. A convex body is said to be in Löwner's position if the minimal volume ellipsoid containing it is the Euclidean unit ball. It is well known (see, for instance, [4, Proposition 4.7]) that a convex body  $K \subseteq \mathbb{R}^n$  is in John's position if and only if  $K^\circ$  is in Löwner's position. Here  $K^\circ$  denotes the polar body of  $K$  defined by

$$K^\circ = \{x \in \mathbb{R}^n : \langle x, y \rangle \leq 1, \forall y \in K\}.$$

Ball observed in [6] that a reverse form of the Brascamp–Lieb inequality would provide that among all convex bodies in Löwner's position the centered regular simplex  $\tilde{S}_n$  with circumradius  $R(\tilde{S}_n) = 1$  has the smallest volume. Moreover, among all centrally symmetric convex bodies in Löwner's position, the  $\ell_1^n$ -ball,  $B_1^n$ , has the smallest volume. The needed reverse form of the Brascamp–Lieb inequality was obtained by Barthe in [9] (see Theorem 2.1 below).

Vaaler showed in [43] that if  $F_0 \in G_{n,k}$  is a  $k$ -dimensional linear subspace of  $\mathbb{R}^n$  in the Grassmannian manifold  $G_{n,k}$ , then  $|B_\infty^n \cap F_0| \geq |B_\infty^k|$ . Ball obtained in [5] a reverse

inequality proving that  $|B_\infty^n \cap F_0| \leq 2^{\frac{n-k}{2}} |B_\infty^k|$ . He also obtained the bound

$$|B_\infty^n \cap F_0|^{1/k} \leq \sqrt{\frac{n}{k}} |B_\infty^k|^{1/k}, \quad (1.1)$$

which is optimal if and only if  $k \mid n$  (see [26]).

It follows from the results of Ball [7] that the  $k$ -dimensional sections of a regular simplex with largest volume are exactly its  $k$ -dimensional faces. Webb showed in [44] that for every hyperplane through the origin  $F_0 \in G_{n,n-1}$ ,

$$|S_n \cap F_0|^{\frac{1}{n-1}} \leq \frac{1}{(\sqrt{2n(n+1)})^{\frac{1}{n-1}}} \sqrt{\frac{n(n+1)}{n-1}} |S_{n-1}|^{\frac{1}{n-1}}. \quad (1.2)$$

There is equality for the sections passing through the origin that contain  $n - 1$  of the vertices.

Dirksen proved in [15, Theorem 6.1] (see also [16]) the following estimate for the volume of  $k$ -dimensional sections of  $S_n$  through the origin:

$$|S_n \cap F_0|^{1/k} \leq \frac{1}{(k+1)^{\frac{n-k}{2k(n+1)}}} \sqrt{\frac{n(n+1)}{k(k+1)}} |S_k|^{1/k} \quad (1.3)$$

for every  $F_0 \in G_{n,k}$ . Besides, this estimate is asymptotically sharp.

The proof of these volume estimates for sections of the cube and the regular simplex follow the lines of Ball's upper bound of the volume ratio. However, only sections of  $B_\infty^n$  and  $S_n$ , which are two particular convex bodies, are being considered. Passing to the general case, we observe that it is not possible to obtain an upper bound for the volume of sections of a general centered convex body  $K$  without any additional assumption since, considering different positions of  $K$ , we can obtain sections with volume as large as desired. However,  $B_\infty^n$  and  $S_n$  are in John's position. Motivated by these, we consider convex bodies in John's position and generalize the above results (and other known results) to sections of such convex bodies. We also obtain some estimates for sections of convex bodies in minimal surface area position.

### 1.1 Volume of sections of convex bodies in John's position

In a recent article [32], Markessinis claimed to have obtained an upper bound for the volume of  $k$ -dimensional central sections of convex bodies in John's position. However, although the estimate given for central sections of centrally symmetric convex bodies in

John's position is correct, the proof in the not necessarily symmetric case is not correct. In the following theorem, we give an upper bound for the volume of central (and non-central)  $k$ -dimensional sections of an arbitrary convex body in John's position.

**Theorem 1.1.** Let  $K \subseteq \mathbb{R}^n$  be a convex body in John's position and  $F_0 \in G_{n,k}$ . Then

$$|K \cap F_0|^{1/k} \leq \frac{1}{(k+1)^{\frac{n-k}{2k(n+1)}}} \sqrt{\frac{n(n+1)}{k(k+1)}} |S_k|^{1/k}.$$

Furthermore, if  $K$  is centrally symmetric

$$|K \cap F_0|^{1/k} \leq \sqrt{\frac{n}{k}} |B_\infty^k|^{1/k}.$$

Moreover, if  $F_h$  is a  $k$ -dimensional affine subspace at distance  $h$  from the origin and  $K$  is a convex body in John's position, then

$$|K \cap F_h|^{1/k} \leq \sqrt{\frac{n(n+1)^{1+\frac{1}{k}}}{k(k+1)^{1+\frac{1}{k}}} \left(\frac{n}{n+h^2}\right)^{\frac{1}{2k}}} |S_k|^{1/k}.$$

**Remark.** The proof in the symmetric case is the same as the one given by Markessinis, which follows Ball's ideas in [5]. Nevertheless, we will reproduce it for the sake of completeness. We can also obtain it as a direct consequence of Theorem 1.5, as well as a consequence of Theorem 8.1 below (see Section 2.6). This estimate is a sharp generalization of Ball's estimate (1.1) for the cube. Moreover, the case  $k = 1$  gives one more proof of John's theorem in the symmetric case: if  $K$  is a centrally symmetric convex body in  $\mathbb{R}^n$  whose maximal volume ellipsoid is  $B_2^n$  then  $K \subseteq \sqrt{n}B_2^n$ .

**Remark.** Many other generalizations and extensions of Ball's estimates for sections of  $B_\infty^n$  have been obtained, for instance, in [28], [30], or [31].

**Remark.** The proof in the non-symmetric case follows Dirksen's ideas from [16]. However, the decomposition of the identity in a linear subspace of  $\mathbb{R}^{n+1}$  in order to apply Theorem 2.1 is not obtained by projecting the vectors in an orthonormal basis of  $\mathbb{R}^{n+1}$ . It comes out by projecting the vectors providing a more general decomposition of the identity in  $\mathbb{R}^{n+1}$ . As a consequence, a different maximization problem from the one in Dirksen's proof has to be considered. Notice that we recover the estimate in (1.3), which is asymptotically sharp for the simplex. Besides, if we take non-central sections by

$k$ -dimensional subspaces a distance  $h = \sqrt{\frac{n(n-k)}{(k+1)}}$  from the origin, which is the distance from the origin to any  $k$ -dimensional face of  $S_n$ , we obtain that

$$|K \cap F_h|^{1/k} \leq \sqrt{\frac{n(n+1)}{k(k+1)}} |S_k|^{1/k},$$

which is exactly the volume of the  $k$ -dimensional faces of  $S_n$ . The estimate for general affine subspaces can also be obtained as a direct consequence of Theorem 8.1 below.

**1.2 Volume of projections of convex bodies in Löwner's position**

Let us recall that if  $F_0 \in G_{n,k}$  is a linear subspace and  $K \cap F_0$  contains the origin in its relative interior, the polar body of  $K \cap F_0$  in  $F_0$  is  $P_{F_0}K^\circ$ , the projection of the polar body of  $K$  onto  $F_0$ . Contrary to the case of the volume of sections of the cube, not much is known about the projections of the cross-polytope. For example, a dual statement to Vaaler's theorem, claiming that if  $F_0 \in G_{n,k}$  then  $|P_{F_0}(B_1^n)| \leq |B_1^k|$ , has only been confirmed if  $k = 2, 3$ , and  $n - 1$  (see [8], [12], and [27]) and a dual statement of Ball's upper bound  $|B_\infty^n \cap F_0| \leq 2^{\frac{n-k}{2}} |B_\infty^k|$  has only been proved when  $k = 2$  or  $k = n - 1$  (see [12] and [27]). Nevertheless, concerning the volume of polar bodies of sections of convex bodies in John's position (i.e., projections of convex bodies in Löwner's position whenever they contain the origin in its interior), it was proved by Barthe in his PhD thesis (see also [1]) that in the case of the  $\ell_p^n$ -balls, if  $1 \leq p \leq 2$  and  $F_0 \in G_{n,k}$ , then

$$|P_{F_0}(B_p^n)|^{1/k} \geq \left(\frac{k}{n}\right)^{\frac{1}{p}-\frac{1}{2}} |B_p^k|^{1/k},$$

where  $P_{F_0}$  denotes the orthogonal projection onto  $F_0$ . In particular, we have the following estimate for the projections of  $B_1^n$ : for every  $F_0 \in G_{n,k}$

$$|P_{F_0}(B_1^n)|^{1/k} \geq \sqrt{\frac{k}{n}} |B_1^k|^{1/k}.$$

We obtain a similar lower bound for the volume of  $k$ -dimensional projections of convex bodies in Löwner's position.

**Theorem 1.2.** Let  $K \subseteq \mathbb{R}^n$  be a convex body in Löwner's position and  $F_0 \in G_{n,k}$ . Then

$$|P_{F_0}(K)|^{1/k} \geq \sqrt{\frac{k}{n}} |\tilde{S}_k|^{1/k}.$$

Furthermore, if  $K$  is centrally symmetric

$$|P_{F_0}(K)|^{1/k} \geq \sqrt{\frac{k}{n}} |B_1^k|^{1/k}.$$

**Remark.** In the symmetric case, the proof follows the idea of the proof of the aforementioned result for  $B_p^n$ . This relies on the use of the reverse Brascamp–Lieb inequality (see Theorem 2.1 below) together with the use of a decomposition of the identity in a linear subspace of  $\mathbb{R}^n$ . As in the proof of Theorem 1.1, unlike in the case in which  $K = B_p^n$ , the decomposition of the identity in the linear subspace does not arise by projecting the canonical basis, but by projecting the vectors in a general decomposition of the identity in  $\mathbb{R}^n$ .

In the non-symmetric case, the proof follows the idea of the proof of Ball’s observation in [6] together with the reverse form of Brascamp–Lieb inequality. Again, in this case, the decomposition of the identity in a linear subspace in  $\mathbb{R}^{n+1}$  arises by projecting the vectors in a decomposition of the identity in  $\mathbb{R}^{n+1}$  rather than an orthonormal basis in  $\mathbb{R}^{n+1}$ . Some estimates of the volume of projections of the regular simplex were obtained in [19]. However, the estimates do not rely on the use of the reverse Brascamp–Lieb inequality.

### 1.3 The mean width

The mean width of a convex body  $K \subseteq \mathbb{R}^n$  is defined by

$$w(K) = \int_{S^{n-1}} h_K(\theta) d\sigma(\theta),$$

where, for every  $x \in \mathbb{R}^n$ ,  $h_K(x) := \sup\{\langle x, y \rangle : y \in K\}$  is the support function of  $K$  at  $x$  and  $d\sigma$  denotes the uniform probability measure on  $S^{n-1}$ . In [40], the authors proved that among all centrally symmetric convex bodies in John’s position in  $\mathbb{R}^n$ ,  $w(K)$  is maximized when  $K = B_\infty^n$ . The not necessarily symmetric case was treated in [10], where it was proved that among all convex bodies in John’s position in  $\mathbb{R}^n$ ,  $w(K)$  is maximized when  $K = S_n$ , where  $S_n$  denotes the regular simplex in John’s position. If we pass to the mean width of sections, then a direct consequence of [11, Theorem 10] is that for any  $k$ -dimensional linear subspace  $F_0 \in G_{n,k}$ ,

$$w(B_\infty^n \cap F_0) \leq \sqrt{\frac{n}{k}} w(B_\infty^k),$$

and this estimate is sharp when  $k \mid n$ .

Furthermore, it was proved in [40] that among all centrally symmetric convex bodies in Löwner's position in  $\mathbb{R}^n$ ,  $w(K)$  is minimized when  $K = B_1^n$ . Finally, in [41], it was proved that among all convex bodies in Löwner's position in  $\mathbb{R}^n$ ,  $w(K)$  is minimized if  $K = \tilde{S}_n$ . We will prove the following results on the mean width of sections of convex bodies in John's position:

**Theorem 1.3.** Let  $K \subseteq \mathbb{R}^n$  be a convex body in John's position and  $F_0 \in G_{n,k}$ . Then

$$w(K \cap F_0) \leq C \frac{n}{k} \sqrt{\frac{\log n}{\log k}} w(S_k),$$

where  $C$  is an absolute constant. Furthermore, if  $K$  is centrally symmetric, then

$$w(K \cap F_0) \leq \sqrt{\frac{n}{k}} w(B_\infty^k).$$

We shall also prove the following result on the mean width of projections of convex bodies in Löwner's position:

**Theorem 1.4.** Let  $K \subseteq \mathbb{R}^n$  be a convex body in Löwner's position. Then, for any  $k$ -dimensional linear subspace  $F_0 \in G_{n,k}$ ,

$$w(P_{F_0}(K)) \geq \sqrt{\frac{k}{n}} w(\tilde{S}_k).$$

Furthermore, if  $K$  is centrally symmetric, then

$$w(P_{F_0}(K)) \geq \sqrt{\frac{k}{n}} w(B_1^k).$$

**Remark.** The proofs of the latter two theorems follow the idea of the previously known results, by applying the Brascamp–Lieb inequality or its reverse form on a linear subspace of  $\mathbb{R}^n$  or  $\mathbb{R}^{n+1}$ . However, we were not able to handle the technical problems, arised from projecting a decomposition of the identity instead of an orthonormal basis, in the non-symmetric case in Theorem 1.3 and a different approach was considered.

### 1.4 The Wills functional

For any compact convex set  $K \subseteq \mathbb{R}^n$ , by Steiner's formula (see [42, Equation (4.1)]), the volume of  $K + tB_2^n$  can be expressed as a polynomial in the variable  $t$

$$|K + tB_2^n| = \sum_{i=0}^n \binom{n}{i} W_i(K) t^i, \quad \forall t \geq 0,$$

where the numbers  $W_i(K)$  are the so-called quermassintegrals of  $K$ . We have that  $W_0(K) = |K|$  is the volume of  $K$ ,  $nW_1(K) = |\partial K|$  is the surface area of  $K$ , and  $W_{n-1} = |B_2^n|w(K)$  is a multiple of the mean width of  $K$ . If  $K$  is contained in a  $k$ -dimensional subspace  $F_0 \in G_{n,k}$ , we can compute its quermassintegrals in  $\mathbb{R}^n$ , but also its quermassintegrals with respect to the subspace  $F_0$ , which we identify with  $\mathbb{R}^k$ . If we denote these quermassintegrals by  $W_i^{(k)}(K)$ , for  $i = 0, \dots, k$ , we have that (see e.g., [39, Property 3.1])

$$W_i^{(k)}(K) = \frac{\binom{n}{i}}{\binom{k}{i}} \frac{|B_2^i|}{|B_2^{n-k+i}|} W_{n-k+i}(K), \quad \forall 0 \leq i \leq k,$$

while  $W_i(K) = 0$  for all  $0 \leq i < n - k$ . In order to avoid the issue that quermassintegrals depend on the space where the convex body is embedded, McMullen [34] defined the intrinsic volumes of a compact convex set  $K \subseteq \mathbb{R}^n$  as

$$V_i(K) = \frac{\binom{n}{i}}{|B_2^{n-i}|} W_{n-i}(K), \quad \forall 0 \leq i \leq n.$$

In [45], Wills introduced the functional

$$\mathcal{W}(K) = \sum_{i=0}^n V_i(K). \tag{1.4}$$

He studied it because of its possible relation with the so-called lattice-point enumerator  $G(K) = \#(K \cap \mathbb{Z}^n)$ . It was proved in [2] that, among symmetric convex bodies in John's position,  $\mathcal{W}(K)$  is maximized if  $K = B_\infty^n$ . Here, we prove the following:

**Theorem 1.5.** Let  $K \subseteq \mathbb{R}^n$  be a centrally symmetric convex body in John's position. Then, for any  $F_0 \in G_{n,k}$  and every  $\lambda \geq 0$ ,

$$\mathcal{W}(\lambda(K \cap F_0)) \leq \mathcal{W}\left(\lambda \sqrt{\frac{n}{k}} B_\infty^k\right).$$



**Remark.** The proof of this Theorem follows the idea of the proof of the result in [2]. What is new here is the consideration of dilations of  $K \cap F_0$  and  $\sqrt{\frac{n}{k}}B_\infty^k$ . This is indeed something different since the Wills functional is not homogeneous. Considering the dilations is important for the applications. Indeed, as direct consequences of obtaining Theorem 1.5 for such dilations, we can obtain the symmetric cases of Theorem 1.1 and Theorem 1.3, providing a different proof in those cases.

We also prove the following estimate for the Wills functional of projections of convex bodies in Löwner's position:

**Theorem 1.6.** Let  $K \subseteq \mathbb{R}^n$  be a centrally symmetric convex body in Löwner's position. Then, for any  $F_0 \in G_{n,k}$ ,

$$\mathcal{W}(P_{F_0}(K)) \geq \frac{1}{k^{k/2}}.$$

**1.5 Sections of convex bodies in minimal surface area position**

The main tool used to obtain most of the estimates above is the fact that a decomposition of the identity operator is associated to any convex body in John's position and that this decomposition allows the use of the Brascamp–Lieb inequality (see Theorem 2.1 below). When  $K$  is a polytope in minimal surface area, then there is again a decomposition of the identity associated to  $K$  (see Section 2.7). A similar use of the Brascamp–Lieb inequality, together with an approximation by polytopes, will lead to similar estimates for sections of convex bodies in minimal surface area position. Namely, we can prove the following:

**Theorem 1.7.** Let  $K \subseteq \mathbb{R}^n$  be a convex body in minimal surface area position and let  $\Pi K$  and  $\Pi^*K$  denote its projection body and polar projection body, respectively. Then, for any  $k$ -dimensional linear subspace  $F_0 \in G_{n,k}$ , we have

- (a)  $|\Pi^*K \cap F_0| \leq \frac{4^k n^k}{k!} \frac{1}{|\partial K|^k},$
- (b)  $|P_{F_0}(\Pi K)| \geq \left(\frac{|\partial K|}{n}\right)^k.$

Furthermore, if  $K$  is centrally symmetric, then for any  $k$ -dimensional linear subspace  $F_0 \in G_{n,k}$ , we have

- (i)  $\mathcal{W}(K \cap F_0) \leq \mathcal{W}\left(\frac{n^2}{k} \frac{|K|}{|\partial K|} B_\infty^k\right),$
- (ii)  $|K \cap F_0|^{1/k} \leq \frac{n^2}{k} \frac{|K|}{|\partial K|} |B_\infty^k|^{1/k},$
- (iii)  $w(K \cap F_0) \leq \frac{n^2}{k} \frac{|K|}{|\partial K|} w(B_\infty^k),$
- (iv)  $|(K \cap F_0)^\circ|^{1/k} \geq \frac{k}{n^2} \frac{|\partial K|}{|K|} |(B_\infty^k)^\circ|^{1/k},$
- (v)  $w((K \cap F_0)^\circ) \geq \frac{k}{n^2} \frac{|\partial K|}{|K|} w((B_\infty^k)^\circ).$

**Remark.** Notice that if  $k = n$  then (a) recovers the right-hand side of (2.6), (b) recovers the left-hand side of (2.7), (ii) recovers the estimate given by Ball's reverse isoperimetric inequality in [6], and (iii) recovers the estimate given in [33, Theorem 7.1].

## 2 Preliminaries

### 2.1 John's position

As mentioned in the introduction, a convex body is said to be in John's position if the maximal volume ellipsoid contained in it is the Euclidean unit ball. A classical theorem of John [29] (see also [7]) states that  $K$  is in John's position if and only if  $B_2^n \subseteq K$  and there exist  $m = O(n^2)$  contact points  $\{u_j\}_{j=1}^m \subseteq \partial K \cap S^{n-1}$  (the intersection of the boundary of  $K$  and the Euclidean unit sphere) and  $\{c_j\}_{j=1}^m$  with  $c_j > 0$  for every  $1 \leq j \leq m$ , such that

$$I_n = \sum_{j=1}^m c_j u_j \otimes u_j, \quad \sum_{j=1}^m c_j u_j = 0, \quad \text{and} \quad \sum_{j=1}^m c_j = n. \quad (2.1)$$

Here  $I_n$  denotes the identity operator in  $\mathbb{R}^n$ ,  $u_j \otimes u_j(x) = \langle x, u_j \rangle u_j$  for every  $x \in \mathbb{R}^n$ , and the third inequality is obtained from the first one by taking traces.

Notice that, for any such decomposition of the identity, we have that for every  $1 \leq k \leq m$

$$1 = |u_k|^2 = \sum_{j=1}^m c_j \langle u_k, u_j \rangle^2 \geq c_k \langle u_k, u_k \rangle^2 = c_k.$$

Thus, all the numbers  $(c_j)_{j=1}^m$  are in the interval  $(0, 1]$ .

### 2.2 Brascamp–Lieb inequality

We will make use of the Brascamp–Lieb inequality (see [13]) and the reverse Brascamp–Lieb inequality due to Barthe (see [9]) in the following form, obtained by Ball (see [5]):

**Theorem 2.1.** Let  $m \geq n$ ,  $\{u_j\}_{j=1}^m \subseteq S^{n-1}$ , and  $\{c_j\}_{j=1}^m \subseteq (0, 1]$  be such that  $I_n = \sum_{j=1}^m c_j u_j \otimes u_j$ . Then, for any integrable functions  $\{f_j\}_{j=1}^m : \mathbb{R} \rightarrow [0, \infty)$ , we have that

$$\int_{\mathbb{R}^n} \prod_{j=1}^m f_j^{c_j}(\langle x, u_j \rangle) dx \leq \prod_{j=1}^m \left( \int_{\mathbb{R}} f_j(t) dt \right)^{c_j} \quad (\text{Brascamp–Lieb inequality}).$$

Besides, for any integrable functions  $\{h_j\}_{j=1}^m : \mathbb{R} \rightarrow [0, \infty)$  and  $h : \mathbb{R}^n \rightarrow [0, \infty)$  satisfying

$$h\left(\sum_{j=1}^m \theta_j c_j u_j\right) \geq \prod_{j=1}^m h_j^{c_j}(\theta_j) \quad \text{for every } \{\theta_j\}_{j=1}^m \subseteq \mathbb{R},$$

we have that

$$\int_{\mathbb{R}^n} h(x) dx \geq \prod_{j=1}^m \left(\int_{\mathbb{R}} h_j(t) dt\right)^{c_j} \quad (\text{Reverse Brascamp-Lieb inequality}).$$

### 2.3 The regular simplex

Let  $\Delta_k$  denote the  $k$ -dimensional regular simplex

$$\Delta_k = \text{conv}\{e_1, \dots, e_{k+1}\} \subseteq H_0,$$

where  $H_0 = \{x \in \mathbb{R}^{k+1} : \sum_{i=1}^{k+1} x_i = 1\}$  is identified with  $\mathbb{R}^k$  and  $(\frac{1}{k+1}, \dots, \frac{1}{k+1})$  is identified with the origin. It is well known that

- $|\Delta_k| = \frac{\sqrt{k+1}}{k!},$
- $r(\Delta_k) = \frac{1}{\sqrt{k(k+1)}},$
- $R(\Delta_k) = \sqrt{\frac{k}{k+1}},$
- $\Delta_k^\circ = -(k+1)\Delta_k.$
- $w(\Delta_k) \simeq \sqrt{\frac{\log k}{k}},$

where  $a \simeq b$  denotes the fact that there exist two positive absolute constants  $c_1, c_2$  such that  $c_1 a \leq b \leq c_2 a$ . Thus,  $\frac{1}{r(\Delta_k)} \Delta_k$  is in John's position and  $\frac{1}{R(\Delta_k)} \Delta_k$  is in Löwner's position. Then, if  $S_k$  denotes the  $k$ -dimensional simplex in John's position and  $\tilde{S}_k$  denotes the  $k$ -dimensional simplex in Löwner's position, we have that

$$S_k = \sqrt{k(k+1)} \Delta_k \quad \text{and} \quad \tilde{S}_k = \sqrt{\frac{k+1}{k}} \Delta_k.$$

Therefore,

$$|S_k|^{1/k} = \frac{\sqrt{k(k+1)^{1+\frac{1}{k}}}}{(k!)^{1/k}} \quad \text{and} \quad |\tilde{S}_k|^{1/k} = \frac{1}{(k!)^{1/k}} \sqrt{\frac{(k+1)^{1+\frac{1}{k}}}{k}}.$$

Moreover,

$$w(S_k) \simeq \sqrt{k \log k} \quad \text{and} \quad w(\tilde{S}_k) \simeq \sqrt{\frac{\log k}{k}}.$$

## 2.4 Mean width

Let  $K \subseteq \mathbb{R}^n$  be a convex body. The mean width of  $K$  is defined by

$$w(K) = \int_{S^{n-1}} h_K(\theta) d\sigma(\theta),$$

where, for every  $\theta \in S^{n-1}$ ,  $h_K(\theta)$  is the support function of  $K$  at  $\theta$  and  $d\sigma$  denotes the uniform probability measure on  $S^{n-1}$ . If we also assume that  $K$  contains the origin in its interior, then  $h_K$  is homogeneous of degree 1. There is a nice representation of the mean width in terms of the standard Gaussian random vector  $G$  in  $\mathbb{R}^n$  (see, for instance, [3, Proof of Theorem 4.2.2]):

$$\mathbb{E}h_K(G) = c_n w(K), \tag{2.2}$$

where  $c_n = \frac{n|B_2^n|\Gamma(\frac{n+1}{2})}{\sqrt{2\pi}^{n/2}} = \frac{\sqrt{2}\Gamma(\frac{n+1}{2})}{\Gamma(\frac{n}{2})}$ . Indeed, integrating in polar coordinates, one has

$$\begin{aligned} \mathbb{E}h_K(G) &= \int_{\mathbb{R}^n} h_K(x) \frac{e^{-\frac{\|x\|_2^2}{2}}}{(2\pi)^{n/2}} dx = n|B_2^n| \int_0^\infty r^n \frac{e^{-\frac{r^2}{2}}}{(2\pi)^{n/2}} \int_{S^{n-1}} h_K(\theta) d\sigma(\theta) \\ &= c_n \int_{S^{n-1}} h_K(\theta) d\sigma(\theta) = c_n w(K). \end{aligned}$$

Likewise, since for any convex body containing the origin in its interior the support function of  $K^\circ$  is  $h_{K^\circ} = \|\cdot\|_K$ , where  $\|\cdot\|_K$  is the Minkowski gauge function of  $K$ , given by

$$\|x\|_K := \inf\{\lambda > 0 : x \in \lambda K\}$$

for all  $x \in \mathbb{R}^n$ , we have that if  $G$  is a standard Gaussian random vector in  $\mathbb{R}^n$

$$\mathbb{E}\|G\|_K = c_n w(K^\circ). \tag{2.3}$$

We would like to refer the reader to [11], [17], [18], or [47] for more information on the use of the Gaussian measure of sections of convex bodies.

## 2.5 Log-concave functions

A function  $f: \mathbb{R}^n \rightarrow [0, \infty)$  is called log-concave if  $f(x) = e^{-v(x)}$  where  $v: \mathbb{R}^n \rightarrow (-\infty, \infty]$  is a convex function. It is well known that any integrable log-concave function  $f: \mathbb{R}^n \rightarrow [0, \infty)$  is bounded and has moments of all orders. If  $K \subseteq \mathbb{R}^n$  is a convex body then its indicator function  $\chi_K$  is integrable and log-concave with integral  $|K|$ . If additionally  $K$  is a convex body containing the origin, then  $e^{-\|\cdot\|_K}$  is integrable and log-concave with integral  $n!|K|$ .

Given a log-concave function  $f = e^{-v}$ , where  $v: \mathbb{R}^n \rightarrow (-\infty, \infty]$  is a convex function, its polar function is the function  $f^\circ: \mathbb{R}^n \rightarrow [0, \infty)$  given by

$$f^\circ(x) = e^{-\mathcal{L}(v)(x)},$$

where  $\mathcal{L}(v)$  denotes the Legendre transform

$$\mathcal{L}(v)(x) = \sup_{y \in \mathbb{R}^n} (\langle x, y \rangle - v(y)), \quad x \in \mathbb{R}^n.$$

For more information on log-concave functions, we refer the reader to [14, Chapter 2].

## 2.6 The Wills functional

Let us recall that for any  $n$ -dimensional convex body  $K$ , its Wills functional is defined by

$$\mathcal{W}(K) = \sum_{i=0}^n V_i(K),$$

where  $V_i(K)$  denotes the  $i$ -th intrinsic volume of  $K$ . Many properties of the Wills functional can be found in [46], [25], [35], or [2]. For our purposes, we emphasize the following two:

- (1) (Hadwiger, see [25, (1.3)]) For any convex body  $K \subseteq \mathbb{R}^n$ ,

$$\mathcal{W}(K) = \int_{\mathbb{R}^n} e^{-\pi d(x,K)^2} dx,$$

where  $d(x, K)$  denotes the Euclidean distance from  $x$  to  $K$ .

- (2) (Hadwiger, see [25, (2.3)]) If  $E$  is a linear subspace of  $\mathbb{R}^n$ ,  $K_1 \subseteq E$  and  $K_2 \subseteq E^\perp$ , then

$$\mathcal{W}(K_1 \times K_2) = \mathcal{W}(K_1)\mathcal{W}(K_2).$$

In particular, if  $K = [-a, a] \subseteq \mathbb{R}$ , we have that

$$\mathcal{W}([-a, a]) = 2a + 2 \int_a^\infty e^{-\pi(x-a)^2} dx = 2a + 1$$

and if  $K = aB_\infty^n \subseteq \mathbb{R}^n$  then  $\mathcal{W}(aB_\infty^n) = (1 + 2a)^n$ .

Let us point out that for any  $\lambda > 0$

$$\mathcal{W}(\lambda K) = \sum_{i=0}^n V_i(\lambda K) = 1 + \sum_{i=1}^n \lambda^i V_i(K).$$

Therefore, if two convex bodies  $K, L \subseteq \mathbb{R}^n$  verify that  $\mathcal{W}(\lambda K) \leq \mathcal{W}(\lambda L)$  for every  $\lambda \geq 0$ , then one immediately obtains that  $V_n(K) \leq V_n(L)$  and  $V_1(K) \leq V_1(L)$  or, equivalently,  $|K| \leq |L|$  and  $w(K) \leq w(L)$ .

Notice that, for any convex body  $K \subseteq \mathbb{R}^n$ , the function given by  $d(x, K)$  for every  $x \in \mathbb{R}^n$  is convex on  $\mathbb{R}^n$  (see [42, Lemma 1.5.9]) and, as the square function is convex on  $\mathbb{R}$ ,  $d(x, K)^2$  is convex on  $\mathbb{R}^n$ . Therefore, the first property above shows that, for any convex body  $K \subseteq \mathbb{R}^n$ , its Will's functional is the integral of the log-concave function  $f_K : \mathbb{R}^n \rightarrow [0, \infty)$  given by

$$f_K(x) = e^{-\pi d(x, K)^2}.$$

Using a double polarity (both in the convex body and in the family of log-concave functions), for any convex body  $K \subseteq \mathbb{R}^n$  containing the origin in its interior, we define the log-concave function  $f_K^\circ$ . It was proved in [2, Lemma 3.1] that for every  $x \in \mathbb{R}^n$

$$f_K^\circ(x) = e^{-\frac{\|x\|_2^2}{4\pi} - \|x\|_K}. \tag{2.4}$$

The following lemma shows that if, for every  $\lambda \geq 0$ , the integral of  $f_{(\lambda K)^\circ}(x)$  is bounded by the integral of  $f_{(\lambda L)^\circ}(x)$ , then  $|K| \leq |L|$ .

**Lemma 2.1.** Let  $K, L \subseteq \mathbb{R}^n$  be two convex bodies containing the origin in their interiors. Assume that there exist two numbers  $A$  and  $\lambda_0 > 0$  such that, for any  $\lambda \in (0, \lambda_0)$ ,

$$\int_{\mathbb{R}^n} f_{(\lambda K)^\circ}^\circ(x) dx \leq A \int_{\mathbb{R}^n} f_{(\lambda L)^\circ}^\circ(x) dx.$$

Then  $|K| \leq A|L|$ .

**Proof.** Notice that for any convex body  $K \subseteq \mathbb{R}^n$  containing the origin in its interior and any  $\lambda > 0$ ,

$$\begin{aligned} \int_{\mathbb{R}^n} f_{(\lambda K)^\circ}^\circ(x) dx &= \int_{\mathbb{R}^n} e^{-\frac{\|x\|_2^2}{4\pi}} e^{-\|x\|_{\lambda K}} dx = \int_{\mathbb{R}^n} e^{-\frac{\|x\|_2^2}{4\pi}} e^{-\frac{\|x\|_K}{\lambda}} dx \\ &= \lambda^n \int_{\mathbb{R}^n} e^{-\frac{\lambda^2 \|x\|_2^2}{4\pi}} e^{-\|x\|_K} dx. \end{aligned}$$

Therefore, we have that for every  $\lambda \in (0, \lambda_0)$ ,

$$\int_{\mathbb{R}^n} e^{-\frac{\lambda^2 \|x\|_2^2}{4\pi}} e^{-\|x\|_K} dx \leq A \int_{\mathbb{R}^n} e^{-\frac{\lambda^2 \|x\|_2^2}{4\pi}} e^{-\|x\|_L} dx$$

and, taking the limit as  $\lambda$  tends to 0 we obtain that

$$n! |K| = \int_{\mathbb{R}^n} e^{-\|x\|_K} dx \leq A \int_{\mathbb{R}^n} e^{-\|x\|_L} dx = n! A|L|.$$

■

### 2.7 Convex bodies in minimal surface area position

A convex body  $K \subseteq \mathbb{R}^n$  is said to be in minimal surface area position if it has minimal surface area among all of its volume preserving affine images. That is, if

$$|\partial K| = \min \{ |\partial T(K)| : T \in SL(n) \},$$

where  $SL(n)$  denotes the set of non-degenerate linear maps  $T \in GL(n)$  with  $|\det T| = 1$ . The surface area measure of a convex body  $K$  is the measure on the sphere defined by

$$\sigma_K(A) := \nu(\{x \in \partial K : \nu_K(x) \in A\}) \quad \forall A \text{ Borel set in } S^{n-1},$$

where  $\nu$  denotes the Hausdorff measure on  $\partial K$  and  $\nu_K(x)$  is the outer normal vector to  $K$  at  $x$ , which is defined  $\nu$ -almost everywhere.

The projection body  $\Pi K$  and its polar, the polar projection body  $\Pi^*K$ , of a convex body  $K$  are the centrally symmetric convex bodies defined by

$$h_{\Pi K}(x) = \|x\|_{\Pi^*K} = |x| |P_{x^\perp}(K)| = \frac{1}{2} \int_{S^{n-1}} |\langle x, \theta \rangle| d\sigma_K(\theta),$$

where, for any  $x \neq 0$ ,  $|P_{x^\perp}(K)|$  denotes the  $(n-1)$ -dimensional volume of the projection of  $K$  onto the hyperplane orthogonal to  $x$  and the last equality is the well-known Cauchy's formula (see, for instance, [42, Equation (5.80)]).

It was proved by Petty [36] (see also [21]) that  $K$  is in minimal surface area position if and only if  $\sigma_K$  is isotropic, that is, if

$$I_n = \frac{n}{|\partial K|} \int_{S^{n-1}} u \otimes u d\sigma_K(u).$$

In [22], it was observed that the latter happens if and only if  $\Pi K$  is in minimal mean width position, that is,

$$w(\Pi K) = \min\{w(T(\Pi K)) : T \in SL(n)\}.$$

Notice that if  $K$  is a polytope with facets  $\{F_j\}_{j=1}^m$  with outer normal vectors  $\{u_j\}_{j=1}^m$ , then the surface area measure of  $K$  is

$$\sigma_K = \sum_{j=1}^m |F_j| \delta_{u_j},$$

where  $\delta_j$  denotes the Dirac delta measure on  $u_j$ . Moreover,  $K$  is in minimal surface area position if and only if

$$I_n = \sum_{j=1}^m \frac{n|F_j|}{|\partial K|} u_j \otimes u_j.$$

In particular, if  $K$  is a polytope with facets  $\{F_j\}_{j=1}^m$  and outer normal vectors  $\{u_j\}_{j=1}^m$ , then for every  $x \in \mathbb{R}^n$

$$h_{\Pi K}(x) = \|x\|_{\Pi^*K} = \frac{1}{2} \sum_{j=1}^m |F_j| |\langle x, u_j \rangle|. \quad (2.5)$$



It was proved in [21] that, as a consequence of a lemma obtained from the Brascamp–Lieb inequality (see [6]), if  $K$  is a convex body in minimal surface area position then

$$|B_2^n| \left( \frac{n|B_2^n|}{|B_2^{n-1}|} \right)^n \frac{1}{|\partial K|^n} \leq |\Pi^* K| \leq \frac{4^n n^n}{n!} \frac{1}{|\partial K|^n}. \tag{2.6}$$

Moreover, if  $K$  is a convex body in minimal surface area position, then

$$\left( \frac{|\partial K|}{n} \right)^n \leq |\Pi K| \leq |B_2^n| \left( \frac{|B_2^{n-1}|}{n|B_2^n|} \right)^n |\partial K|^n. \tag{2.7}$$

This can be seen as a consequence of the Blaschke–Santaló inequality and its exact reverse for zonoids (see [23] and [38]), or as a direct consequence of the reverse form of Brascamp–Lieb inequality (see [22]).

### 3 General Setting

In this section, we introduce the notation for a setting that will be used in several of our proofs. We distinguish the cases in which we are dealing with non-symmetric convex bodies in John’s position, symmetric convex bodies in John’s position, or polytopes in minimal surface area position.

#### 3.1 Non-symmetric convex bodies in John’s position

Let  $K \subseteq \mathbb{R}^n$  be a (not necessarily symmetric) convex body in John’s position and let  $\{u_j\}_{j=1}^m$  and  $\{c_j\}_{j=1}^m$  be the contact points in  $\partial K \cap S^{n-1}$  and positive weights satisfying John’s condition (2.1). We will denote by  $C \subseteq \mathbb{R}^n$  the convex body

$$C = \{x \in \mathbb{R}^n : \langle x, u_j \rangle \leq 1, \forall 1 \leq j \leq m\}. \tag{3.1}$$

It is easily verified that  $K \subseteq C$ . We will denote, for every  $1 \leq j \leq m$ ,

- $v_j = \sqrt{\frac{n}{n+1}}(-u_j, \frac{1}{\sqrt{n}}) \in S^n$ , and
- $\delta_j = \frac{n+1}{n}c_j$ .

These vectors satisfy

$$I_{n+1} = \sum_{j=1}^m \delta_j v_j \otimes v_j, \quad \sum_{j=1}^m \delta_j v_j = (0, \sqrt{n+1}) \quad \text{and} \quad \sum_{j=1}^m \delta_j = n+1. \tag{3.2}$$

Therefore, as seen in Section 2.1,  $\delta_j \in (0, 1]$  for every  $1 \leq j \leq m$ . Introduce the cone

$$L := \{y = (x, r) \in \mathbb{R}^{n+1} : \langle y, v_j \rangle \geq 0, \forall 1 \leq j \leq m\}. \quad (3.3)$$

The next lemma, which was proved in [6], relates  $L$  and  $C$ . We include its proof here for the sake of completeness.

**Lemma 3.1.** Let  $K \subseteq \mathbb{R}^n$  be a convex body in John's position and let  $L$  be defined as in equation (3.3). Then

$$L = \left\{ (x, r) \in \mathbb{R}^{n+1} : r \geq 0, x \in \frac{r}{\sqrt{n}}C \right\}.$$

**Proof.** Let  $y = (x, r) \in L$ . By the definition of  $v_j$  we have that for each  $1 \leq j \leq m$

$$\langle y, v_j \rangle = -\sqrt{\frac{n}{n+1}} \langle x, u_j \rangle + \frac{r}{\sqrt{n+1}}.$$

Assume that  $r < 0$ . Then, since  $\langle y, v_j \rangle \geq 0$  for every  $1 \leq j \leq m$ , we have that

$$-\sqrt{\frac{n}{n+1}} \langle x, u_j \rangle + \frac{r}{\sqrt{n+1}} \geq 0 \quad \forall 1 \leq j \leq m.$$

Then,  $\langle x, u_j \rangle < 0$  for every  $1 \leq j \leq m$ . As a consequence, since  $\{c_j\}_{j=1}^m \subseteq (0, \infty)$ ,

$$\sum_{j=1}^m c_j \langle x, u_j \rangle < 0,$$

which contradicts the fact that  $\sum_{j=1}^m c_j u_j = 0$ . Therefore, if  $y = (x, r) \in L$  then  $r \geq 0$ .

For any  $r \geq 0$  and every  $1 \leq j \leq m$  we have  $\langle y, v_j \rangle \geq 0$  if and only if  $\langle x, u_j \rangle \leq \frac{r}{\sqrt{n}}$ . The latter condition is true for every  $1 \leq j \leq m$  if and only if  $x \in \frac{r}{\sqrt{n}}C$ .

Conversely, assume that  $y = (x, r)$  verifies that  $r \geq 0$  and  $x \in \frac{r}{\sqrt{n}}C$ , which happens if and only if  $\langle x, u_j \rangle \leq \frac{r}{\sqrt{n}}$  for every  $1 \leq j \leq m$ . Then, for every  $1 \leq j \leq m$ ,

$$\langle y, v_j \rangle = -\sqrt{\frac{n}{n+1}} \langle x, u_j \rangle + \frac{r}{\sqrt{n+1}} \geq 0.$$

Thus,  $y \in L$ . ■

Given any  $k$ -dimensional affine subspace  $F_h$  in  $\mathbb{R}^n$  at distance  $h$  from the origin, we will consider the linear  $(k + 1)$ -dimensional subspace in  $\mathbb{R}^{n+1}$

$$H = \text{span}\{(x, \sqrt{n}) : x \in F_h\} \in G_{n+1,k+1}. \tag{3.4}$$

Notice that if  $F_0 \in G_{n,k}$  is a linear subspace then  $H$  equals the cartesian product  $H = F_0 \times \mathbb{R} = \{(x, r) \in \mathbb{R}^{n+1} : x \in F_0, r \in \mathbb{R}\}$ . Furthermore, assume that  $F_h$  is at distance  $h$  from the origin and  $f : L \cap H \rightarrow [0, \infty)$  is an integrable function. By Lemma 3.1, and taking into account that  $\mathbb{R}^n \times \{0\}$  and  $P_H(\{0\} \times \mathbb{R})$  provide an orthogonal decomposition of  $H$ , we have that

$$\int_{L \cap H} f(x, r) dr dx = \int_0^\infty \int_{\frac{r}{\sqrt{n}}(C \cap F_h) \times \{r\}} f(x, r) dx \sqrt{\frac{n + h^2}{n}} dr. \tag{3.5}$$

Set  $J = \{1 \leq j \leq m : P_H v_j \neq 0\}$  and, for every  $j \in J$ , we define

- $w_j = \frac{P_H v_j}{\|P_H v_j\|_2}$ ,
- $\kappa_j = \frac{n+1}{n} c_j \|P_H v_j\|_2^2 = \delta_j \|P_H v_j\|_2^2$ .

Then, we have that

$$I_H = \sum_{j \in J} \kappa_j w_j \otimes w_j \quad \text{and} \quad \sum_{j \in J} \kappa_j = k + 1, \tag{3.6}$$

where  $I_H$  denotes the identity in the linear subspace  $H$ . Furthermore, denoting by  $s_j = \frac{1}{\|P_H v_j\|_2}$  for every  $j \in J$ , one has that for every  $y = (x, r) \in H \subseteq \mathbb{R}^{n+1}$

$$\begin{aligned} \sum_{j \in J} \kappa_j s_j \langle y, w_j \rangle &= \sum_{j \in J} \delta_j \langle y, P_H v_j \rangle = \sum_{j=1}^m \delta_j \langle y, P_H v_j \rangle = \sum_{j=1}^m \delta_j \langle y, v_j \rangle \\ &= r \sqrt{n + 1}. \end{aligned} \tag{3.7}$$

The following lemma shows that, whenever  $F_0$  is a linear subspace, we have a strictly positive lower bound for the Euclidean norm of  $P_H v_j$  for every  $1 \leq j \leq m$ . Consequently, if  $F_0 \in G_{n,k}$ , the set  $J$  defined above equals  $J = \{1, \dots, m\}$ .

**Lemma 3.2.** Let  $\{u_j\}_{j=1}^m \subseteq S^{n-1}$ ,  $\{c_j\}_{j=1}^m$  be such that (2.1) holds,  $F_0 \in G_{n,k}$ ,  $H = F_0 \times \mathbb{R} \in G_{n+1,k+1}$ , and  $\{v_j\}_{j=1}^m \subseteq S^n$  be defined as in (3.2). Then, for every  $1 \leq j \leq m$ , we have

$$\frac{1}{n + 1} \leq \|P_H v_j\|_2^2 \leq 1.$$

**Proof.** Let  $c = \left(0, \frac{1}{\sqrt{n+1}}\right) \in H$  and notice that for every  $1 \leq j \leq m$

$$\langle P_H(v_j - c), c \rangle = \langle v_j - c, c \rangle = \frac{1}{n+1} - \frac{1}{n+1} = 0.$$

Since  $c \in H$ , we have that  $P_H c = c$  and then  $P_H v_j = c + P_H(v_j - c)$ . Thus,

$$\|P_H v_j\|_2^2 = \|c + (P_H(v_j - c))\|_2^2 = \|c\|_2^2 + \|P_H(v_j - c)\|_2^2 \geq \|c\|_2^2 = \frac{1}{n+1}.$$

Thus, for every  $1 \leq j \leq m$ , we have that

$$\frac{1}{n+1} \leq \|P_H v_j\|_2^2 \leq 1.$$

■

### 3.2 Symmetric convex bodies in John's position

Let  $K \subseteq \mathbb{R}^n$  be a centrally symmetric convex body in John's position and, like in the not necessarily symmetric case, let  $\{u_j\}_{j=1}^m$  and  $\{c_j\}_{j=1}^m$  be the contact points of  $\partial K$  and  $S^{n-1}$  and positive weights satisfying (2.1). We will also denote by  $C_0$  the symmetric convex body

$$C_0 = \{x \in \mathbb{R}^n : |\langle x, u_j \rangle| \leq 1, \forall 1 \leq j \leq m\}. \quad (3.8)$$

Clearly,  $K$  is a subset of  $C_0$ . If  $F_0 \in G_{n,k}$  is a linear subspace, we set  $J_0 = \{1 \leq j \leq m : P_{F_0} u_j \neq 0\}$  and for every  $j \in J_0$ , we define

- $v_j^0 = \frac{P_{F_0} u_j}{\|P_{F_0} u_j\|_2} \in S^{n-1} \cap F_0$ ,
- $\delta_j^0 = c_j \|P_{F_0} u_j\|_2^2$ .

Then, we have that

$$I_{F_0} = \sum_{j \in J_0} \delta_j^0 v_j^0 \otimes v_j^0 \quad \text{and} \quad \sum_{j \in J_0} \delta_j^0 = k, \quad (3.9)$$

where  $I_{F_0}$  denotes the identity operator in  $F_0$ , and also

$$\begin{aligned}
 K \cap F_0 \subseteq C_0 \cap F_0 &= \{x \in F_0 : |\langle x, u_j \rangle| \leq 1, \forall 1 \leq j \leq m\} \\
 &= \{x \in F_0 : |\langle x, P_{F_0} u_j \rangle| \leq 1, \forall 1 \leq j \leq m\} \\
 &= \{x \in F_0 : |\langle x, P_{F_0} u_j \rangle| \leq 1, \forall j \in J_0\} \\
 &= \left\{ x \in F_0 : |\langle x, v_j^0 \rangle| \leq t_j, \forall j \in J_0 \right\},
 \end{aligned}$$

where  $t_j = \frac{1}{\|P_{F_0} u_j\|_2} = \left(\frac{c_j}{\delta_j^0}\right)^{1/2}$  for every  $j \in J_0$ . Furthermore,

$$\begin{aligned}
 (K \cap F_0)^\circ &\supseteq (C_0 \cap F_0)^\circ = P_{F_0}(C_0^\circ) = P_{F_0}(\text{conv}\{\pm u_j : 1 \leq j \leq m\}) \\
 &= \text{conv}\{\pm P_{F_0} u_j : 1 \leq j \leq m\} \\
 &= \text{conv}\{\pm P_{F_0} u_j : j \in J_0\}.
 \end{aligned}$$

Thus,

$$K \cap F_0 \subseteq C_0 \cap F_0 = \{x \in F_0 : |\langle x, v_j^0 \rangle| \leq t_j, \forall j \in J_0\} \quad (3.10)$$

and

$$(K \cap F_0)^\circ \supseteq (C_0 \cap F_0)^\circ = \text{conv}\{\pm P_{F_0} u_j : j \in J_0\}. \quad (3.11)$$

### 3.3 Polytopes in minimal surface area position

Let  $K$  be a (not necessarily centrally symmetric) polytope in minimal surface area position with facets  $\{F_j\}_{j=1}^m$  and outer normal vectors  $\{u_j\}_{j=1}^m$ , and let  $F_0 \in G_{n,k}$  be a  $k$ -dimensional linear subspace. Then,

$$K = \{x \in \mathbb{R}^n : \langle x, u_j \rangle \leq h_K(u_j), \forall 1 \leq j \leq m\}$$

and

$$I_n = \sum_{j=1}^m \frac{n|F_j|}{|\partial K|} u_j \otimes u_j = \sum_{j=1}^m c_j u_j \otimes u_j, \quad (3.12)$$

where  $c_j = \frac{n|F_j|}{|\partial K|}$  for every  $1 \leq j \leq m$ . Besides (see, for instance, [24, Theorem 18.2])

$$\sum_{j=1}^m c_j u_j = \frac{n}{|\partial K|} \sum_{j=1}^m |F_j| u_j = 0.$$

and

$$\sum_{j=1}^m c_j h_K(u_j) = \sum_{j=1}^m \frac{n|F_j|}{|\partial K|} h_K(u_j) = \frac{n^2|K|}{|\partial K|}. \tag{3.13}$$

Note also that if  $K$  is a centrally symmetric polytope in minimal surface area position, with facets  $\{F_j\}_{j=1}^m$  and outer normal vectors  $\{u_j\}_{j=1}^m$ , and if  $F_0 \in G_{n,k}$  is a  $k$ -dimensional linear subspace, then

$$K = \{x \in \mathbb{R}^n : |\langle x, u_j \rangle| \leq h_K(u_j), \forall 1 \leq j \leq m\}.$$

As in the case where the decomposition of the identity comes from a centrally symmetric convex body in John’s position, we set  $J_0 = \{1 \leq j \leq m : P_{F_0} u_j \neq 0\}$  and, for every  $j \in J_0$ , we define

- $v_j^0 = \frac{P_{F_0} u_j}{\|P_{F_0} u_j\|_2}$ ,
- $\delta_j^0 = c_j \|P_{F_0} u_j\|_2^2 = \frac{n|F_j| \|P_{F_0} u_j\|_2^2}{|\partial K|}$ .

We have that

$$I_{F_0} = \sum_{j=1}^m c_j P_{F_0} u_j \otimes P_{F_0} u_j = \sum_{j \in J_0} \delta_j^0 v_j^0 \otimes v_j^0. \tag{3.14}$$

Besides, if we denote  $t_j = \frac{1}{\|P_{F_0} u_j\|_2} = \left(\frac{c_j}{\delta_j^0}\right)^{1/2}$  for every  $j \in J_0$ , then

$$K \cap F_0 = \{x \in F_0 : |\langle x, v_j^0 \rangle| \leq t_j h_K(u_j), \forall j \in J_0\} \tag{3.15}$$

and

$$\begin{aligned} (K \cap F_0)^\circ &= \text{conv} \left\{ \pm \frac{P_{F_0}(u_j)}{h_K(u_j)} : 1 \leq j \leq m \right\} \\ &= \text{conv} \left\{ \pm \frac{P_{F_0}(u_j)}{h_K(u_j)} : j \in J_0 \right\}. \end{aligned} \tag{3.16}$$

### 4 Volume of Sections of Convex Bodies in John's Position

In this section, we will give the proof of Theorem 1.1.

**Proof of Theorem 1.1.** Let us start with the symmetric case. Assume that  $K$  is a centrally symmetric convex body in John's position and  $F_0 \in G_{n,k}$  is a linear  $k$ -dimensional subspace. We follow the notation in Section 3.2. By (3.10), we have that

$$K \cap F_0 \subseteq C_0 \cap F_0 = \{x \in F_0 : |\langle x, v_j^0 \rangle| \leq t_j, \forall j \in J_0\},$$

where  $t_j = \frac{1}{\|P_{F_0} u_j\|_2} = \left(\frac{c_j}{\delta_j^0}\right)^{1/2}$  for all  $j \in J_0$ . Therefore, by (3.9) and the Brascamp–Lieb inequality (Theorem 2.1),

$$\begin{aligned} |K \cap F_0| &\leq |C_0 \cap F_0| = \int_{F_0} \prod_{j \in J_0} \chi_{[-t_j, t_j]}(\langle x, v_j^0 \rangle) dx = \int_{F_0} \prod_{j \in J_0} \chi_{[-t_j, t_j]}^{\delta_j^0}(\langle x, v_j^0 \rangle) dx \\ &\leq \prod_{j \in J_0} \left( \int_{\mathbb{R}} \chi_{[-t_j, t_j]}(t) dt \right)^{\delta_j^0} = \prod_{j \in J_0} (2t_j)^{\delta_j^0} = 2^k \prod_{j \in J_0} \left( \frac{c_j}{\delta_j^0} \right)^{\frac{\delta_j^0}{2}}. \end{aligned}$$

By the arithmetic-geometric mean inequality and (2.1), we get

$$\prod_{j \in J_0} \left( \frac{c_j}{\delta_j^0} \right)^{\frac{\delta_j^0}{k}} \leq \sum_{j \in J_0} \frac{\delta_j^0}{k} \frac{c_j}{\delta_j^0} = \frac{1}{k} \sum_{j \in J_0} c_j \leq \frac{1}{k} \sum_{j=1}^m c_j = \frac{n}{k}.$$

Hence,

$$|K \cap F_0|^{1/k} \leq 2\sqrt{\frac{n}{k}} = \sqrt{\frac{n}{k}} |B_\infty^k|^{1/k}.$$

Assume now that  $K \subseteq \mathbb{R}^n$  is a (not necessarily symmetric) convex body in John's position and  $F_0 \in G_{n,k}$  a  $k$ -dimensional linear subspace. We follow the notation introduced in Section 3.1. Applying Lemma 3.1, one gets

$$L \cap H = \left\{ (x, r) \in F_0 \times \mathbb{R} : r \geq 0, x \in \frac{r}{\sqrt{n}}(C \cap F_0) \right\}.$$

Denote  $s_j = \frac{1}{\|P_H v_j\|_2}$  as in (3.7). Using the Brascamp–Lieb inequality (Theorem 2.1), we get

$$\begin{aligned} \int_{L \cap H} e^{-\sum_{j=1}^m \kappa_j s_j \langle Y, w_j \rangle} dy &= \int_H \prod_{j=1}^m \left( \chi_{[0, \infty)}(\langle Y, v_j \rangle) e^{-s_j \langle Y, w_j \rangle} \right)^{\kappa_j} dy \\ &= \int_H \prod_{j=1}^m \left( \chi_{[0, \infty)}(\langle Y, P_H v_j \rangle) e^{-s_j \langle Y, w_j \rangle} \right)^{\kappa_j} dy \\ &= \int_H \prod_{j=1}^m \left( \chi_{[0, \infty)}(\langle Y, w_j \rangle) e^{-s_j \langle Y, w_j \rangle} \right)^{\kappa_j} dy \leq \prod_{j=1}^m \left( \int_0^\infty e^{-s_j t} dt \right)^{\kappa_j} \\ &= \prod_{j=1}^m \|P_H v_j\|_2^{\delta_j \|P_H v_j\|_2^2}. \end{aligned}$$

On the other hand, taking into account (3.7), we see that

$$\begin{aligned} \int_{L \cap H} e^{-\sum_{j=1}^m \kappa_j s_j \langle Y, w_j \rangle} dy &= \int_0^\infty \int_{\frac{r}{\sqrt{n}}(C \cap F_0)} e^{-r\sqrt{n+1}} dx dr \\ &= \int_0^\infty \frac{r^k}{n^{\frac{k}{2}}} |C \cap F_0| e^{-r\sqrt{n+1}} dr = \frac{k!}{n^{\frac{k}{2}}(n+1)^{\frac{k+1}{2}}} |C \cap F_0| \\ &= \frac{k^{\frac{k}{2}}(k+1)^{\frac{k+1}{2}}}{n^{\frac{k}{2}}(n+1)^{\frac{k+1}{2}}} \frac{|C \cap F_0|}{|S_k|}. \end{aligned}$$

Let us maximize  $\prod_{j=1}^m \|P_H v_j\|_2^{\delta_j \|P_H v_j\|_2^2}$  under the constraints

- $\frac{1}{n+1} \leq \|P_H v_j\|_2^2 \leq 1 \quad \forall 1 \leq j \leq m,$
- $\sum_{j=1}^m \delta_j \|P_H v_j\|_2^2 = k+1,$
- $\sum_{j=1}^m \delta_j = n+1,$
- $0 \leq \delta_j \leq 1.$

Equivalently, let us maximize  $F(x, \delta) = \frac{1}{2} \sum_{j=1}^m \delta_j x_j \log x_j$  under the constraints

- $\frac{1}{n+1} \leq x_j \leq 1 \quad \forall 1 \leq j \leq m,$
- $\sum_{j=1}^m \delta_j x_j = k+1,$
- $\sum_{j=1}^m \delta_j = n+1,$
- $0 \leq \delta_j \leq 1.$

First notice that the function  $F(x, \delta)$  is continuous on a compact domain  $M$  in  $\mathbb{R}^{2m}$ , which is given by the constraints. Therefore, it attains its maximum. For every  $x = (x_1, \dots, x_m)$



with  $\frac{1}{n+1} \leq x_j \leq 1$  for all  $1 \leq j \leq m$ , let  $F_x(\delta)$  be the function

$$F_x(\delta) = \frac{1}{2} \sum_{j=1}^m \delta_j x_j \log x_j.$$

Notice that  $F_x$  is a convex function. Since the set

$$A = \left\{ \delta \in \mathbb{R}^m : \sum_{j=1}^m \delta_j x_j = k + 1, \sum_{j=1}^m \delta_j = n + 1, 0 \leq \delta_j \leq 1 \forall 1 \leq j \leq m \right\}$$

is a compact convex set,  $F_x$  attains its maximum on some extreme point of  $A$ . These are the points of intersection of the 2-dimensional faces of the cube

$$\{ \delta \in \mathbb{R}^m : 0 \leq \delta_j \leq 1 \forall 1 \leq j \leq m \}$$

with the  $(m - 2)$ -dimensional affine subspace

$$\left\{ \delta \in \mathbb{R}^m : \sum_{j=1}^m \delta_j x_j = k + 1, \sum_{j=1}^m \delta_j = n + 1 \right\}.$$

Therefore, a maximizer of the function  $F_x$  has to be a point of the form

$$\delta_\lambda = (\underbrace{1, 1, \dots, 1}_n, \underbrace{\lambda, 1 - \lambda, 0, \dots, 0}_{m-n-2})$$

for some  $\frac{1}{2} \leq \lambda \leq 1$  (or a permutation of it), such that  $\sum_{j=1}^m \delta_j x_j = k + 1$  is satisfied. For every  $\delta_\lambda$  with  $\frac{1}{2} \leq \lambda \leq 1$ , we will find the maximizer of the function

$$F_{\delta_\lambda}(x) = \frac{1}{2} \sum_{j=1}^m \delta_{\lambda,j} x_j \log x_j$$

on the compact convex set

$$B_\lambda = \left\{ x \in \mathbb{R}^m : \sum_{j=1}^m \delta_{\lambda,j} x_j = k + 1, \frac{1}{n + 1} \leq x_j \leq 1 \forall 1 \leq j \leq m \right\}.$$

If  $\delta_\lambda^*$  is the decreasing rearrangement of  $\delta_\lambda$ , we can assume without loss of generality that  $\delta_\lambda = \delta_\lambda^*$ . Let

$$D = \frac{k+1}{n+1}$$

and

$$\tilde{x} = \left( \underbrace{1, 1, \dots, 1}_k, D, \underbrace{\frac{1}{n+1}, \dots, \frac{1}{n+1}}_{m-k-1} \right).$$

We check that  $\frac{1}{n+1} \leq D \leq 1$  and

$$\sum_{j=1}^m \delta_{\lambda,j} \tilde{x}_j = k + D + \frac{n-k}{n+1} = k + 1.$$

For every  $x = (x_1, \dots, x_m) \in B_\lambda$ , we have  $\tilde{x} \succ (x_1, \dots, x_m)$ , since the first  $k+1$  coordinates of  $\tilde{x}$  are as large as they can. Here, the notation  $\tilde{x} \succ (x_1, \dots, x_m)$  means that

- $\sum_{j=1}^m \delta_{\lambda,j} \tilde{x}_j = \sum_{j=1}^m \delta_{\lambda,j} x_j = k + 1$ ,
- $\sum_{j=1}^l \delta_{\lambda,j} \tilde{x}_j \geq \sum_{j=1}^l \delta_{\lambda,j} x_j \quad \forall 1 \leq l \leq m$ .

Therefore, by the weighted Karamata's inequality (see [20]), we have that for every  $x \in B_\lambda$

$$F_{\delta_\lambda}(x) \leq F_{\delta_\lambda}(\tilde{x}) \leq \max_{(\delta,x) \in M} F(\delta, x).$$

Since

$$\max_{(\delta,x) \in M} F(\delta, x) \leq \max_{\lambda \in [\frac{1}{2}, 1], x \in B_\lambda} F_{\delta_\lambda}(x),$$

we see that

$$\begin{aligned} \max_{(\delta,x) \in M} F(\delta, x) &= \max_{\lambda \in [\frac{1}{2}, 1]} F_{\delta_\lambda}(\tilde{x}) = \max_{\lambda \in [\frac{1}{2}, 1]} \left\{ \frac{1}{2} D \log D + \frac{n-k}{2(n+1)} \log \left( \frac{1}{n+1} \right) \right\} \\ &= \frac{1}{2} D \log D - \frac{n-k}{n+1} \log(n+1). \end{aligned}$$

Thus,

$$\prod_{j=1}^m \|P_H v_j\|_2^{\delta_j} \leq e^{\frac{1}{2} D \log D - \frac{n-k}{2(n+1)} \log(n+1)} = \frac{(k+1)^{\frac{k+1}{2(n+1)}}}{(n+1)^{\frac{n+1}{2(n+1)}}}.$$

Since  $|K \cap F_0| \leq |C \cap F_0|$ , it follows that

$$|K \cap F|^{1/k} \leq \frac{1}{(k+1)^{\frac{n-k}{2k(n+1)}}} \sqrt{\frac{n(n+1)}{k(k+1)}} |S_k|^{1/k}.$$

Finally, assume now that  $K \subseteq \mathbb{R}^n$  is a (not necessarily symmetric) convex body in John's position and  $F_h$  is a  $k$ -dimensional affine subspace at distance  $h$  from 0. We continue to follow the notation introduced in Section 3.1. Given the  $k$ -dimensional affine subspace  $F_h$ , we take the linear subspace  $H = \text{span}\{(x, \sqrt{n}) : x \in F_h\} \in G_{n+1, k+1}$  as in (3.4). By Lemma 3.1, we see that

$$L \cap H = \left\{ (x, r) \in \mathbb{R}^{n+1} : r \geq 0, x \in \frac{r}{\sqrt{n}}(C \cap F_h) \right\}.$$

Recall that  $J = \{1 \leq j \leq m : P_H v_j \neq 0\}$  and  $s_j = \frac{1}{\|P_H v_j\|_2}$  for all  $j \in J$ . Using (3.6) and the Brascamp–Lieb inequality (Theorem 2.1), we have that

$$\begin{aligned} \int_{L \cap H} e^{-\sum_{j \in J} \kappa_j s_j \langle y, w_j \rangle} dy &= \int_H \prod_{j \in J} (\chi_{[0, \infty)}(\langle y, v_j \rangle) e^{-s_j \langle y, w_j \rangle})^{\kappa_j} dy \\ &= \int_H \prod_{j \in J} (\chi_{[0, \infty)}(\langle y, P_H v_j \rangle) e^{-s_j \langle y, w_j \rangle})^{\kappa_j} dy \\ &= \int_H \prod_{j \in J} (\chi_{[0, \infty)}(\langle y, w_j \rangle) e^{-s_j \langle y, w_j \rangle})^{\kappa_j} dy \\ &\leq \prod_{j \in J} \left( \int_0^\infty e^{-s_j t} dt \right)^{\kappa_j} = \prod_{j \in J} \|P_H v_j\|_2^{\delta_j \|P_H v_j\|_2^2} \leq 1. \end{aligned}$$

Taking into account (3.5) and (3.7), we obtain

$$\begin{aligned} \int_{L \cap H} e^{-\sum_{j \in J} \kappa_j s_j \langle y, w_j \rangle} dy &= \int_0^\infty \int_{\frac{r}{\sqrt{n}}(C \cap F_h)} e^{-r\sqrt{n+1}} dx \sqrt{\frac{n+h^2}{n}} dr \\ &= \int_0^\infty \frac{r^k (n+h^2)^{\frac{1}{2}}}{n^{\frac{k+1}{2}}} |C \cap F_h| e^{-r\sqrt{n+1}} dr \\ &= \frac{(n+h^2)^{\frac{1}{2}} k!}{n^{\frac{k+1}{2}} (n+1)^{\frac{k+1}{2}}} |C \cap F_h|. \end{aligned}$$

Since  $|K \cap F_h| \leq |C \cap F_h|$ , we get

$$|K \cap F_h| \leq \frac{n^{\frac{k}{2}}(n+1)^{\frac{k+1}{2}}}{k!} \sqrt{\frac{n}{n+h^2}}$$

or, equivalently,

$$|K \cap F_h|^{1/k} \leq \sqrt{\frac{n(n+1)^{1+\frac{1}{k}}}{k(k+1)^{1+\frac{1}{k}}} \left(\frac{n}{n+h^2}\right)^{\frac{1}{2k}}} |S_k|^{1/k}.$$

■

## 5 Volume of Projections of Convex Bodies in Löwner's Position

In this section, we will give the proof of Theorem 1.2.

**Proof of Theorem 1.2.** Let us start with the symmetric case. Assume that  $K$  is a centrally symmetric convex body in John's position and  $F_0 \in G_{n,k}$  a  $k$ -dimensional linear subspace. We follow the notation in Section 3.2. By (3.11), we get that  $K \cap F_0 \subseteq C_0 \cap F_0$ . This implies that

$$(K \cap F_0)^\circ \supseteq (C_0 \cap F_0)^\circ = \text{conv}\{\pm P_{F_0} u_j : j \in J_0\}.$$

It follows that for every  $x \in F_0$

$$\begin{aligned} h_{K \cap F_0}(x) &\leq h_{C_0 \cap F_0}(x) = \|x\|_{(C_0 \cap F_0)^\circ} = \inf \left\{ \sum_{j \in J_0} |\alpha_j| : x = \sum_{j \in J_0} \alpha_j P_{F_0} u_j \right\} \\ &= \inf \left\{ \sum_{j \in J_0} |\alpha_j| : x = \sum_{j \in J_0} \alpha_j \|P_{F_0} u_j\|_2 v_j^0 \right\} = \inf \left\{ \sum_{j \in J_0} \frac{|\beta_j|}{\|P_{F_0} u_j\|_2} : x = \sum_{j \in J_0} \beta_j v_j^0 \right\} \\ &= \inf \left\{ \sum_{j \in J_0} \delta_j^0 |\theta_j| t_j : x = \sum_{j \in J_0} \delta_j^0 \theta_j v_j^0 \right\}, \end{aligned}$$

where  $t_j = \frac{1}{\|P_{F_0} u_j\|_2} = \left(\frac{c_j}{\delta_j^0}\right)^{\frac{1}{2}}$  for all  $j \in J_0$ . For every  $j \in J_0$ , we set

$$f_j(t) := e^{-|t|t_j}, \quad t \in \mathbb{R}.$$

Then, if  $x = \sum_{j \in J_0} \delta_j^0 \theta_j v_j^0$  for some  $\{\theta_j\}_{j \in J_0} \subseteq \mathbb{R}$ , we have

$$\prod_{j \in J_0} f_j^{\delta_j^0}(\theta_j) = e^{-\sum_{j \in J_0} \delta_j |\theta_j| t_j} \leq e^{-h_{K \cap F_0}(x)}.$$

Using the reverse Brascamp–Lieb inequality (Theorem 2.1), one obtains

$$\begin{aligned} k! |(K \cap F_0)^\circ| &= \int_F e^{-h_{K \cap F_0}(x)} dx \geq \prod_{j \in J} \left( \int_{\mathbb{R}} e^{-|t| t_j} dt \right)^{\delta_j} \\ &= \frac{2^k}{\prod_{j \in J_0} t_j^{\delta_j}} = \frac{2^k}{\prod_{j \in J_0} \left(\frac{c_j}{\delta_j}\right)^{\frac{\delta_j}{2}}}. \end{aligned}$$

As we have seen in the proof of Theorem 1.1

$$\prod_{j \in J_0} \left(\frac{c_j}{\delta_j^0}\right)^{\frac{\delta_j}{k}} \leq \sum_{j \in J_0} \frac{\delta_j^0}{k} \frac{c_j}{\delta_j^0} = \frac{1}{k} \sum_{j \in J_0} c_j \leq \frac{1}{k} \sum_{j=1}^m c_j = \frac{n}{k}.$$

Taking into account that  $|(B_\infty^k)^\circ| = |B_1^k| = \frac{2^k}{k!}$ , we obtain

$$|(K \cap F_0)^\circ|^{1/k} \geq \sqrt{\frac{k}{n}} |(B_\infty^k)^\circ|^{1/k}.$$

Assume now that  $K \subseteq \mathbb{R}^n$  is a (not necessarily symmetric) convex body in John's position and  $F_0 \in G_{n,k}$  is a  $k$ -dimensional linear subspace. We follow the notation introduced in Section 3.1. For  $H = F_0 \times \mathbb{R} \in G_{n+1,k+1}$  as in (3.4), we have that

$$\begin{aligned} (C \cap F_0)^\circ &= P_{F_0}(C^\circ) = P_{F_0}(\text{conv}\{u_j : 1 \leq j \leq m\}) \\ &= \text{conv}\{P_{F_0} u_j : 1 \leq j \leq m\}. \end{aligned}$$

For any  $y = (x, r) \in H = F_0 \times \mathbb{R}$ , we write

$$N(y) = \inf \left\{ \sum_{j=1}^m \frac{\kappa_j \theta_j}{\sqrt{n \|P_{F_0} u_j\|_2^2 + 1}} : \theta_j \geq 0, y = \sum_{j=1}^m \kappa_j \theta_j w_j \right\},$$

where the latter infimum is understood as  $\infty$  if there do not exist  $\{\theta_j\}_{j=1}^m$  with  $\theta_j \geq 0$  such that  $y = \sum_{j=1}^m \kappa_j \theta_j w_j$ . Notice that for any  $\{\theta_j\}_{j=1}^m \subseteq \mathbb{R}$ ,

$$\begin{aligned} y = \sum_{j \in J} \kappa_j \theta_j w_j &\Leftrightarrow (x, r) = \left( -\sum_{j=1}^m \frac{\kappa_j \theta_j P_{F_0}(u_j)}{\sqrt{\|P_{F_0} u_j\|_2^2 + \frac{1}{n}}}, \sum_{j=1}^m \frac{\kappa_j \theta_j}{\sqrt{n \|P_{F_0} u_j\|_2^2 + 1}} \right) \\ &\Leftrightarrow (x, r) = \left( -r \sqrt{n} \sum_{j=1}^m \frac{\kappa_j \theta_j P_{F_0}(u_j)}{r \sqrt{n \|P_{F_0} u_j\|_2^2 + 1}}, \sum_{j=1}^m \frac{\kappa_j \theta_j}{\sqrt{n \|P_{F_0} u_j\|_2^2 + 1}} \right). \end{aligned}$$

Then there exist  $\{\theta_j\}_{j=1}^m \subseteq \mathbb{R}$  with  $\theta_j \geq 0$  for every  $1 \leq j \leq m$  such that the latter equality holds if and only if

$$(x, r) \in L_1 := \{(x, r) \in F_0 \times \mathbb{R} : r \geq 0 : x \in -r \sqrt{n} (C \cap F_0)^\circ\},$$

and for all such  $y = (x, r) \in L_1$ , we have that  $N(y) = r$ . Therefore, for every  $y \in H$ ,

$$\sup_{y = \sum_{j=1}^m \kappa_j \theta_j w_j} \prod_{j=1}^m \left( \chi_{[0, \infty)}(\theta_j) e^{-\frac{\theta_j}{\sqrt{n \|P_{F_0} u_j\|_2^2 + 1}}} \right)^{\kappa_j} = e^{-N(y)}.$$

Thus, by (3.6) and the reverse Brascamp–Lieb inequality (Theorem 2.1),

$$\int_H e^{-N(y)} dy \geq \prod_{j=1}^m \left( \int_0^\infty e^{-\frac{t}{\sqrt{n \|P_{F_0} u_j\|_2^2 + 1}}} dt \right)^{\kappa_j}.$$

On the one hand,

$$\int_H e^{-N(y)} dy = \int_0^\infty e^{-r} |-r \sqrt{n} (C \cap F_0)^\circ| dr = k! n^{k/2} |(C \cap F_0)^\circ|.$$

On the other hand, for every  $1 \leq j \leq m$ ,

$$\int_0^\infty e^{-\frac{t}{\sqrt{n \|P_{F_0} u_j\|_2^2 + 1}}} dt = \sqrt{n \|P_{F_0} u_j\|_2^2 + 1} = \sqrt{n+1} \|P_H v_j\|_2.$$

Since  $(K \cap F_0)^\circ \supseteq (C \cap F_0)^\circ$ , we obtain

$$\begin{aligned} |(K \cap F_0)^\circ| &\geq \frac{(n+1)^{\frac{k+1}{2}} \prod_{j=1}^m \|P_{H^j} v_j\|_2^{\delta_j \|P_{H^j} v_j\|_2^2}}{k! n^{k/2}} \\ &= \left(\frac{n+1}{k+1}\right)^{\frac{k+1}{2}} \left(\frac{k}{n}\right)^{k/2} \prod_{j=1}^m \|P_{H^j} v_j\|_2^{\delta_j \|P_{H^j} v_j\|_2^2} |S_k^\circ|. \end{aligned}$$

For the convex function  $f(x) = x \log x$ , we apply Jensen's inequality to get

$$\sum_{j=1}^m \frac{\delta_j}{n+1} \|P_{H^j} v_j\|_2^2 \log \|P_{H^j} v_j\|_2^2 \geq f\left(\sum_{j=1}^m \frac{\delta_j \|P_{H^j} v_j\|_2^2}{n+1}\right) = f\left(\frac{k+1}{n+1}\right).$$

Thus,

$$\prod_{j=1}^m \|P_{H^j} v_j\|_2^{\delta_j \|P_{H^j} v_j\|_2^2} = e^{\frac{n+1}{2} \sum_{j=1}^m \frac{\delta_j}{n+1} \|P_{H^j} v_j\|_2^2 \log \|P_{H^j} v_j\|_2^2} \geq \left(\frac{k+1}{n+1}\right)^{\frac{k+1}{2}}.$$

Therefore,

$$|(K \cap F_0)^\circ|^{\frac{1}{k}} \geq \sqrt{\frac{k}{n}} |S_k^\circ|^{\frac{1}{k}}.$$

■

### 6 Mean Width of Sections of Convex Bodies in John's Position

In this section, we will prove Theorem 1.3.

**Proof of Theorem 1.3.** Let us start with the symmetric case. Assume that  $K$  is a centrally symmetric convex body in John's position and  $F_0 \in G_{n,k}$  is a  $k$ -dimensional linear subspace. We follow the notation in Section 3.2. By (3.11), we have that  $K \cap F_0 \subseteq C_0 \cap F_0$  and

$$(K \cap F_0)^\circ \supseteq (C_0 \cap F_0)^\circ = \text{conv}\{\pm P_{F_0} u_j : j \in J_0\}.$$

It follows that for every  $x \in F_0$

$$\begin{aligned} h_{K \cap F_0}(x) &\leq h_{C_0 \cap F_0}(x) = \inf \left\{ \sum_{j \in J_0} |\alpha_j| : x = \sum_{j \in J_0} \alpha_j P_{F_0} u_j \right\} \\ &= \inf \left\{ \sum_{j \in J_0} |\alpha_j| : x = \sum_{j \in J_0} \alpha_j \|P_{F_0} u_j\|_2 v_j^0 \right\} = \inf \left\{ \sum_{j \in J_0} |\beta_j| t_j : x = \sum_{j \in J_0} \beta_j v_j^0 \right\}, \end{aligned}$$

where  $t_j = \frac{1}{\|P_{F_0} u_j\|_2}$  for all  $j \in J_0$ . For every  $x \in F_0$ , we write  $x = \sum_{j \in J_0} \delta_j^0 \langle x, v_j^0 \rangle v_j^0$ , therefore

$$h_{K \cap F_0}(x) \leq \sum_{j \in J_0} \delta_j^0 t_j |\langle x, v_j^0 \rangle|. \quad (6.1)$$

If  $G_1$  is a standard Gaussian random vector in  $F_0$  and  $G_2$  is a standard Gaussian random vector on  $\mathbb{R}^k$ , using (6.1), we get

$$\begin{aligned} \mathbb{E} h_{K \cap F_0}(G_1) &\leq \sum_{j \in J_0} \delta_j^0 t_j \mathbb{E} |\langle G_1, v_j \rangle| = \mathbb{E} |\langle G_2, e_1 \rangle| \sum_{j \in J_0} \delta_j^0 t_j \\ &= \frac{1}{k} \mathbb{E} \|G_2\|_1 \sum_{j \in J_0} \delta_j^0 t_j = \frac{1}{k} \sum_{j \in J_0} \delta_j^0 t_j \mathbb{E} h_{B_\infty^k}(G_2). \end{aligned}$$

By Hölder's inequality and (3.9),

$$\begin{aligned} \frac{1}{k} \sum_{j \in J_0} \delta_j^0 t_j &= \frac{1}{k} \sum_{j \in J_0} c_j \|P_{F_0} u_j\|_2 \leq \frac{1}{k} \left( \sum_{j \in J_0} c_j \right)^{\frac{1}{2}} \left( \sum_{j \in J_0} c_j \|P_{F_0} u_j\|_2^2 \right)^{\frac{1}{2}} \\ &\leq \frac{1}{k} \left( \sum_{j=1}^m c_j \right)^{\frac{1}{2}} \left( \sum_{j=1}^m c_j \|P_{F_0} u_j\|_2^2 \right)^{\frac{1}{2}} = \frac{\sqrt{nk}}{k} = \sqrt{\frac{n}{k}}. \end{aligned}$$

Hence,

$$\mathbb{E} h_{K \cap F_0}(G_1) \leq \sqrt{\frac{n}{k}} \mathbb{E} h_{B_\infty^k}(G_2).$$

Equivalently, by (2.2),

$$w(K \cap F_0) \leq \sqrt{\frac{n}{k}} w(B_\infty^k).$$



Let us now assume that  $K$  is a not necessarily symmetric convex body in John's position and let  $F_0 \in G_{n,k}$ . We follow the notation introduced in Section 3.1. For every  $x \in F_0$ , we have

$$h_{K \cap F_0}(x) \leq h_{C \cap F_0}(x) = \inf \left\{ \sum_{j=1}^m a_j : x = \sum_{j=1}^m a_j P_{F_0} u_j, a_j \geq 0 \right\}.$$

Let  $\theta \in S^{n-1} \cap F_0$ . By (2.1), we have  $\sum_{j=1}^m c_j P_{F_0} u_j = 0$ , so we may write

$$\theta = \sum_{j=1}^m c_j \left( \langle \theta, P_{F_0} u_j \rangle - \min_{1 \leq k \leq m} \langle \theta, P_{F_0} u_k \rangle \right) P_{F_0} u_j.$$

Setting (like in the symmetric case before)  $J_0 = \{1 \leq j \leq m : P_{F_0} u_j \neq 0\}$  and  $v_j^0 = \frac{P_{F_0} u_j}{\|P_{F_0} u_j\|_2}$  for  $j \in J_0$ , we get

$$\begin{aligned} w(K \cap F_0) &\leq w(C \cap F_0) = \int_{S^{n-1} \cap F_0} h_{C \cap F_0}(\theta) d\sigma(\theta) \\ &\leq \int_{S^{n-1} \cap F_0} \sum_{j=1}^m c_j \left( \langle \theta, P_{F_0} u_j \rangle - \min_{1 \leq k \leq m} \langle \theta, P_{F_0} u_k \rangle \right) d\sigma(\theta) \\ &= n \int_{S^{n-1} \cap F_0} \max_{1 \leq k \leq m} \langle \theta, -P_{F_0} u_k \rangle d\sigma(\theta) \leq n \int_{S^{n-1} \cap F_0} \max_{1 \leq k \leq m} |\langle \theta, P_{F_0} u_k \rangle| d\sigma(\theta) \\ &\leq n \int_{S^{n-1} \cap F_0} \max_{1 \leq k \leq m} \|P_{F_0} u_k\|_2 \max_{k \in J_0} |\langle \theta, v_k^0 \rangle| d\sigma(\theta) \leq n \int_{S^{n-1} \cap F_0} \max_{k \in J_0} |\langle \theta, v_k^0 \rangle| d\sigma(\theta). \end{aligned}$$

It is a well-known fact (see, for instance, [3, Proposition 9.1.5 and Lemma 5.2.11]) that for any  $\{\theta_i\}_{i=1}^N \subseteq S^{n-1}$ , one has that

$$\int_{S^{n-1}} \max_{1 \leq k \leq N} |\langle \theta, \theta_k \rangle| d\sigma(\theta) \simeq \sqrt{\frac{\log N}{n}}.$$

Therefore, there exists an absolute constant  $C_1$  such that

$$\int_{S^{n-1} \cap F} \max_{k \in J_0} |\langle \theta, v_k^0 \rangle| d\sigma(\theta) \leq C_1 \sqrt{\frac{\log m}{k}}.$$

Taking into account that  $m = O(n^2)$  and that  $w(S_k) \simeq \sqrt{k \log k}$ , we obtain that there exists an absolute constant  $C_2 > 0$  such that

$$w(K \cap F_0) \leq C_2 \frac{n}{k} \sqrt{\frac{\log n}{\log k}} w(S_k). \quad \blacksquare$$

## 7 Mean Width of Projections of Convex Bodies in Löwner's Position

In this section, we will prove Theorem 1.4. We will make use of the following lemma.

**Lemma 7.1.** Let  $K \subseteq \mathbb{R}^n$  be a (not necessarily symmetric) convex body and  $F_h$  be a  $k$ -dimensional affine subspace at distance  $h$  from the origin. Take some  $\alpha \in \mathbb{R}$  and  $\beta \leq 0$ . Identifying the origin with the closest point in  $F_h$  to 0, we identify  $F_h$  with  $\mathbb{R}^k$  and let  $\gamma_k$  be the  $k$ -dimensional Gaussian measure on  $F_h$ . Then,

$$\begin{aligned} & \sqrt{\frac{n+h^2}{n}} \int_0^\infty \frac{e^{-\frac{(r-\alpha\sqrt{n+1})^2}{2}}}{\sqrt{2\pi}} e^{\beta r\sqrt{n+1}} \gamma_k \left( \frac{r}{\sqrt{n}} (C \cap F_h) \right) dr \leq \\ & \leq \int_0^\infty \frac{e^{-\frac{(r-\alpha d_1)^2}{2}}}{\sqrt{2\pi}} e^{\beta r\sqrt{k+1}} \gamma_k \left( r\sqrt{k+1} \Delta_k \right) dr, \end{aligned}$$

where  $C$  is defined as in (3.1),  $\Delta_k$  denotes the regular  $k$ -dimensional simplex as introduced in Section 2.3,  $d_1 = \frac{1}{\sqrt{k+1}} \sum_{j \in J} \delta_j \|P_H v_j\|_2$ , and  $H$  is defined as in (3.4),  $\delta_j$  and  $v_j$  as in (3.2), and  $J$  as in (3.6).

**Proof.** Following the notation introduced in Section 3.1, let  $L$  be the cone defined in (3.3). By Lemma 3.1, we have that

$$L \cap H = \left\{ (x, r) : r \geq 0, x \in \frac{r}{\sqrt{n}} (C \cap F_h) \right\}.$$

For any  $\alpha, \beta \in \mathbb{R}$ , let  $\mu_{\alpha, \beta}$  be the measure on  $H$  whose density with respect to the Lebesgue measure at a point  $y = (x, r)$  is

$$d\mu_{\alpha, \beta}(y) = \frac{e^{-\frac{\|y\|_2^2}{2}}}{(2\pi)^{\frac{k+1}{2}}} e^{(\alpha+\beta)r\sqrt{n+1}} dy.$$

For any  $\alpha, \beta \in \mathbb{R}$ , taking into account (3.5), we have that

$$\begin{aligned} \mu_{\alpha, \beta}(L \cap H) &= \int_0^\infty \frac{e^{-\frac{r^2}{2}}}{\sqrt{2\pi}} e^{\alpha r\sqrt{n+1}} e^{\beta r\sqrt{n+1}} \gamma_k \left( \frac{r}{\sqrt{n}} (C \cap F_h) \right) \sqrt{\frac{n+h^2}{n}} dr \\ &= e^{\frac{\alpha^2(n+1)}{2}} \int_0^\infty \frac{e^{-\frac{(r-\alpha\sqrt{n+1})^2}{2}}}{\sqrt{2\pi}} e^{\beta r\sqrt{n+1}} \gamma_k \left( \frac{r}{\sqrt{n}} (C \cap F_h) \right) \sqrt{\frac{n+h^2}{n}} dr. \end{aligned}$$

Recall that  $J = \{1 \leq j \leq m : P_H v_j \neq 0\}$ ,  $s_j = \frac{1}{\|P_H v_j\|_2}$  for every  $j \in J$  and  $y = (x, r) \in H$ . Using the definition of  $L$ , the identity (3.7), and the Brascamp–Lieb inequality (Theorem 2.1), we have

$$\begin{aligned} \mu_{\alpha, \beta}(L \cap H) &= \int_H \frac{e^{-\frac{\|y\|_2^2}{2}}}{(2\pi)^{(k+1)/2}} e^{\alpha r\sqrt{n+1}} e^{\beta r\sqrt{n+1}} \prod_{j \in J} \chi_{[0, \infty)}(\langle y, v_j \rangle) dy \\ &= \int_H \frac{e^{-\frac{\|y\|_2^2}{2}}}{(2\pi)^{(k+1)/2}} e^{\alpha r\sqrt{n+1}} e^{\beta r\sqrt{n+1}} \prod_{j \in J} \chi_{[0, \infty)}(\langle y, w_j \rangle) dy \\ &= \int_H \frac{e^{-\frac{\sum_{j \in J} \kappa_j \langle y, w_j \rangle^2}{2}}}{(2\pi)^{(k+1)/2}} e^{\sum_{j \in J} \kappa_j (\alpha + \beta) s_j \langle y, w_j \rangle} \prod_{j \in J} \chi_{[0, \infty)}(\langle y, w_j \rangle) dy \\ &= \int_H \prod_{j \in J} \left( \frac{e^{-\frac{\langle y, w_j \rangle^2}{2}}}{\sqrt{2\pi}} e^{(\alpha + \beta) s_j \langle y, w_j \rangle} \chi_{[0, \infty)}(\langle y, w_j \rangle) \right)^{\kappa_j} dy \\ &\leq \prod_{j \in J} \left( \int_0^\infty \frac{e^{-\frac{t^2}{2}}}{\sqrt{2\pi}} e^{\alpha s_j t} e^{\beta s_j t} dt \right)^{\kappa_j} = e^{\frac{\alpha^2 \sum_{j \in J} \kappa_j s_j^2}{2}} \prod_{j=1}^m \left( \int_0^\infty \frac{e^{-\frac{(t-s_j)^2}{2}}}{\sqrt{2\pi}} e^{\beta s_j t} dt \right)^{\kappa_j} \\ &= e^{\frac{\alpha^2(n+1)}{2}} \prod_{j \in J} \left( \int_0^\infty \frac{e^{-\frac{(t-s_j)^2}{2}}}{\sqrt{2\pi}} e^{\beta s_j t} dt \right)^{\kappa_j} \end{aligned}$$

Therefore, for any  $\alpha, \beta \in \mathbb{R}$ ,

$$\begin{aligned} &\sqrt{\frac{n+h^2}{n}} \int_0^\infty \frac{e^{-\frac{(r-\alpha\sqrt{n+1})^2}{2}}}{\sqrt{2\pi}} e^{\beta r\sqrt{n+1}} \gamma_k \left( \frac{r}{\sqrt{n}} (C \cap F_h) \right) dr \\ &\leq \prod_{j \in J} \left( \int_0^\infty \frac{e^{-\frac{(t-s_j)^2}{2}}}{\sqrt{2\pi}} e^{\beta s_j t} dt \right)^{\kappa_j}. \end{aligned}$$

Notice that  $s_j = \frac{1}{\|P_H v_j\|_2} \geq 1$  for every  $j \in J$ , which implies that, for any  $\beta \leq 0$ , one has  $\beta s_j \leq \beta$  for every  $j \in J$ . Using this inequality in the second following inequality and the Prékopa–Leindler inequality (see [37, Lemma 1.2]) in the third following inequality, we have that for any  $\beta \leq 0$

$$\begin{aligned}
& \sqrt{\frac{n+h^2}{n}} \int_0^\infty \frac{e^{-\frac{(r-\alpha\sqrt{n+1})^2}{2}}}{\sqrt{2\pi}} e^{\beta r\sqrt{n+1}} \gamma_k \left( \frac{r}{\sqrt{n}} (C \cap F_h) \right) dr \\
& \leq \prod_{j \in J} \left( \int_0^\infty \frac{e^{-\frac{(t-\alpha s_j)^2}{2}}}{\sqrt{2\pi}} e^{\beta s_j t} dt \right)^{\kappa_j} \\
& \leq \prod_{j \in J} \left( \int_0^\infty \frac{e^{-\frac{(t-\alpha s_j)^2}{2}}}{\sqrt{2\pi}} e^{\beta t} dt \right)^{\kappa_j} \leq \left( \int_0^\infty \frac{e^{-\frac{(t-\frac{\alpha}{k+1} \sum_{j \in J} \kappa_j s_j)^2}{2}}}{\sqrt{2\pi}} e^{\beta t} dt \right)^{k+1} \\
& = \int_{[0, \infty)^{k+1}} \prod_{i=1}^{k+1} \frac{e^{-\frac{(t_i - \frac{\alpha d_1}{\sqrt{k+1}})^2}{2}}}{\sqrt{2\pi}} e^{\beta t_i} dt \\
& = \int_{[-\frac{\alpha d_1}{\sqrt{k+1}}, \infty)^{k+1}} \prod_{i=1}^{k+1} \frac{e^{-\frac{\|t\|_2^2}{2}}}{\sqrt{2\pi}} e^{\beta \sqrt{k+1}(t, v_0)} e^{\beta \alpha d_1 \sqrt{k+1}} dt,
\end{aligned}$$

where  $v_0 = \left( \frac{1}{\sqrt{k+1}}, \dots, \frac{1}{\sqrt{k+1}} \right)$ . Therefore, for any  $\alpha \in \mathbb{R}$  and any  $\beta \leq 0$ ,

$$\begin{aligned}
& \sqrt{\frac{n+h^2}{n}} \int_0^\infty \frac{e^{-\frac{(r-\alpha\sqrt{n+1})^2}{2}}}{\sqrt{2\pi}} e^{\beta r\sqrt{n+1}} \gamma_k \left( \frac{r}{\sqrt{n}} (C \cap F_h) \right) dr \\
& \leq \int_{[-\frac{\alpha d_1}{\sqrt{k+1}}, \infty)^{k+1}} \prod_{i=1}^{k+1} \frac{e^{-\frac{\|t\|_2^2}{2}}}{\sqrt{2\pi}} e^{\beta \sqrt{k+1}(t, v_0)} e^{\beta \alpha d_1 \sqrt{k+1}} dt \\
& = \int_{-\alpha d_1}^\infty \frac{e^{-\frac{t^2}{2}}}{\sqrt{2\pi}} e^{\beta \sqrt{k+1}t} e^{\beta \alpha d_1 \sqrt{k+1}} \gamma_k \left( (t + \alpha d_1) \sqrt{k+1} \Delta_k \right) dt \\
& = \int_0^\infty \frac{e^{-\frac{(r-\alpha d_1)^2}{2}}}{\sqrt{2\pi}} e^{\beta \sqrt{k+1}r} \gamma_k \left( r \sqrt{k+1} \Delta_k \right) dr,
\end{aligned}$$

where  $\Delta_k$  denotes the regular  $k$ -dimensional simplex as introduced in Section 2.3. ■

**Proof of Theorem 1.4.** Let us start with the symmetric case. Assume that  $K$  is a centrally symmetric convex body in John's position and  $F_0 \in G_{n,k}$  is a  $k$ -dimensional linear subspace. We want to prove that

$$w((K \cap F_0)^\circ) \geq w\left(\left(\sqrt{\frac{n}{k}}B_\infty^k\right)^\circ\right).$$

Equivalently, by (2.3), we want to prove that

$$\mathbb{E}\|G_1\|_{K \cap F_0} \geq \mathbb{E}\|G_2\|_{\sqrt{\frac{n}{k}}B_\infty^k},$$

where  $G_1$  is a standard Gaussian random vector on  $F_0$  and  $G_2$  is a standard Gaussian random vector on  $\mathbb{R}^k$ . If  $L \subseteq \mathbb{R}^k$  is a convex body containing the origin in its interior and  $G$  is a standard Gaussian random vector then

$$\mathbb{E}\|G\|_L = \int_0^\infty \mathbb{P}(\|G\|_L \geq t) dt = \int_0^\infty \gamma_k(\mathbb{R}^n \setminus tL) dt, \tag{7.1}$$

where  $\gamma_k(A)$  denotes the Gaussian measure of the  $k$ -dimensional set  $A$ . Therefore, the statement we want to prove is equivalent to

$$\int_0^\infty \gamma_k(F_0 \setminus t(K \cap F_0)) dt \geq \int_0^\infty \gamma_k\left(\mathbb{R}^n \setminus t\sqrt{\frac{n}{k}}B_\infty^k\right) dt$$

or, equivalently,

$$\int_0^\infty (1 - \gamma_k(t(K \cap F_0))) dt \geq \int_0^\infty \left(1 - \gamma_k\left(t\sqrt{\frac{n}{k}}B_\infty^k\right)\right) dt.$$

We are going to prove that for any  $t \geq 0$

$$\gamma_k(t(K \cap F_0)) \leq \gamma_k\left(t\sqrt{\frac{n}{k}}B_\infty^k\right),$$

which implies the latter inequality.

We follow the notation in Section 3.2. By (3.10) we have that

$$K \cap F_0 \subseteq C_0 \cap F_0 = \left\{x \in F_0 : |\langle x, v_j^0 \rangle| \leq t_j, \forall j \in J_0\right\},$$

where  $t_j = \frac{1}{\|P_{F_0} u_j\|_2} = \left(\frac{c_j}{\delta_j^0}\right)^{1/2}$  for all  $j \in J_0$ . Therefore, for every  $t \geq 0$ ,

$$t(K \cap F_0) \subseteq t(C_0 \cap F_0) = \left\{x \in F_0 : |\langle x, v_j^0 \rangle| \leq t t_j, \forall j \in J_0\right\}.$$

By (3.9) and the Brascamp–Lieb inequality (Theorem 2.1),

$$\begin{aligned} \gamma_k(t(K \cap F_0)) &\leq \gamma_k(t(C_0 \cap F_0)) = \int_{F_0} \left( \prod_{j \in J} \chi_{[-t t_j, t t_j]}(\langle x, v_j^0 \rangle) \right) \frac{e^{-\frac{\|x\|_2^2}{2}}}{(2\pi)^{k/2}} dx \\ &= \int_{F_0} \left( \prod_{j \in J_0} \chi_{[-t t_j, t t_j]}(\langle x, v_j^0 \rangle) \right) \frac{e^{-\frac{\sum_{j \in J_0} \delta_j^0 \langle x, v_j^0 \rangle^2}{2}}}{(2\pi)^{k/2}} dx \\ &= \int_{F_0} \prod_{j \in J_0} \left( \chi_{[-t t_j, t t_j]}(\langle x, v_j^0 \rangle) \frac{e^{-\frac{\langle x, v_j^0 \rangle^2}{2}}}{\sqrt{2\pi}} \right)^{\delta_j^0} dx \\ &\leq \prod_{j \in J_0} \left( \int_{-t t_j}^{t t_j} \frac{e^{-\frac{t^2}{2}}}{\sqrt{2\pi}} dt \right)^{\delta_j^0} = \left( \prod_{j \in J_0} \gamma_1(t t_j [-e_1, e_1])^{\frac{\delta_j^0}{k}} \right)^k. \end{aligned}$$

Since  $\gamma_1$  is log-concave, we obtain

$$\gamma_k(t(K \cap F_0)) \leq \gamma_1 \left( \left( t \sum_{j \in J_0} \frac{t_j \delta_j^0}{k} \right) [-e_1, e_1] \right)^k = \gamma_k \left( \left( t \sum_{j \in J_0} \frac{t_j \delta_j^0}{k} \right) B_\infty^k \right).$$

By Hölder's inequality and (3.9), we have that

$$\begin{aligned} \sum_{j \in J_0} \frac{t_j \delta_j^0}{k} &= \sum_{j \in J_0} \frac{\sqrt{c_j \delta_j^0}}{k} \leq \frac{1}{k} \left( \sum_{j \in J_0} c_j \right)^{1/2} \left( \sum_{j \in J_0} \delta_j^0 \right)^{1/2} \\ &\leq \frac{1}{k} \left( \sum_{j=1}^m c_j \right)^{1/2} \left( \sum_{j \in J_0} \delta_j^0 \right)^{1/2} = \sqrt{\frac{n}{k}}. \end{aligned}$$

Thus, for every  $t \geq 0$ ,

$$\gamma_k(t(K \cap F_0)) \leq \gamma_k \left( t \sqrt{\frac{n}{k}} B_\infty^k \right).$$

Assume now that  $K \subseteq \mathbb{R}^n$  is a (not necessarily symmetric) convex body in John's position and  $F_0 \in G_{n,k}$  is a linear subspace. Following the notation introduced in Section 3.1, we denote by  $H \in G_{n+1,k+1}$  the  $(k + 1)$ -dimensional linear subspace  $H = \text{span}\{(x, \sqrt{n}) : x \in F\} = F_0 \times \mathbb{R}$ , as in (3.4). By Lemma 3.1, we have that

$$L \cap H = \left\{ (x, r) \in F_0 \times \mathbb{R} : r \geq 0, x \in \frac{r}{\sqrt{n}}(C \cap F_0) \right\}.$$

Using Lemma 7.1 for  $\beta = 0$ , the linear subspace  $F_0 \in G_{n,k}$  and an arbitrary  $\alpha \in \mathbb{R}$ , we get

$$\int_0^\infty \frac{e^{-\frac{(r-\alpha\sqrt{n+1})^2}{2}}}{\sqrt{2\pi}} \gamma_k \left( \frac{r}{\sqrt{n}}(C \cap F_0) \right) dr \leq \int_0^\infty \frac{e^{-\frac{(r-\alpha d_1)^2}{2}}}{\sqrt{2\pi}} \gamma_k \left( r\sqrt{k+1}\Delta_k \right) dr,$$

where  $d_1 = \frac{1}{\sqrt{k+1}} \sum_{j \in J} \delta_j \|P_H v_j\|_2$ ,  $\delta_j$  and  $v_j$  are defined as in (3.2), and  $J$  as in (3.6). Applying the latter inequality to  $-\alpha$ , we also get

$$\int_0^\infty \frac{e^{-\frac{(r+\alpha\sqrt{n+1})^2}{2}}}{\sqrt{2\pi}} \gamma_k \left( \frac{r}{\sqrt{n}}(C \cap F_0) \right) dr \leq \int_0^\infty \frac{e^{-\frac{(r+\alpha d_1)^2}{2}}}{\sqrt{2\pi}} \gamma_k \left( r\sqrt{k+1}\Delta_k \right) dr$$

or, equivalently,

$$\int_{-\infty}^0 \frac{e^{-\frac{(r-\alpha\sqrt{n+1})^2}{2}}}{\sqrt{2\pi}} \gamma_k \left( \frac{|r|}{\sqrt{n}}(C \cap F_0) \right) dr \leq \int_{-\infty}^0 \frac{e^{-\frac{(r-\alpha d_1)^2}{2}}}{\sqrt{2\pi}} \gamma_k \left( |r|\sqrt{k+1}\Delta_k \right) dr.$$

Therefore, for any  $\alpha \in \mathbb{R}$ ,

$$\int_{-\infty}^\infty \frac{e^{-\frac{(r-\alpha\sqrt{n+1})^2}{2}}}{\sqrt{2\pi}} \gamma_k \left( \frac{|r|}{\sqrt{n}}(C \cap F_0) \right) dr \leq \int_{-\infty}^\infty \frac{e^{-\frac{(r-\alpha d_1)^2}{2}}}{\sqrt{2\pi}} \gamma_k \left( |r|\sqrt{k+1}\Delta_k \right) dr.$$

Hence,

$$\int_{-\infty}^\infty \frac{e^{-\frac{(r-\alpha\sqrt{n+1})^2}{2}}}{\sqrt{2\pi}} \gamma_k(F_0 \setminus \left( \left( \frac{|r|}{\sqrt{n}}(C \cap F_0) \right) \right)) dr \geq \int_{-\infty}^\infty \frac{e^{-\frac{(r-\alpha d_1)^2}{2}}}{\sqrt{2\pi}} \gamma_k \left( \mathbb{R}^k \setminus (|r|\sqrt{k+1}\Delta_k) \right) dr.$$

Integrating in  $\alpha \in \mathbb{R}$ , we obtain

$$\frac{1}{\sqrt{n+1}} \int_{-\infty}^\infty \gamma_k \left( F_0 \setminus \left( \frac{|r|}{\sqrt{n}}(C \cap F_0) \right) \right) dr \geq \frac{1}{d_1} \int_{-\infty}^\infty \gamma_k \left( \mathbb{R}^k \setminus (|r|\sqrt{k+1}\Delta_k) \right) dr.$$

Equivalently,

$$\frac{1}{\sqrt{n+1}} \int_0^\infty \gamma_k \left( F_0 \setminus \left( \frac{r}{\sqrt{n}} (C \cap F_0) \right) \right) dr \geq \frac{1}{d_1} \int_0^\infty \gamma_k \left( \mathbb{R}^k \setminus (r\sqrt{k+1}\Delta_k) \right) dr,$$

or

$$\sqrt{\frac{n}{n+1}} \int_0^\infty \gamma_k (F_0 \setminus (r(C \cap F_0))) dr \geq \frac{1}{d_1 \sqrt{k+1}} \int_0^\infty \gamma_k (\mathbb{R}^k \setminus (r\Delta_k)) dr.$$

Using (7.1), (2.3), and the fact that  $K \subseteq C$ , we obtain

$$w((K \cap F_0)^\circ) \geq \frac{1}{d_1} \sqrt{\frac{n+1}{n(k+1)}} w((\Delta_k)^\circ).$$

If  $S_k$  denotes the  $k$ -dimensional regular simplex in John's position, then

$$\sqrt{k(k+1)}\Delta_k = S_k.$$

Therefore, for any  $k$ -dimensional linear subspace  $F_0$ , we have

$$w((K \cap F_0)^\circ) \geq \sqrt{\frac{k(n+1)}{n}} \frac{1}{d_1} w((S_k)^\circ).$$

By Hölder's inequality and (3.6), we have that

$$d_1 \sqrt{k+1} = \sum_{j \in J} \delta_j \|P_H v_j\|_2 \leq \left( \sum_{j=1}^m \delta_j \right)^{\frac{1}{2}} \left( \sum_{j=1}^m \delta_j \|P_H v_j\|_2^2 \right)^{\frac{1}{2}} = \sqrt{(n+1)(k+1)}.$$

Thus,

$$w((K \cap F_0)^\circ) \geq \sqrt{\frac{k}{n}} w(S_k^\circ). \quad \blacksquare$$

## 8 The Wills Functional of Sections of Convex Bodies in John's Position

In this section, we will give the proof of Theorem 1.5.

**Proof of Theorem 1.5.** Let  $K$  be a centrally symmetric convex body in John's position and  $F_0 \in G_{n,k}$  a  $k$ -dimensional linear subspace. We follow the notation in Section 3.2.



By (3.10), we have that for every  $\lambda \geq 0$

$$\lambda(K \cap F_0) \subseteq \lambda(C_0 \cap F_0) = \left\{ x \in F_0 : |\langle x, v_j^0 \rangle| \leq \lambda t_j, \forall j \in J_0 \right\},$$

where  $t_j = \frac{1}{\|P_{F_0} u_j\|_2} = \left( \frac{c_j}{\delta_j^0} \right)^{1/2}$  for all  $j \in J_0$ . For every  $j \in J_0$ , we define  $f_j : \mathbb{R} \rightarrow [0, \infty)$  to be the function

$$f_j(t) = e^{-\pi d(tv_j^0, P_{\langle v_j^0 \rangle}(\lambda(C_0 \cap F_0)))^2} \quad \forall t \in \mathbb{R},$$

where  $\langle v_j^0 \rangle$  denotes the 1-dimensional subspace spanned by  $v_j^0$ . Then,

$$\int_{\mathbb{R}} f_j(t) dt = \int_{\mathbb{R}} e^{-\pi d(tv_j^0, P_{\langle v_j^0 \rangle}(\lambda(C_0 \cap F_0)))^2} dt = \mathcal{W}(P_{\langle v_j^0 \rangle}(\lambda(C_0 \cap F_0)))$$

and

$$P_{\langle v_j^0 \rangle}(\lambda(C_0 \cap F_0)) \subseteq [-\lambda t_j, \lambda t_j] v_j.$$

It follows that for every  $j \in J_0$ ,

$$\int_{\mathbb{R}} f_j(t) dt \leq \mathcal{W}([- \lambda t_j, \lambda t_j] v_j) = (1 + 2\lambda t_j).$$

Therefore, by (3.9) and the Brascamp–Lieb inequality (Theorem 2.1),

$$\begin{aligned} \int_{F_0} e^{-\pi \sum_{j \in J_0} \delta_j^0 d((x, v_j^0) v_j^0, P_{\langle v_j^0 \rangle}(\lambda(C_0 \cap F_0)))^2} &= \int_{F_0} \prod_{j \in J_0} f_j^{\delta_j^0}(\langle x, v_j^0 \rangle) dx \\ &\leq \prod_{j \in J_0} \left( \int_{\mathbb{R}} f_j(t) dt \right)^{\delta_j^0} \leq \prod_{j \in J_0} (1 + 2\lambda t_j)^{\delta_j^0}. \end{aligned}$$

By the arithmetic–geometric mean inequality, (2.1), and (3.9), we have

$$\begin{aligned}
 \prod_{j \in J_0} (1 + 2\lambda t_j)^{\frac{\delta_j^0}{k}} &\leq \sum_{j \in J_0} \frac{\delta_j^0}{k} (1 + 2\lambda t_j) \leq 1 + \frac{2\lambda}{k} \sum_{j \in J_0} c_j \|P_{F_0} u_j\|_2 \\
 &\leq 1 + \frac{2\lambda}{k} \left( \sum_{j \in J_0} c_j \right)^{\frac{1}{2}} \left( \sum_{j \in J_0} c_j \|P_{F_0} u_j\|_2^2 \right)^{\frac{1}{2}} \\
 &\leq 1 + \frac{2\lambda}{k} \left( \sum_{j=1}^m c_j \right)^{\frac{1}{2}} \left( \sum_{j \in J_0} c_j \|P_{F_0} u_j\|_2^2 \right)^{\frac{1}{2}} \\
 &= 1 + 2\lambda \sqrt{\frac{n}{k}}.
 \end{aligned}$$

It follows that

$$\prod_{j \in J_0} (1 + 2\lambda t_j)^{\delta_j^0} \leq \left( 1 + 2\lambda \sqrt{\frac{n}{k}} \right)^k = \mathcal{W} \left( \lambda \sqrt{\frac{n}{k}} B_\infty^k \right).$$

On the other hand, let  $x_0 \in \lambda(C_0 \cap F_0)$ . Then, for every  $x \in F_0$  and every  $j \in J_0$ ,

$$d \left( \langle x, v_j^0 \rangle v_j^0, P_{\langle v_j^0 \rangle}(\lambda(C_0 \cap F_0)) \right)^2 \leq d \left( \langle x, v_j^0 \rangle v_j^0, \langle x_0, v_j^0 \rangle v_j^0 \right)^2 = \langle x - x_0, v_j \rangle^2.$$

Thus, for every  $x_0 \in \lambda(C_0 \cap F_0)$  and every  $x \in F_0$ ,

$$\sum_{j \in J_0} \delta_j^0 d \left( \langle x, v_j^0 \rangle v_j^0, P_{\langle v_j^0 \rangle}(\lambda(C_0 \cap F_0)) \right)^2 \leq \sum_{j \in J_0} \delta_j^0 \langle x - x_0, v_j \rangle^2 = |x - x_0|^2.$$

Hence, for every  $x \in F_0$ ,

$$\sum_{j \in J_0} \delta_j^0 d \left( \langle x, v_j^0 \rangle v_j^0, P_{\langle v_j^0 \rangle}(\lambda(C_0 \cap F_0)) \right)^2 \leq d(x, \lambda(C_0 \cap F_0))^2.$$

Consequently,

$$\begin{aligned}
 \mathcal{W}(\lambda(C_0 \cap F_0)) &= \int_F e^{-\pi d(x, \lambda(C_0 \cap F_0))^2} dx \\
 &\leq \int_F e^{-\pi \sum_{j \in J_0} \delta_j^0 d \left( \langle x, v_j^0 \rangle v_j^0, P_{\langle v_j^0 \rangle}(\lambda(C_0 \cap F_0)) \right)^2} dx \leq \mathcal{W} \left( \lambda \sqrt{\frac{n}{k}} B_\infty^k \right).
 \end{aligned}$$

Since  $K \cap F_0 \subseteq C_0 \cap F_0$  and by the monotonicity of the Wills functional, we get

$$\mathcal{W}(\lambda(K \cap F_0)) \leq \mathcal{W}(\lambda(C_0 \cap F_0)) \leq \mathcal{W}\left(\lambda\sqrt{\frac{n}{k}}B_\infty^k\right). \quad \blacksquare$$

The following result gives a similar upper bound for a quantity defined via a double polarity, both on the convex body and on the log-concave function. For any  $k$  dimensional affine subspace  $F_h$  and any convex body  $K \subseteq F_h$ , we will consider

$$f_K(x) = e^{-\pi d^2(x,K)} \quad \forall x \in F_h,$$

as defined in Section 2.6.

**Theorem 8.1.** Let  $K \subseteq \mathbb{R}^n$  be a convex body in John's position and let  $F_h$  be a  $k$ -dimensional affine subspace at distance  $h$  from 0. Assume that the closest point to the origin in  $F_h$  belongs to the relative interior to  $K \cap F_h$ . Then, for every  $\lambda > 0$ ,

$$\int_{F_h} f_{(\lambda(K \cap F_h))^\circ}(x) dx \leq \frac{n+1}{k+1} \sqrt{\frac{n}{n+h^2}} \int_{\mathbb{R}^k} f_{\left(\lambda\sqrt{\frac{n(n+1)}{k(k+1)}}S_k\right)^\circ}(x) dx.$$

Here we consider  $K \cap F_h$  as a subset of  $F_h$ , which we identify with  $\mathbb{R}^k$  and the polarity is taken on  $F_h$ .

Furthermore, if  $K$  is centrally symmetric and  $F_0 \in G_{n,k}$  is a  $k$ -dimensional linear subspace then, for every  $\lambda > 0$ ,

$$\int_{F_0} f_{(\lambda(K \cap F_0))^\circ}(x) dx \leq \int_{\mathbb{R}^k} f_{\left(\lambda\sqrt{\frac{n}{k}}B_\infty^k\right)^\circ}(x) dx.$$

**Proof.** Let  $K \subseteq \mathbb{R}^n$  be a centrally symmetric convex body in John's position and let  $F_0 \in G_{n,k}$  be a  $k$ -dimensional linear subspace. From the definition of  $f_{(\lambda(K \cap F_0))^\circ}$  and (2.4), we have that, for every  $\lambda > 0$ ,

$$\begin{aligned} \int_{F_0} f_{(\lambda(K \cap F_0))^\circ}(x) dx &= \int_{F_0} e^{-\frac{\|x\|_2^2}{4\pi}} e^{-\|x\|_{\lambda(K \cap F_0)}} dx = \int_{F_0} e^{-\frac{\|x\|_2^2}{4\pi}} \int_{\|x\|_{\lambda(K \cap F_0)}}^\infty e^{-t} dt dx \\ &= \int_0^\infty e^{-t} \int_{t\lambda(K \cap F_0)} e^{-\frac{\|x\|_2^2}{4\pi}} dt dx \\ &= (2\pi)^k \int_0^\infty e^{-t} \int_{\frac{t\lambda}{\sqrt{2\pi}}(K \cap F_0)} \frac{e^{-\frac{\|x\|_2^2}{2}}}{(\sqrt{2\pi})^k} dx dt \\ &= (2\pi)^k \int_0^\infty e^{-t} \gamma_k\left(\frac{t\lambda}{\sqrt{2\pi}}(K \cap F_0)\right) dt. \end{aligned}$$

Similarly, for every  $\lambda > 0$ ,

$$\int_{\mathbb{R}^k} f_{(\lambda\sqrt{\frac{n}{k}}B_\infty^k)}^\circ(x) dx = (2\pi)^k \int_0^\infty e^{-t} \gamma_k \left( \frac{t\lambda}{\sqrt{2\pi}} \sqrt{\frac{n}{k}} B_\infty^k \right) dt.$$

As we have seen in the proof of Theorem 1.4, for every  $t \geq 0$  and every  $\lambda > 0$ ,

$$\gamma_k \left( \frac{t\lambda}{\sqrt{2\pi}} (K \cap F_0) \right) \leq \gamma_k \left( \frac{t\lambda}{\sqrt{2\pi}} \sqrt{\frac{n}{k}} B_\infty^k \right).$$

Therefore, for every  $\lambda > 0$ ,

$$\int_{F_0} f_{(\lambda(K \cap F_0))^\circ}^\circ(x) dx \leq \int_{\mathbb{R}^k} f_{(\lambda\sqrt{\frac{n}{k}}B_\infty^k)}^\circ(x) dx.$$

Assume now that  $K \subseteq \mathbb{R}^n$  is a (not necessarily symmetric) convex body in John's position,  $F_h$  is a  $k$ -dimensional affine subspace at distance  $h$  from the origin, and the closest point to the origin in  $F_h$  belongs to the relative interior to  $K \cap F_h$ . We will identify  $F_h$  with  $\mathbb{R}^k$  and the closest point in  $F_h$  to the origin in  $\mathbb{R}^n$  with the origin in  $F_h$  (identified with  $\mathbb{R}^k$ ). As before, we have that, for every  $\lambda > 0$ ,

$$\int_{F_h} f_{(\lambda(K \cap F_h))^\circ}^\circ(x) dx = (2\pi)^k \int_0^\infty e^{-t} \gamma_k \left( \frac{t\lambda}{\sqrt{2\pi}} (K \cap F_h) \right) dt.$$

We will follow the notation in Section 3.1. By Lemma 7.1, we have that for any  $\alpha \in \mathbb{R}$  and any  $\beta \leq 0$ ,

$$\begin{aligned} \sqrt{\frac{n+h^2}{n}} \int_0^\infty \frac{e^{-\frac{(r-\alpha\sqrt{n+1})^2}{2}}}{\sqrt{2\pi}} e^{\beta r\sqrt{n+1}} \gamma_k \left( \frac{r}{\sqrt{n}} (C \cap F_h) \right) dr &\leq \\ \int_0^\infty \frac{e^{-\frac{(r-\alpha d_1)^2}{2}}}{\sqrt{2\pi}} e^{\beta r\sqrt{k+1}} \gamma_k \left( r\sqrt{k+1} \Delta_k \right) dr, \end{aligned}$$

where  $d_1 = \frac{1}{\sqrt{k+1}} \sum_{j \in J} \delta_j \|P_H v_j\|_2$ . Integrating with respect to  $\alpha \in \mathbb{R}$ , we see that for any  $\beta \leq 0$ ,

$$\sqrt{\frac{n+h^2}{n(n+1)}} \int_0^\infty e^{\beta r\sqrt{n+1}} \gamma_k \left( \frac{r}{\sqrt{n}} (C \cap F_h) \right) dr \leq \frac{1}{d_1} \int_0^\infty e^{\beta r\sqrt{k+1}} \gamma_k \left( r\sqrt{k+1} \Delta_k \right) dr.$$

Equivalently, changing variables  $u = \sqrt{\frac{n}{2\pi}}r$ , for any  $\beta \leq 0$ ,

$$\sqrt{\frac{n+h^2}{n+1}} \int_0^\infty e^{\frac{\beta u \sqrt{n(n+1)}}{\sqrt{2\pi}}} \gamma_k \left( \frac{u}{\sqrt{2\pi}} (C \cap F_h) \right) du \leq \frac{1}{d_1 \sqrt{k+1}} \int_0^\infty e^{\frac{\beta u}{\sqrt{2\pi}}} \gamma_k \left( \frac{u}{\sqrt{2\pi}} \Delta_k \right) du.$$

For any  $\lambda > 0$ , take  $\beta = -\frac{1}{\lambda} \sqrt{\frac{2\pi}{n(n+1)}}$  to obtain

$$\sqrt{\frac{n+h^2}{n+1}} \int_0^\infty e^{-\frac{u}{\lambda}} \gamma_k \left( \frac{u}{\sqrt{2\pi}} (C \cap F_h) \right) du \leq \frac{1}{d_1 \sqrt{k+1}} \int_0^\infty e^{\frac{-u}{\lambda \sqrt{n(n+1)}}} \gamma_k \left( \frac{u}{\sqrt{2\pi}} \Delta_k \right) du,$$

or, equivalently, changing variables  $u = \lambda v$  in the integral on the left-hand side and  $u = \lambda \sqrt{n(n+1)}v$  in the integral on the right-hand side and renaming  $v$  as  $u$ ,

$$\sqrt{\frac{n+h^2}{n+1}} \int_0^\infty e^{-u} \gamma_k \left( \frac{u\lambda}{\sqrt{2\pi}} (C \cap F_h) \right) du \leq \frac{\sqrt{n(n+1)}}{d_1 s \sqrt{k+1}} \int_0^\infty e^{-u} \gamma_k \left( \frac{u\lambda \sqrt{n(n+1)}}{\sqrt{2\pi}} \Delta_k \right) du.$$

Since  $\sqrt{k(k+1)}\Delta_k = S_k$ , we see that for every  $\lambda > 0$ ,

$$\int_0^\infty e^{-u} \gamma_k \left( \frac{u\lambda}{\sqrt{2\pi}} (C \cap F_h) \right) du \leq \frac{(n+1)\sqrt{n}}{d_1 \sqrt{(k+1)(n+h^2)}} \int_0^\infty e^{-u} \gamma_k \left( \frac{u\lambda}{\sqrt{2\pi}} \sqrt{\frac{n(n+1)}{k(k+1)}} S_k \right) du.$$

Consequently, for any  $\lambda > 0$ , taking polars with respect to the closest point in  $F_h$  to the origin that we assumed to belong to the relative interior of  $K \cap F_h$ , we get

$$\begin{aligned} \int_{F_h} f_{(\lambda(K \cap F_h))^\circ}^\circ(x) dx &\leq \int_{F_h} f_{(\lambda(C \cap F_h))^\circ}^\circ(x) dx \\ &\leq \frac{\sqrt{n(n+1)}}{d_1 \sqrt{(k+1)(n+h^2)}} \int_{\mathbb{R}^k} f_{(\lambda \sqrt{\frac{n(n+1)}{k(k+1)}} S_k)^\circ}^\circ(x) dx. \end{aligned}$$

For every  $j \in J$ , denote  $\kappa_j$  as in (3.6) and  $s_j$  as in (3.7). Then, by (3.6), we have that

$$d_1 \sqrt{k+1} = \sum_{j \in J} \kappa_j s_j \geq \sum_{j \in J} \kappa_j = \sum_{j=1}^m = k+1.$$

Thus, for every  $\lambda > 0$ ,

$$\int_{F_h} f_{(\lambda(K \cap F_h))^\circ}^\circ(x) dx \leq \frac{n+1}{k+1} \sqrt{\frac{n}{n+h^2}} \int_{\mathbb{R}^k} f_{(\lambda \sqrt{\frac{n(n+1)}{k(k+1)}} S_k)^\circ}^\circ(x) dx. \quad \blacksquare$$

## 9 The Wills Functional of Projections of Convex Bodies in Löwner's Position

In this section, we will give the proof of Theorem 1.6.

**Proof of Theorem 1.6.** Let  $K$  be a centrally symmetric convex body in John's position and let  $F_0 \in G_{n,k}$  be a  $k$ -dimensional linear subspace. We follow the notation in Section 3.2. By (3.11), we have that

$$(K \cap F_0)^\circ \supseteq (C_0 \cap F_0)^\circ = \text{conv}\{\pm P_{F_0} u_j, j \in J_0\} = \text{conv}\{\pm \|P_{F_0} u_j\|_2 v_j, j \in J_0\}.$$

Since the function  $d(\cdot, (C_0 \cap F_0)^\circ)^2$  is convex, for any  $x \in F_0$  and any  $\{\theta_j\}_{j \in J_0} \subseteq \mathbb{R}$  such that  $x = \sum_{j \in J_0} \delta_j^0 \theta_j v_j^0$ , we have that

$$\begin{aligned} d(x, (K \cap F_0)^\circ)^2 &\leq d(x, (C_0 \cap F_0)^\circ)^2 = d\left(\sum_{j \in J_0} \frac{\delta_j^0}{k} k \theta_j v_j^0, (C_0 \cap F_0)^\circ\right)^2 \\ &\leq \frac{1}{k} \sum_{j \in J_0} \delta_j^0 d\left(k \theta_j v_j^0, (C_0 \cap F_0)^\circ\right)^2 \\ &\leq \frac{1}{k} \sum_{j \in J_0} \delta_j^0 d\left(k \theta_j v_j^0, \left[-\|P_{F_0} u_j\|_2 v_j^0, \|P_{F_0} u_j\|_2 v_j^0\right]\right)^2 \\ &= \sum_{j \in J_0} \delta_j^0 d\left(\sqrt{k} \theta_j v_j^0, \left[-\frac{\|P_{F_0} u_j\|_2}{\sqrt{k}} v_j^0, \frac{\|P_{F_0} u_j\|_2}{\sqrt{k}} v_j^0\right]\right)^2. \end{aligned}$$

For every  $j \in J_0$ , we set

$$f_j(t) = e^{-\pi d\left(\sqrt{k} t v_j^0, \left[-\frac{\|P_{F_0} u_j\|_2}{\sqrt{k}} v_j^0, \frac{\|P_{F_0} u_j\|_2}{\sqrt{k}} v_j^0\right]\right)^2}, \quad \forall t \in \mathbb{R}.$$

Moreover, for any  $x \in F_0$  and any  $\{\theta_j\}_{j \in J_0} \subseteq \mathbb{R}$  such that  $x = \sum_{j \in J_0} \delta_j^0 \theta_j v_j^0$ , we have

$$\prod_{j \in J_0} f_j^{\delta_j^0}(\theta_j) \leq e^{-\pi d(x, (K \cap F_0)^\circ)^2}.$$

Therefore, by (3.9) and the reverse Brascamp–Lieb inequality (Theorem 2.1),

$$\begin{aligned}
 \mathcal{W}((K \cap F_0)^\circ) &= \int_{F_0} e^{-\pi d(x, (K \cap F_0)^\circ)^2} dx \geq \prod_{j \in J_0} \left( \int_{\mathbb{R}} f_j(t) dt \right)^{\delta_j^0} \\
 &= \prod_{j \in J_0} \left( \frac{1}{\sqrt{k}} \int_{\mathbb{R}} e^{-\pi d\left( tv_j^0, \left[ -\frac{\|P_{F_0} u_j\|_2}{\sqrt{k}} v_j^0, \frac{\|P_{F_0} u_j\|_2}{\sqrt{k}} v_j^0 \right] \right)^2} dt \right)^{\delta_j^0} \\
 &= \frac{1}{k^{k/2}} \prod_{j \in J_0} \mathcal{W} \left( \left[ -\frac{\|P_{F_0} u_j\|_2}{\sqrt{k}} v_j^0, \frac{\|P_{F_0} u_j\|_2}{\sqrt{k}} v_j^0 \right] \right)^{\delta_j^0} \\
 &= \frac{1}{k^{k/2}} \prod_{j \in J_0} \left( 1 + \frac{2\|P_{F_0} u_j\|_2}{\sqrt{k}} \right)^{\delta_j^0} \geq \frac{1}{k^{k/2}}.
 \end{aligned}$$

■

### 10 Sections of Convex Bodies in Minimal Surface Area Position

In this section, we are going to prove Theorem 1.7. Let us start assuming that  $K$  is a centrally symmetric polytope in minimal surface area position and  $F_0 \in G_{n,k}$ . By an approximation argument, the inequalities we obtain will also be true for any centrally symmetric convex body in minimal surface area position. We will follow the notation introduced in Section 3.3.

Let  $J_0$  and, for every  $j \in J_0$ ,  $c_j$ ,  $\delta_j^0$ , and  $v_j^0$  be as in (3.14). Let for every  $j \in J_0$ ,  $f_j : \mathbb{R} \rightarrow [0, \infty)$  be the function

$$f_j(t) = e^{-\pi d(tv_j^0, P_{\langle v_j^0 \rangle}(K \cap F_0))^2} \quad \forall t \in \mathbb{R},$$

where  $\langle v_j^0 \rangle$  denotes the 1-dimensional subspace spanned by  $v_j^0$ . Notice that, for every  $j \in J_0$ ,

$$\int_{\mathbb{R}} f_j(t) dt = \int_{\mathbb{R}} e^{-\pi d(tv_j^0, P_{\langle v_j^0 \rangle}(K \cap F_0))^2} dt = \mathcal{W}(P_{\langle v_j^0 \rangle}(K \cap F_0)).$$

For every  $j \in J_0$ , we have that

$$P_{\langle v_j^0 \rangle}(K \cap F_0) \subseteq \left[ -t_j h_K(u_j), t_j h_K(u_j) \right] v_j^0,$$

where  $t_j$  is defined as in (3.15). Then, for every  $j \in J_0$ ,

$$\int_{\mathbb{R}} f_j(t) dt \leq \mathcal{W} \left( [-t_j h_K(u_j), t_j h_K(u_j)] v_j \right) = (1 + 2t_j h_K(u_j)).$$

Therefore, by (3.14) and the Brascamp–Lieb inequality (Theorem 2.1),

$$\begin{aligned} \int_{F_0} e^{-\pi \sum_{j \in J_0} \delta_j^0 d \left( \langle x, v_j^0 \rangle v_j^0, P_{(v_j^0)}(C_0 \cap F_0) \right)^2} dx &= \int_{F_0} \prod_{j \in J_0} f_j^{\delta_j^0}(\langle x, v_j^0 \rangle) dx \\ &\leq \prod_{j \in J_0} \left( \int_{\mathbb{R}} f_j(t) dt \right)^{\delta_j^0} \leq \prod_{j \in J_0} (1 + 2t_j h_K(u_j))^{\delta_j^0}. \end{aligned}$$

By the arithmetic–geometric mean inequality and (3.13), we have

$$\begin{aligned} \prod_{j \in J_0} (1 + 2t_j h_K(u_j))^{\frac{\delta_j^0}{k}} &\leq \sum_{j \in J_0} \frac{\delta_j^0}{k} (1 + 2t_j h_K(u_j)) \\ &= 1 + \frac{2}{k} \sum_{j \in J_0} \frac{n |F_j| \|P_{F_0} u_j\|_2 h_K(u_j)}{|\partial K|} \leq 1 + \frac{2n}{k |\partial K|} \sum_{j \in J_0} |F_j| h_K(u_j) \\ &\leq 1 + \frac{2n}{k |\partial K|} \sum_{j=1}^m |F_j| h_K(u_j) = 1 + 2 \frac{n^2 |K|}{k |\partial K|}. \end{aligned}$$

Thus,

$$\prod_{j \in J_0} (1 + 2t_j h_K(u_j))^{\delta_j^0} \leq \left( 1 + 2 \frac{n^2 |K|}{k |\partial K|} \right)^k = \mathcal{W} \left( \frac{n^2 |K|}{k |\partial K|} B_{\infty}^k \right).$$

Let  $x_0 \in K \cap F_0$ . For every  $x \in F_0$  and every  $j \in J_0$ , we have

$$d \left( \langle x, v_j^0 \rangle v_j^0, P_{(v_j^0)}(K \cap F_0) \right)^2 \leq d \left( \langle x, v_j^0 \rangle v_j^0, \langle x_0, v_j^0 \rangle v_j^0 \right)^2 = \langle x - x_0, v_j \rangle^2.$$

Thus, for every  $x_0 \in C_0 \cap F_0$  and every  $x \in F_0$ ,

$$\sum_{j \in J_0} \delta_j^0 d \left( \langle x, v_j^0 \rangle v_j^0, P_{(v_j^0)}(K \cap F_0) \right)^2 \leq \sum_{j \in J_0} \delta_j^0 \langle x - x_0, v_j^0 \rangle^2 = |x - x_0|^2.$$



Hence, for every  $x \in F_0$ ,

$$\sum_{j \in J_0} \delta_j^0 d \left( \langle x, v_j^0 \rangle v_j^0, P_{\langle v_j^0 \rangle} (K \cap F_0) \right)^2 \leq d(x, K \cap F_0)^2.$$

Consequently,

$$\begin{aligned} \mathcal{W}(K \cap F_0) &= \int_{F_0} e^{-\pi d(x, K \cap F_0)^2} dx \leq \int_{F_0} e^{-\pi \sum_{j \in J_0} \delta_j^0 d \left( \langle x, v_j^0 \rangle v_j^0, P_{\langle v_j^0 \rangle} (K \cap F_0) \right)^2} dx \\ &\leq \mathcal{W} \left( \frac{n^2 |K|}{k |\partial K|} B_\infty^k \right), \end{aligned}$$

which proves (i).

Notice that for every  $\lambda \geq 0$  we have that  $\lambda K$  is in minimal surface area position. Therefore, we can use (i) to get that for every  $\lambda \geq 0$

$$\mathcal{W}(\lambda(K \cap F_0)) \leq \mathcal{W} \left( \lambda \frac{n^2 |K|}{k |\partial K|} B_\infty^k \right).$$

As explained in Section 2.6, the last one implies that  $V_1(K \cap F_0) \leq V_1 \left( \lambda \frac{n^2 |K|}{k |\partial K|} B_\infty^k \right)$  and  $V_n((K \cap F_0) \leq V_n \left( \lambda \frac{n^2 |K|}{k |\partial K|} B_\infty^k \right)$ , which are equivalent to (ii) and (iii), respectively.

Now, using (3.16), we observe that for every  $x \in F_0$

$$\begin{aligned} h_{K \cap F_0}(x) &= \inf \left\{ \sum_{j \in J_0} |\alpha_j| : x = \sum_{j \in J_0} \frac{\alpha_j}{h_K(u_j)} P_{F_0} u_j \right\} \\ &= \inf \left\{ \sum_{j \in J_0} |\beta_j| t_j h_K(u_j) : x = \sum_{j \in J_0} \beta_j v_j^0 \right\}, \end{aligned}$$

where  $t_j = \frac{1}{\|P_{F_0} u_j\|_2} = \left( \frac{c_j}{\delta_j^0} \right)^{\frac{1}{2}}$  for all  $j \in J_0$ . For every  $j \in J_0$ , we define

$$f_j(t) := e^{-|t| t_j h_K(u_j)}, \quad t \in \mathbb{R}.$$

Then, if  $x = \sum_{j \in J_0} \delta_j^0 \theta_j v_j^0$  for some  $\{\theta_j\}_{j \in J_0} \subseteq \mathbb{R}$ , we have

$$\prod_{j \in J_0} f_j^{\delta_j^0}(\theta_j) = e^{-\sum_{j \in J_0} \delta_j |\theta_j| t_j h_K(u_j)} \leq e^{-h_{K \cap F_0}(x)}.$$

Therefore, by (3.14) and the reverse Brascamp–Lieb inequality (Theorem 2.1),

$$\begin{aligned} k! |(K \cap F_0)^\circ| &= \int_{F_0} e^{-h_{K \cap F_0}(x)} dx \geq \prod_{j \in J} \left( \int_{\mathbb{R}} e^{-|t|t_j h_K(u_j)} dt \right)^{\delta_j} \\ &= \frac{2^k}{\prod_{j \in J_0} (t_j h_K(u_j))^{\delta_j}}. \end{aligned}$$

By the arithmetic–geometric mean inequality and (3.13),

$$\begin{aligned} \prod_{j \in J_0} (t_j h_K(u_j))^{\frac{\delta_j^0}{k}} &\leq \sum_{j \in J_0} \frac{\delta_j^0}{k} t_j h_K(u_j) = \sum_{j \in J_0} \frac{n |F_j| \|P_{F_0} u_j\|_2 h_K(u_j)}{k |\partial K|} \\ &\leq \frac{n}{k |\partial K|} \sum_{j=1}^m |F_j| h_K(u_j) = \frac{n^2 |K|}{k |\partial K|}. \end{aligned}$$

Taking into account that  $|(B_\infty^k)^\circ| = |B_1^k| = \frac{2^k}{k!}$ , we obtain

$$|(K \cap F_0)^\circ|^{1/k} \geq \frac{k |\partial K|}{n^2 |K|} |(B_\infty^k)^\circ|^{1/k},$$

which gives us (iv).

Finally, from (3.15), observe that for every  $t \geq 0$

$$t(K \cap F_0) = \{x \in F_0 : |\langle x, v_j^0 \rangle| \leq t t_j h_K(u_j), \forall j \in J_0\}.$$

By (3.14) and the Brascamp–Lieb inequality (Theorem 2.1), we have that

$$\begin{aligned} \gamma_k(t(K \cap F_0)) &= \int_{\mathbb{R}^n} \prod_{j \in J_0} \chi_{[-t t_j h_K(u_j), t t_j h_K(u_j)]}(\langle x, v_j^0 \rangle) \frac{e^{-\sum_{j \in J_0} \frac{\delta_j^0 \langle x, v_j^0 \rangle^2}{2}}}{(2\pi)^{k/2}} dx \\ &\leq \prod_{j \in J_0} \left( \int_{-t t_j h_K(u_j)}^{t t_j h_K(u_j)} \frac{e^{-\frac{s^2}{2}}}{\sqrt{2\pi}} ds \right)^{\delta_j^0} = \prod_{j \in J_0} \gamma_1 \left( [-t t_j h_K(u_j), t t_j h_K(u_j)] \right)^{\delta_j^0}. \end{aligned}$$

Since the 1-dimensional Gaussian measure is log-concave, we obtain that

$$\begin{aligned} \prod_{j \in J_0} \gamma_1 \left( \left[ -tt_j h_K(u_j), tt_j h_K(u_j) \right] \right)^{\delta_j^0} &\leq \gamma_1 \left( \left( \sum_{j \in J_0} \frac{\delta_j^0 tt_j h_K(u_j)}{k} \right) [-e_1, e_1] \right)^k \\ &= \gamma_k \left( t \left( \sum_{j \in J_0} \frac{\delta_j^0 t_j h_K(u_j)}{k} \right) B_\infty^k \right). \end{aligned}$$

Therefore, by (3.13),

$$\begin{aligned} \sum_{j \in J_0} \frac{\delta_j^0 t_j h_K(u_j)}{k} &= \sum_{j \in J_0} \frac{n|F_j| \|P_{F_0} u_j\|_2 h_K(u_j)}{k|\partial K|} \leq \sum_{j \in J_0} \frac{n|F_j| h_K(u_j)}{k|\partial K|} \\ &\leq \sum_{j=1}^m \frac{n|F_j| h_K(u_j)}{k|\partial K|} = \frac{n^2}{k} \frac{|K|}{|\partial K|}. \end{aligned}$$

Thus, for any  $t \geq 0$ ,

$$\gamma_k(t(K \cap F_0)) \leq \gamma_k \left( t \frac{n^2}{k} \frac{|K|}{|\partial K|} B_\infty^k \right).$$

Therefore,

$$w((K \cap F_0)^\circ) \geq w \left( \left( \frac{n^2}{k} \frac{|K|}{|\partial K|} B_\infty^k \right)^\circ \right) = \frac{k}{n^2} \frac{|\partial K|}{|K|} w((B_\infty^k)^\circ),$$

and we obtain (v).

Let us now assume that  $K$  is a (not necessarily symmetric) polytope in minimal surface area position and  $F_0 \in G_{n,k}$ . Again, by approximation, the inequalities we obtain will be true for any convex body. By (2.5), we have that for any  $x \in F_0$ ,

$$\begin{aligned} \|x\|_{\Pi^* K \cap F_0} &= \frac{1}{2} \sum_{j=1}^m |F_j| |\langle x, u_j \rangle| = \frac{1}{2} \sum_{j \in J_0} |F_j| \|P_{F_0} u_j\|_2 \left| \langle x, v_j^0 \rangle \right| \\ &= \sum_{j \in J_0} \frac{|\partial K| \delta_j^0 t_j}{2n} \left| \langle x, v_j^0 \rangle \right|, \end{aligned}$$

where the vectors  $u_j$  are defined as in (3.12) and, for every  $j \in J_0$ ,  $\delta_j^0$  and  $v_j^0$  are defined as in (3.14) and  $t_j$  is defined as in (3.15). Therefore, by (3.14) and the Brascamp–Lieb

inequality (Theorem 2.1),

$$\begin{aligned}
 k! |\Pi^*K \cap F_0| &= \int_{F_0} e^{-\|x\|_{\Pi^*K \cap F_0}} dx = \int_{F_0} e^{-\sum_{j \in J_0} \frac{|\partial K| \delta_j^0 t_j}{2n} |\langle x, v_j^0 \rangle|} dx \\
 &= \int_{F_0} \prod_{j \in J_0} \left( e^{-\frac{|\partial K| t_j}{2n} |\langle x, v_j^0 \rangle|} \right)^{\delta_j^0} dx \\
 &\leq \prod_{j \in J_0} \left( \int_{\mathbb{R}} e^{-\frac{|\partial K| t_j}{2n} t} dt \right)^{\delta_j^0} = \prod_{j \in J_0} \left( \frac{4n}{|\partial K| t_j} \right)^{\delta_j^0} \\
 &= \left( \frac{4n}{|\partial K|} \right)^k \prod_{j \in J_0} (\|P_{F_0} u_j\|)^{\delta_j^0} \leq \left( \frac{4n}{|\partial K|} \right)^k.
 \end{aligned}$$

To prove the remaining inequality, we start observing that  $P_{F_0} \Pi K = (\Pi^*K \cap F_0)^\circ$ . Then, for every  $x \in F_0$ ,

$$\|x\|_{P_{F_0} \Pi K} = \inf \left\{ \max_{j \in J_0} |\tau_j| : x = \sum_{j \in J_0} \frac{|\partial K| \delta_j^0 t_j \tau_j}{2n} v_j^0 \right\}.$$

Any decomposition of  $x$  of the form  $x = \sum_{j \in J_0} \delta_j^0 \theta_j v_j^0$  with  $|\theta_j| \leq \frac{|\partial K| t_j}{2n}$  gives a decomposition of  $x$  of the form

$$x = \sum_{j \in J_0} \frac{|\partial K| \delta_j^0 t_j \tau_j}{2n} v_j^0 \quad \text{with} \quad \tau_j = \frac{2n \theta_j}{|\partial K| t_j}.$$

Since  $\max_{j \in J_0} |\tau_j| \leq 1$ , we get that the functions  $h_j = \chi_{\left[-\frac{|\partial K| t_j}{2n}, \frac{|\partial K| t_j}{2n}\right]}$ ,  $j \in J_0$ , and the function  $h = \chi_{P_{F_0} \Pi K}$  have the property that

$$h \left( \sum_{j \in J_0} \delta_j^0 \theta_j v_j^0 \right) \geq \prod_{j \in J_0} h_j^{\delta_j^0}(\theta_j),$$

for every  $\{\theta_j\}_{j \in J_0} \subseteq \mathbb{R}$ . Hence, by (3.14) and the reverse Brascamp–Lieb inequality (Theorem 2.1),

$$\begin{aligned}
 |P_{F_0} \Pi K| &= \int_{F_0} h(x) dx \geq \prod_{j \in J_0} \left( \int_{\mathbb{R}} h_j(t) dt \right)^{\delta_j^0} = \prod_{j \in J_0} \left( \frac{|\partial K| t_j}{n} \right)^{\delta_j^0} \\
 &= \left( \frac{|\partial K|}{n} \right)^k \prod_{j \in J_0} \left( \frac{1}{\|P_{F_0} u_j\|} \right)^{\delta_j^0} \geq \left( \frac{|\partial K|}{n} \right)^k.
 \end{aligned}$$

## Funding

This work was partially supported by the Ministerio de Ciencia e Innovacion of Spain [Project PID-105979-GB-I00 to D. A.-G.]; the Diputación General de Aragón [Project E48\_20R to D. A.-G.]; and the Hellenic Foundation for Research and Innovation [Project Number 1849 to S.B.].

## Acknowledgments

We would like to thank the anonymous referee for his/her careful reading of the paper and for his/her suggestions, which were very helpful in order to improve the presentation of the paper.

## References

- [1] Alonso-Gutiérrez, D. "Volume estimates for  $L_p$ -zonotopes and best constants in Brascamp–Lieb inequalities." *Mathematika* 56 (2010): 45–60.
- [2] Alonso-Gutiérrez, D., M. A. Hernández Cifre, and J. Yepes Nicolás. "Further inequalities for the (generalized) Will's functional." *Commun. Contemp. Math.* 23, no. 3 (2021): 2050011.
- [3] Artstein-Avidan, S., A. Giannopoulos, and V. D. Milman. "Asymptotic Geometric Analysis, Part I." In *Mathematical Surveys and Monographs*, vol. 202. Providence, Rhode Island: AMS, 2015. ISBN 978-1-4704-2193-9.
- [4] Aubrun, G. and S. J. Szarek. "Alice and Bob Meet Banach: The Interface of Asymptotic Geometric Analysis and Quantum Information Theory." In *Mathematical Surveys and Monographs*, vol. 223. Providence, Rhode Island: AMS, 2017. ISBN: 978-1-4704-3468-7.
- [5] Ball, K. M. "Volumes of Sections of Cubes and Related Problems." In *Lecture Notes in Mathematics*, vol. 1376. 251–60. Berlin: Springer, 1989.
- [6] Ball, K. M. "Volume ratios and a reverse isoperimetric inequality." *J. London Math. Soc.* 44, no. 2 (1991): 351–9.
- [7] Ball, K. M. "Ellipsoids of maximal volume in convex bodies." *Geom. Dedicata* 41 (1992): 241–50.
- [8] Ball, K. M. "Mahler's conjecture and wavelets." *Discrete Comput. Geom.* 13, no. 3–4 (1995): 271–7.
- [9] Barthe, F. "Inégalités de Brascamp–Lieb et convexité." *C. R. Acad. Sci. Paris* 324 (1997): 885–8.
- [10] Barthe, F. "An extremal property of the mean width of the simplex." *Math. Ann.* 310 (1998): 685–93.
- [11] Barthe, F., O. Guédon, S. Mendelson, and A. Naor. "A probabilistic approach to the geometry of the  $\ell_p^N$ -ball." *Ann. Probab.* 33, no. 2 (2005): 480–513.
- [12] Barthe, F. and A. Naor. "Hyperplane projections of the unit ball of  $\ell_p^N$ ." *Discrete Comput. Geom.* 27, no. 2 (2002): 215–26.
- [13] Brascamp, H. J. and E. H. Lieb. "Best constants in Young's inequality, its converse and its generalization to more than three functions." *Adv. Math.* 20 (1976): 151–73.
- [14] Brazitikos, S., A. Giannopoulos, P. Valettas, and B.-H. Vritsiou. "Geometry of Isotropic Convex Bodies." In *Mathematical Surveys and Monographs*, vol. 196. Providence, Rhode Island: AMS, 2014. ISBN 978-1-4704-1456-6.

- [15] Dirksen, H. "Sections of simplices and cylinders—volume formulas and estimates." PhD diss., Christian-Albrechts-Universität zu Kiel, 2015.
- [16] Dirksen, H. "Sections of the regular simplex—volume formulas and estimates." *Math. Nachr.* 290, no. 16 (2017): 2567–84.
- [17] Eszenakis, A., P. Nayar, and T. Tkocz. "Gaussian mixtures: entropy and geometric inequalities." *Ann. Probab.* 46, no. 5 (2018): 2908–45.
- [18] Eszenakis, A. "On extremal sections of subspaces of  $L_p$ ." *Discrete Comput. Geom.* 65 (2021): 489–509.
- [19] Filliman, P. "The extreme projections of the regular simplex." *Trans. AMS* 317, no. 2 (1990): 611–29.
- [20] Fuchs, L. "A new proof of an inequality of Hardy–Littlewood–Pólya." *Math. Tidsskr. B.* (1947): 53–4.
- [21] Giannopoulos, A. and M. Papadimitrakis. "Isotropic surface area measures." *Mathematika* 46 (1999): 1–13.
- [22] Giannopoulos, A., V. D. Milman, and M. Rudelson. "Convex Bodies With Minimal Mean Width." In *Geometric Aspects of Functional Analysis*. Lecture Notes in Mathematics, vol. 1745. 81–93. 2000.
- [23] Gordon, Y., M. Meyer, and S. Reisner. "Zonoids with minimal volume product: a new proof." *Proc. AMS* 104 (1988): 273–6.
- [24] Gruber, P. M. *Convex and Discrete Geometry*. Berlin Heidelberg: Springer, 2007.
- [25] Hadwiger, H. "Das Wills'sche Funktional." *Monatsh. Math.* 79 (1975): 213–21.
- [26] Ivanov, G. "On the volume of the John–Löwner ellipsoid." *Discrete Comput. Geom.* 63, no. 2 (2020): 455–9.
- [27] Ivanov, G. "On the volume of projections of the cross-polytope." *Discrete Math.* 344, no. 5 (2021): 112312.
- [28] Ivanov, G., and I. Tsiutsiurupa "On the volume of sections of the cube." *Anal. Geom. Metr. Spaces* 9, no. 1 (2021): 1–18.
- [29] John, F. "Extremum Problems With Inequalities as Subsidiary Conditions." In *Courant Anniversary Volume*. 187–204. New York: Inter-science, 1948.
- [30] König, H. and A. Koldobsky. "On the Maximal Measure of Sections of the n-Cube." *Geometric Analysis, Mathematical Relativity, and Nonlinear Partial Differential Equations*. Contemp. Math., vol. 599. 123–55. 2012.
- [31] Koldobsky, A. "An application of the Fourier transform to sections of star bodies." *Israel J. Math.* 106, no. 1 (1998): 157–64.
- [32] Markessinis, E. "Central sections of a convex body with ellipsoid of maximal volume  $B_2^n$ ." *Math. Ann.* 378 (2020): 233–41.
- [33] Markessinis, E., G. Paouris, and C. Saroglou. "Comparing the M-position with some classical positions of convex bodies." *Math. Proc. Cambridge Philos. Soc.* 152 (2012): 131–52.
- [34] McMullen, P. "Non-linear angle-sum relations for polyhedral cones and polytopes." *Math. Proc. Cambridge Philos. Soc.* 78 (1975): 247–61.

- [35] McMullen, P. "Inequalities between intrinsic volumes." *Monatsh. Math.* 111, no. 1 (1991): 47–53.
- [36] Petty, C. M. "Surface area of a convex body under affine transformations." *Proc. AMS* 12 (1961): 824–8.
- [37] Pisier, G. "The Volume of Convex Bodies and Banach Space Geometry." *Cambridge Tracts in Mathematics*, vol. 94. Cambridge Univ. Press, 1989. ISBN 978–0–521–36465–2.
- [38] Reisner, S. "Zonoids with minimal volume product." *Math. Z.* 192 (1986): 339–46.
- [39] Sangwine-Yager, J. R. "Mixed Volumes." In *Handbook of Convex Geometry*, edited by P. M. Gruber and J. M. Wills 43–71. North-Holland: Amsterdam, 1993.
- [40] Schechtman, G. and M. Schmuckenschläger. "A Concentration Inequality for Harmonic Measures on the Sphere." In *Geometric Aspects of Functional Analysis (Israel, 1992–1994)*, vol. 77. 255–73. Oper. Theory Adv. Appl. Basel: Birkhäuser, 1995.
- [41] Schmuckenschläger, M. "An Extremal Property of the Regular Simplex." In *Convex Geometric Analysis (Berkeley, CA, 1996)*, vol. 34. 199–202. Cambridge: MSRI Publications, Cambridge Univ. Press, 1999.
- [42] Schneider, R. "Convex Bodies: The Brunn–Minkowski Theory," 2nd expanded ed. In *Encyclopedia of Mathematics and its Applications*, vol. 151. Cambridge: Cambridge University Press, 2014.
- [43] Vaaler, J. D. "A Geometric inequality with applications to linear forms." *Pacific J. Math.* 83, no. 2 (1979): 543–53.
- [44] Webb, S. "Central slices of the regular simplex." *Geom. Dedicata* 61 (1996): 19–28.
- [45] Wills, J. M. "Zur Gitterpunktanzahl konvexer Mengen." *Elem. Math.* 28 (1973): 57–63.
- [46] Wills, J. M. "Nullstellenverteilung zweier konvexgeometrischer Polynome." *Beiträge Algebra Geom.* 29 (1989): 51–9.
- [47] Zvavitch, A. "Gaussian measure of sections of dilates and translations of convex bodies." *Adv. Appl. Math.* 41, no. 2 (2008): 247–54.