

Grassmann sheaves and the classification of vector sheaves

M. H. Papatriantafillou – E. Vassiliou*

Abstract

Given a sheaf of unital commutative and associative algebras \mathcal{A} , first we construct the k -th Grassmann sheaf $\mathcal{G}_{\mathcal{A}}(k, n)$ of \mathcal{A}^n , whose sections induce vector subsheaves of \mathcal{A}^n of rank k . Next we show that every vector sheaf over a paracompact space is a subsheaf of \mathcal{A}^∞ . Finally, applying the preceding results to the universal Grassmann sheaf $\mathcal{G}_{\mathcal{A}}(n)$, we prove that vector sheaves of rank n over a paracompact space are classified by the global sections of $\mathcal{G}_{\mathcal{A}}(n)$.

Introduction

Let \mathcal{A} be a sheaf of unital commutative and associative algebras over the ring \mathbb{R} or \mathbb{C} . A vector sheaf \mathcal{E} is a locally free \mathcal{A} -module. For instance, the sections of a vector bundle provide such a sheaf. However, a vector sheaf is not necessarily free, as is the case of the sections of a non trivial vector bundle.

Recently, vector sheaves gained a particular interest because they serve as the platform to abstract the classical geometry of vector bundles and their connections within a non smooth framework. This point of view has already been developed in [7] (see also [8] for applications to physics, and [10] for the reduction of the geometry of vector sheaves to the general setting of principal sheaves).

A fundamental result of the classical theory is the homotopy classification of vector bundles (of rank, say, n) over a fixed base. The construction of the classifying space, and the subsequent classification, are based on the

2000 *Mathematics Subject Classification*. Primary: 18F20. *keywords*. Vector sheaves, Grassmann sheaves.

*The authors were partially supported by University of Athens Research Grands 70/4/5639 and 70/4/3410, respectively.

Grassmann manifold (or variety) $G_k(\mathbb{R}^n)$ of k -dimensional subspaces of \mathbb{R}^n . In this respect we refer, e.g., to [5] and [6]. However, considering vector sheaves, we see that a homotopy classification is not possible, since the pull backs of a vector sheaf by homotopic maps need not be isomorphic, even in the trivial case of the free \mathcal{A} -module \mathcal{A} , as we prove in Section 1. Consequently, any attempt to classify vector sheaves (over a fixed space X) should not involve pull-backs and homotopy.

In this paper we develop a classification scheme based on a sort of universal Grassmann sheaf. More explicitly, for fixed $k \leq n \in \mathbb{N}$, in Section 2 we construct –in two equivalent ways– a sheaf $\mathcal{G}_{\mathcal{A}}(k, n)$, legitimately called the k -th Grassmann sheaf of \mathcal{A}^n , whose sections coincide (up to isomorphism) with vector subsheaves of \mathcal{A}^n of rank k (Proposition 2.3). Then, inducing in Section 3 the vector sheaf \mathcal{A}^∞ , we show that every vector sheaf over a paracompact space is a subsheaf of \mathcal{A}^∞ (Theorem 3.1). A direct application of the previous ideas leads us to the construction of the universal Grassmann sheaf $\mathcal{G}_{\mathcal{A}}(n)$ of rank n . The main result here (Theorem 3.5) asserts that arbitrary vector sheaves of rank n , over a paracompact base space, coincide –up to isomorphism– with the sections of $\mathcal{G}_{\mathcal{A}}(n)$.

1. Vector sheaves and homotopy

For the general theory of sheaves we refer to standard sources such as [1], [2], [4], and [9]. In what follows we recall a few definitions in order to fix the notations and terminology of the present paper.

Throughout the paper \mathcal{A} denotes a fixed sheaf of *unital commutative and associative* \mathbb{K} -algebras ($\mathbb{K} = \mathbb{R}, \mathbb{C}$) over a topological space X . An \mathcal{A} -module $\mathcal{E} \equiv (\mathcal{E}, \pi, X)$ is a sheaf whose stalks \mathcal{E}_x are \mathcal{A}_x -modules so that the respective operations of addition and scalar multiplication

$$\mathcal{E} \times_X \mathcal{E} \longrightarrow \mathcal{E} \quad \text{and} \quad \mathcal{A} \times_X \mathcal{E} \longrightarrow \mathcal{E}$$

are continuous. In particular, a *vector sheaf of rank n* is an \mathcal{A} -module \mathcal{E} , locally isomorphic to \mathcal{A}^n . This means there is an open covering $\mathcal{U} = \{U_\alpha\}$, $\alpha \in I$, of X and $\mathcal{A}|_{U_\alpha}$ -isomorphisms

$$\psi_\alpha: \mathcal{E}|_{U_\alpha} \longrightarrow \mathcal{A}^n|_{U_\alpha}, \quad \alpha \in I.$$

The category of vector sheaves of rank n over X is denoted by $\mathcal{V}^n(X)$. More details, examples and applications of vector sheaves can be found in [7], [8], and [10].

As already mentioned in the Introduction, we shall show, by a concrete counterexample, that homotopic maps do not yield isomorphic pull-backs, even in the simplest case of the free \mathcal{A} -module \mathcal{A} . In fact, we consider two *non-isomorphic* algebras A_0 and A_1 and a morphism of algebras $\rho: A_0 \rightarrow A_1$. Given now a topological space X and a fixed point $x_0 \in X$, for every open $U \subseteq X$ we set

$$A(U) := \begin{cases} A_0, & x_0 \in U, \\ A_1, & x_0 \notin U, \end{cases}$$

while, for every open $V \subseteq U$, $\rho_V^U: A(U) \rightarrow A(V)$ denotes the corresponding (restriction) map, defined by

$$\rho_V^U := \begin{cases} id: A_0 \rightarrow A_0, & \text{if } x_0 \in V, \\ \rho: A_0 \rightarrow A_1, & \text{if } U \ni x_0 \notin V, \\ id: A_1 \rightarrow A_1, & \text{if } x_0 \notin U. \end{cases}$$

It is not difficult to show that $(A(U), \rho_V^U)$ is a presheaf whose sheafification is a sheaf of algebras, denoted by $\mathcal{A} \equiv (\mathcal{A}, \pi, X)$. It is clear that

$$\mathcal{A}_{x_0} = \varinjlim_{U \in \mathcal{N}(x_0)} A(U) = A_0.$$

On the other hand, if there is an $x_1 \in X$ admitting a neighborhood $V \in \mathcal{N}(x_1)$ with $x_0 \notin V$ (a fact always ensured if X is a T_1 -space), then

$$\mathcal{A}_{x_1} = \varinjlim_{x_1 \in W \subseteq V} A(W) = A_1,$$

which is not isomorphic to \mathcal{A}_{x_0} . Hence, a vector sheaf, even a free one, need not have locally isomorphic fibres.

Let now $\alpha: [0, 1] \rightarrow X$ be a continuous path with $\alpha(0) = x_0$ and $\alpha(1) = x_1$. For any topological space Y , we define the map

$$f: [0, 1] \times Y \longrightarrow X \quad \text{with} \quad f(t, y) := \alpha(t).$$

Obviously, this is a homotopy between the constant maps $f_0 = x_0: Y \rightarrow X$ and $f_1 = x_1: Y \rightarrow X$. As a result, taking the pull-backs of \mathcal{A} by the latter, we see that, for every $y \in Y$,

$$\begin{aligned} f_0^*(\mathcal{A})_y &= \{(y, a) \mid a \in \mathcal{A}_{x_0}\} = \{y\} \times A_0, \\ f_1^*(\mathcal{A})_y &= \{(y, b) \mid b \in \mathcal{A}_{x_1}\} = \{y\} \times A_1; \end{aligned}$$

that is, we obtain two non-isomorphic stalks, thus proving the claim.

2. The Grassmann sheaf of rank k in \mathcal{A}^n

As in Section 1, \mathcal{A} is a sheaf of unital commutative and associative \mathbb{K} -algebras over a given topological space X . We denote by \mathfrak{T}_X the topology of X and fix $n \in \mathbb{N}$. For $k \in \mathbb{N}$ with $k \leq n$ and any $U \in \mathfrak{T}_X$, we define the set

$$G_{\mathcal{A}}(k, n)(U) := \{\mathcal{S} \text{ subsheaf of } \mathcal{A}^n|_U : \mathcal{S} \cong \mathcal{A}^k|_U\},$$

that is, $G_{\mathcal{A}}(k, n)(U)$ consists of the free submodules of $\mathcal{A}^n|_U$ of rank k . If, for every $U, V \in \mathfrak{T}_X$ with $V \subseteq U$,

$$\rho_V^U : G_{\mathcal{A}}(k, n)(U) \longrightarrow G_{\mathcal{A}}(k, n)(V) : \mathcal{S} \mapsto \mathcal{S}|_V$$

denotes the natural restriction, it is clear that the collection

$$(2.1) \quad G_{\mathcal{A}}(k, n) := (G_{\mathcal{A}}(k, n)(U), \rho_V^U)$$

determines a presheaf. Moreover, it is a *monopresheaf*. Indeed, if $U = \bigcup_{i \in I} U_i$ and $\mathcal{E}_1, \mathcal{E}_2 \in G_{\mathcal{A}}(k, n)(U)$ with $\mathcal{E}_1|_{U_i} = \mathcal{E}_2|_{U_i}$, for all $i \in I$, then $\mathcal{E}_1 = \mathcal{E}_2$. However, it is not complete: If $\mathcal{E}_i \in G_{\mathcal{A}}(k, n)(U_i)$, with $\mathcal{E}_i|_{U_i \cap U_j} = \mathcal{E}_j|_{U_i \cap U_j}$, then $\mathcal{E} := \bigcup_{i \in I} \mathcal{E}_i$ is a vector sheaf over U , but not necessarily a free $\mathcal{A}|_U$ -module.

Definition 2.1. The k -th Grassmann sheaf of \mathcal{A}^n , denoted by $\mathcal{G}_{\mathcal{A}}(k, n)$, is defined to be the sheaf generated by the presheaf $G_{\mathcal{A}}(k, n)$.

Since $G_{\mathcal{A}}(k, n)$ is not complete, it does not coincide with the complete presheaf

$$(2.2) \quad (\mathcal{G}_{\mathcal{A}}(k, n)(U), r_V^U),$$

of (continuous) sections of $\mathcal{G}_{\mathcal{A}}(k, n)$. We shall describe $\mathcal{G}_{\mathcal{A}}(k, n)$ via another complete presheaf. As a matter of fact, we consider the presheaf

$$(2.3) \quad V_{\mathcal{A}}(k, n) := (V_{\mathcal{A}}(k, n)(U), \lambda_V^U),$$

where now

$$V_{\mathcal{A}}(k, n)(U) := \{\mathcal{S} \text{ subsheaf of } \mathcal{A}^n|_U : \mathcal{S} \in \mathcal{V}^k(U)\}$$

and

$$\lambda_V^U : V_{\mathcal{A}}(k, n)(U) \longrightarrow V_{\mathcal{A}}(k, n)(V) : \mathcal{S} \mapsto \mathcal{S}|_V$$

are the natural restrictions. In contrast to $G_{\mathcal{A}}(k, n)$, $V_{\mathcal{A}}(k, n)$ is obviously a complete presheaf.

Lemma 2.2. *The sheaf generated by $V_{\mathcal{A}}(k, n)$ is isomorphic to $\mathcal{G}_{\mathcal{A}}(k, n)$.*

Proof. Clearly, for every $U \in \mathfrak{T}_X$,

$$G_{\mathcal{A}}(k, n)(U) \subseteq V_{\mathcal{A}}(k, n)(U),$$

that is, $G_{\mathcal{A}}(k, n)$ is a sub-presheaf of $V_{\mathcal{A}}(k, n)$. Besides, for every $\mathcal{E} \in V_{\mathcal{A}}(k, n)(U)$ and every $x \in U$, there is $V \in \mathfrak{T}_X$ with $x \in V \subseteq U$, so that $\mathcal{E}|_V$ is free of rank k , namely $\mathcal{E}|_V \in G_{\mathcal{A}}(k, n)(V)$. Thus, $G_{\mathcal{A}}(k, n)(U)$ and $V_{\mathcal{A}}(k, n)(U)$ define the same sheaf $\mathcal{G}_{\mathcal{A}}(k, n)$. \square

Since $V_{\mathcal{A}}(k, n)$ is complete, it is isomorphic with the sheaf of sections of $\mathcal{G}_{\mathcal{A}}(k, n)$, thus we have the following interpretation of the elements of $\mathcal{G}_{\mathcal{A}}(k, n)(X)$.

Proposition 2.3. *The global sections of the k -th Grassmann sheaf $\mathcal{G}_{\mathcal{A}}(k, n)$ coincide –up to isomorphism– with the vector subsheaves of \mathcal{A}^n of rank k .*

3. The universal Grassmann sheaf

The preliminary results of the preceding section hold for every base space X . Here we prove that if X is a paracompact space, then *any* vector sheaf can be interpreted as a section of an appropriate universal Grassmann sheaf.

First we prove a Whitney-type embedding theorem. To this end, for every sheaf of algebras \mathcal{A} , we consider the presheaf

$$U \longmapsto \prod_{i \in \mathbb{N}} \mathcal{A}_i(U), \quad U \in \mathfrak{T}_X,$$

where $\mathcal{A}_i = \mathcal{A}$, for every $i \in \mathbb{N}$, with the obvious restrictions. This presheaf generates the infinite *fibre* product

$$\mathcal{A}^\infty := \prod_{i \in \mathbb{N}} \mathcal{A}_i,$$

which is a free \mathcal{A} -module. Then, we obtain:

Theorem 3.1. *Let X be a paracompact space. Then every vector sheaf \mathcal{E} of finite rank over X is a subsheaf of \mathcal{A}^∞ .*

Proof. Let \mathcal{E} be a vector sheaf of rank, say, k . Since X is paracompact, a reasoning similar to that of [5, Proposition 5.4] proves that \mathcal{E} is free over a countable open covering $\{U_i\}_{i \in \mathbb{N}}$ of X . Let $\psi_i: \mathcal{E}|_{U_i} \rightarrow \mathcal{A}^k|_{U_i}$, $i \in \mathbb{N}$, be the

respective family of \mathcal{A} -module isomorphisms. The same open covering has a countable locally finite open refinement ([3, Ch. VIII, Theorem 1.4]), with a subordinate partition of unity $\{\alpha_i: X \rightarrow \mathbb{R}\}_{i \in \mathbb{N}}$ (ibid., Ch. VIII, Theorem 4.2). For every $i \in \mathbb{N}$, we define the map $\alpha_i \psi_i: \mathcal{E} \rightarrow \mathcal{A}^k$ by

$$\alpha_i \psi_i(u) := \begin{cases} \alpha_i(\pi(u)) \psi_i(u), & \pi(u) \in U_i, \\ 0, & \pi(u) \notin U_i. \end{cases}$$

Therefore, $\alpha_i \psi_i$ is an \mathcal{A} -module morphism, whose restriction to the interior of $\text{supp } \alpha_i \subseteq U_i$ is an isomorphism.

We consider the fibre product $\prod_{i \in \mathbb{N}} (\mathcal{A}^k)_i$, where $(\mathcal{A}^k)_i \equiv \mathcal{A}^k$, for every $i \in \mathbb{N}$, and we denote by

$$p_i: \prod_{i \in \mathbb{N}} (\mathcal{A}^k)_i \longrightarrow (\mathcal{A}^k)_i$$

the corresponding projections. The universal property of the product ensures the existence of a unique \mathcal{A} -morphism

$$\psi: \mathcal{E} \longrightarrow \prod_{i \in \mathbb{N}} (\mathcal{A}^k)_i,$$

such that

$$p_i \circ \psi = \alpha_i \psi_i.$$

Then ψ is a monomorphism. In fact, let $0 \neq u \in \mathcal{E}_x$ with $\psi_x(u) = 0$. There is $i \in \mathbb{N}$, with $\alpha_i(x) > 0$, thus $\alpha_i(x) \psi_{i,x}(u) \neq 0$, a contradiction. Hence, \mathcal{E} is identified with its image $\psi(\mathcal{E}) \leq \prod_{i \in \mathbb{N}} (\mathcal{A}^k)_i$. Since

$$\prod_{i \in \mathbb{N}} (\mathcal{A}^k)_i \equiv \prod_{i \in \mathbb{N}} \mathcal{A}_i,$$

where $\mathcal{A}_i = \mathcal{A}$, for every $i \in \mathbb{N}$, the assertion is proven. \square

We shall show that a further restriction on the topology of X leads to an embedding of \mathcal{E} into a smaller sheaf. To this end, assume that \mathcal{E} is a vector sheaf of rank k , which is free over a *finite* open covering $\{U_i\}_{1 \leq i \leq n}$ of X . Let $\psi_i: \mathcal{E}|_{U_i} \rightarrow \mathcal{A}^k|_{U_i}$ be the respective \mathcal{A} -module isomorphisms, and $\{\alpha_i: X \rightarrow \mathbb{R}\}_{1 \leq i \leq n}$ a subordinate partition of unity. Considering the maps $\alpha_i \psi_i: \mathcal{E} \rightarrow \mathcal{A}^k$, as before, we obtain the sheaf morphism

$$f: \mathcal{E} \longrightarrow \mathcal{A}^{kn}: u \mapsto (\alpha_1(\pi(u)) \cdot \psi_1(u), \dots, \alpha_n(\pi(u)) \cdot \psi_n(u))$$

which embeds \mathcal{E} into the free \mathcal{A} -module \mathcal{A}^{kn} . Therefore we have proved the following:

Proposition 3.2. *Over a compact space X , every vector sheaf of finite rank is a subsheaf of a free \mathcal{A} -module of finite rank.*

Clearly, every free \mathcal{A} -module \mathcal{A}^n is a submodule of the free \mathcal{A} -module $\bigoplus_{i \in \mathbb{N}} \mathcal{A}_i$, with $\mathcal{A}_i = \mathcal{A}$, for every $i \in \mathbb{N}$. Thus we obtain:

Corollary 3.3. *Over a compact space X , every vector sheaf of finite rank is a subsheaf of the free \mathcal{A} -module $\bigoplus_{i \in \mathbb{N}} \mathcal{A}_i$.*

We are now in a position to repeat the constructions of Section 2 in a more general way. For every $n \in \mathbb{N}$, we define the set

$$G_{\mathcal{A}}(n)(U) := \{ \mathcal{S} \text{ subsheaf of } \mathcal{A}^\infty|_U : \mathcal{S} \cong \mathcal{A}^n|_U \}.$$

Then the collection

$$G_{\mathcal{A}}(n) := (G_{\mathcal{A}}(n)(U), \rho_V^U), \quad U \in \mathfrak{T}_X,$$

where ρ_V^U denotes the obvious restriction, is a non-complete monopresheaf.

Definition 3.4. The sheaf $\mathcal{G}_{\mathcal{A}}(n)$, generated by the presheaf $G_{\mathcal{A}}(n)$, is called *the universal Grassmann sheaf of rank n* .

The respective complete presheaf of the sections of $\mathcal{G}_{\mathcal{A}}(n)$ is isomorphic to the presheaf

$$V_{\mathcal{A}}(n) := (V_{\mathcal{A}}(n)(U), \lambda_V^U), \quad U \in \mathfrak{T}_X,$$

where now

$$V_{\mathcal{A}}(n)(U) := \{ \mathcal{S} \text{ subsheaf of } \mathcal{A}^\infty|_U : \mathcal{S} \in \mathcal{V}^n(U) \}$$

and λ_V^U are the natural restrictions.

As a result, adapting the proof of Proposition 2.3 to the present situation, we obtain the main result of this work, namely the following classification of vector sheaves:

Theorem 3.5. *If X is a paracompact space, then the vector sheaves of rank n (over X) coincide –up to isomorphism– with the global sections of the universal Grassmann sheaf $\mathcal{G}_{\mathcal{A}}(n)$.*

References

- [1] G. L. Bredon: *Sheaf Theory*, 2nd edition, GTM **170**, Springer-Verlag, New York, 1997.
- [2] C. H. Dowker: *Lectures on Sheaf Theory*, Tata Inst. Fund. Research, Bombay, 1962.
- [3] J. Dugundji: *Topology*, Allyn and Bacon, Boston, 1966.
- [4] R. Godement: *Topologie Algébrique et Théorie des Faisceaux*, 3ème édition, Hermann, Paris, 1973.
- [5] D. Husemoller: *Fibre Bundles*, 3rd edition, Springer, New York, 1994.
- [6] D. W. Kahn: *Introduction to Global Analysis*, Academic Press, New York, 1980.
- [7] A. Mallios: *Geometry of Vector Sheaves. An axiomatic approach to differential geometry* (in 2 volumes), Kluwer, Dordrecht, 1998.
- [8] A. Mallios: *Modern Differential Geometry in Gauge Theories* (in 2 volumes), Birkhäuser, New York, 2006.
- [9] B. R. Tennison: *Sheaf Theory*, London Mathematical Society Lecture Note Series **20**, Cambridge University Press, Cambridge, 1975.
- [10] E. Vassiliou: *Geometry of Principal Sheaves*, Springer, New York, 2005.