

DICHOTOMIES OF THE SET OF TEST MEASURES OF A HAAR-NULL SET

PANDELIS DODOS

ABSTRACT. We prove that if X is a Polish space and F is a face of $P(X)$ with the Baire property, then F is either a meager or a co-meager subset of $P(X)$. As a consequence we show that for every abelian Polish group X and every analytic Haar-null set $A \subseteq X$, the set of test measures $T(A)$ of A is either meager or co-meager. We characterize the non-locally-compact groups as the ones for which there exists a closed Haar-null set $F \subseteq X$ with $T(F)$ is meager. Moreover, we answer negatively a question of Mycielski by showing that for every non-locally-compact abelian Polish group and every σ -compact subgroup G of X there exists a G -invariant F_G subset of X which is neither prevalent nor Haar-null.

1. INTRODUCTION AND AUXILIARY LEMMAS

A universally measurable subset A of an abelian Polish group X is called *Haar-null* if there exists a probability measure μ on X , called a *test measure* of A , such that $\mu(x + A) = 0$ for every $x \in X$. This definition is due to Christensen [C] and extends the usual notion of a Haar-measure zero set. The same class of sets has been also considered independently by Hunt, Sauer and Yorke [HSY]. They used the term *shy* instead of Haar-null. The complements of Haar-null sets are called *prevalent*. In Christensen's paper [C] a number of important properties of Haar-null sets were established. In particular, he showed that the class of Haar-null sets is a σ -ideal, which in the case of non-locally-compact groups contains all compact sets. On the other hand, he proved that if X is locally-compact, then Haar-null sets are precisely the Haar-measure zero sets and so his definition is indeed a generalization. There are, however, a number of properties of Haar-null sets which differentiate the locally from the non-locally compact case (see [BL] and the references therein). An important example is the countable chain condition, which is not satisfied if X is non-locally-compact—this is due to Dougherty [Dou] for a large class of abelian groups, and to Solecki [S1] in general.

Note that the test measure of a Haar-null set is not unique. Our motivation for this paper was to investigate the structure of the set of test measures $T(A)$ of a Haar-null set A . One natural question is about its size. Our first result states that there are only two extreme possibilities.

Research supported by a grant of EPEAEK program “Pythagoras”.

Theorem A. *Let X be an abelian Polish group. Then for every analytic Haar-null subset A of X the set*

$$T(A) = \{\mu \in P(X) : \mu(x + A) = 0 \text{ for every } x \in X\}$$

is either a meager or a co-meager subset of $P(X)$.

Actually, Theorem A is a consequence of a general property shared by the faces of $P(X)$. Specifically, we prove the following zero-one law.

Theorem B. *Let X be a Polish space and let F a face of $P(X)$ with the Baire property. Then F is either a meager or a co-meager subset of $P(X)$.*

The crucial point in the proof of Theorem B is the fact that convex averaging is open in $P(X)$.

Theorem A justifies the following definition. An analytic Haar-null set A is called *strongly Haar-null* if $T(A)$ is co-meager. Otherwise, it is called *weakly Haar-null*. In every abelian Polish group there exist strongly and weakly Haar-null sets. However the existence of a closed weakly Haar-null set characterizes the non-locally-compact groups.

Theorem C. *Let X be an abelian Polish group. Then X is non-locally-compact if and only if it contains a closed weakly Haar-null set.*

The proof of the above theorem is descriptive set-theoretic and uses the results of Solecki in [S2].

We also deal with a problem of Mycielski. He asked in [My] the following: if X is a non-locally-compact abelian Polish group, G is a countable dense subgroup of X and $A \subseteq X$ is a G -invariant universally measurable set, then is it true that A is either prevalent or Haar-null? The problem was answered negatively by Dougherty in $\mathbb{R}^{\mathbb{N}}$ (see [Dou]). Using a result of Matoušková and Zelený [MZ] we show that for every non-locally-compact abelian Polish group X and every σ -compact subgroup G of X , there exists a G -invariant F_σ subset of X which is neither prevalent nor Haar-null.

Finally, a measure-theoretic analogue of Theorem A is given on the last section. It is based on the fact that $T(A)$ is invariant under the group of homeomorphisms $\mu \rightarrow \mu * \delta_x$ where $x \in X$.

Notation. In what follows let X denote a Polish space (additional assumptions will be stated explicitly). By d we denote a compatible complete metric of X . If $x \in X$ and $r > 0$, then by $B(x, r)$ we denote the set $\{y \in X : d(x, y) < r\}$. By $P(X)$ we denote the space of all Borel probability measures on X , while by $M_+(X)$ the space of all positive finite Borel measures. Both $P(X)$ and $M_+(X)$ equipped with the weak topology are Polish spaces (see, e.g., [Pa]). If d is a compatible complete metric of X , then a compatible complete metric of $P(X)$ is the so-called

Lévy metric ϱ , defined by the rule

$$\rho(\mu, \nu) = \inf\{\delta \geq 0 : \mu(A) \leq \nu(A_\delta) + \delta \text{ and } \nu(A) \leq \mu(A_\delta) + \delta\}$$

where $A_\delta = \{x \in X : d(x, A) \leq \delta\}$. All balls in $P(X)$ are taken with respect to the Lévy metric ρ . That is, if $\mu \in P(X)$ and $r > 0$, then by $B(\mu, r)$ we denote the set $\{\nu \in P(X) : \rho(\mu, \nu) < r\}$ (from the context it will be clear whether we refer to a ball in $P(X)$ or in X). Finally, for every $\mu \in P(X)$ by $\text{supp}(\mu)$ we denote the support of the measure μ . All the other pieces of notation we use are standard (for more information we refer to [Ke]).

2. FACES OF $P(X)$

Through this section by X we denote Polish space. Recall that a subset F of $P(X)$ is called a *face* if F is an extreme convex subset of $P(X)$. For every $t \in (0, 1)$ consider the function $T_t : P(X) \times P(X) \rightarrow P(X)$ defined by

$$T_t(\mu, \nu) = t\mu + (1-t)\nu.$$

Clearly, every T_t is continuous. The following lemma provides an estimate for their range.

Lemma 1. *Let $r > 0$ and $m, \mu \in P(X)$ such that $\rho(m, \mu) < r$. Also let $t \in (0, 1)$. Then for every $\nu \in P(X)$ we have $\rho(T_t(\nu, m), \mu) \leq t + r$.*

Proof. Let $A \subseteq X$ be Borel. Then note that

$$\begin{aligned} t\nu(A) + (1-t)m(A) &\leq t + m(A) &\leq t + \mu(A_r) + r \\ &\leq \mu(A_{t+r}) + t + r. \end{aligned}$$

Similarly, we have

$$\begin{aligned} \mu(A) &\leq (1-t)\mu(A) + t\mu(A) \leq (1-t)m(A_r) + r + t \\ &\leq t\nu(A_{t+r}) + (1-t)m(A_{t+r}) + t + r. \end{aligned}$$

By the above inequalities, we obtain that $\rho(t\nu + (1-t)m, \mu) \leq t + r$. \square

The crucial property of T_t is that they are open. This result is due to Eifler [E1]. For the sake of completeness we include a proof.

Proposition 2. *For every $t \in (0, 1)$ the function T_t is open.*

Proof. Let us introduce some notation. Given $\mu, \nu \in M_+(X)$ we write $\nu \leq \mu$ if $\nu(A) \leq \mu(A)$ for every $A \subseteq X$ Borel (note that $\nu \leq \mu$ implies that $\nu \ll \mu$). Moreover, for every $\mu \in M_+(X)$ and every $f \in C_b(X)$ by $f\mu$ we denote the measure defined by $f\mu(A) = \int_A f(x) d\mu(x)$.

Claim 3. *Let $\mu, \nu \in M_+(X)$ with $\nu \leq \mu$ and let (μ_n) in $M_+(X)$ with $\mu_n \rightarrow \mu$. Then there exist a sequence (ν_k) in $M_+(X)$ and a subsequence (μ_{n_k}) of (μ_n) such that $\nu_k \rightarrow \nu$ and $\nu_k \leq \mu_{n_k}$ for every k .*

Proof of Claim 3. Let $g = d\nu/d\mu \in L^1(\mu)$. Since $\nu \leq \mu$, we have that $0 \leq g(x) \leq 1$ μ -a.e. Therefore, we may select a sequence (f_k) in $C_b(X)$ such that $f_k \rightarrow g$ in $L^1(\mu)$ and $0 \leq f_k(x) \leq 1$ for every $x \in X$ and every k . It follows that $f_k\mu \rightarrow g\mu = \nu$. Moreover, for any fixed k , we have $f_k\mu_n \rightarrow f_k\mu$ as $n \rightarrow \infty$. We select a subsequence (μ_{n_k}) of (μ_n) such that $f_k\mu_{n_k} \rightarrow \nu$ and we set $\nu_k = f_k\mu_{n_k}$. The proof of Claim 3 is completed. \square

Fix $0 < t < 1$ and set $w := t\mu + (1-t)\nu$ where $\mu, \nu \in P(X)$. It is enough to show that for every sequence (w_n) in $P(X)$ with $w_n \rightarrow w$ there exists a subsequence (w_{n_k}) of (w_n) and sequences $(\mu_k), (\nu_k)$ in $P(X)$ such that $\mu_k \rightarrow \mu$, $\nu_k \rightarrow \nu$ and $w_{n_k} = t\mu_k + (1-t)\nu_k$. So, let (w_n) be one. Since $t\mu \leq w$, by the above claim, there exist a sequence (m_k) in $M_+(X)$ and a subsequence (w_{n_k}) of (w_n) such that $m_k \rightarrow t\mu$ and $m_k \leq w_{n_k}$. Set $t_k := m_k(X)$. Then $t_k \rightarrow t$. By passing to further subsequences if necessary, we may assume that the sequence (t_k) is monotone. Without loss of generality suppose that $t_k \uparrow t$ (the case $t_k \downarrow t$ is similar). Set $m'_k := m_k/t_k$ and $\nu_k := (w_{n_k} - m_k)/(1-t_k)$. Then $m'_k, \nu_k \in P(X)$ and

$$\begin{aligned} w_{n_k} &= t_k m'_k + (1-t_k)\nu_k = (t-t-t_k)m'_k + (1-t+t-t_k)\nu_k \\ &= t\left(m'_k - \frac{t-t_k}{t}m'_k + \frac{t-t_k}{t}\nu_k\right) + (1-t)\nu_k. \end{aligned}$$

We set

$$\mu_k := m'_k - \frac{t-t_k}{t}m'_k + \frac{t-t_k}{t}\nu_k.$$

Then $\mu_k \in P(X)$. As $\mu_k \rightarrow \mu$ and $\nu_k \rightarrow \nu$ the proof is completed. \square

Remark 1. Note that the above proof is valid for all metrizable spaces, as long as we restrict ourselves to Radon probability measures. In this direction the more general result is contained in [E2].

Finally, we need the following elementary property of continuous, open functions.

Lemma 4. *Let Z, Y be Baire spaces and $f : Z \rightarrow Y$ a continuous, open function. Then $f^{-1}(G)$ is co-meager in Z for every $G \subseteq Y$ co-meager.*

We are ready to give our main result of this section.

Theorem 5. *Let X be a Polish space and let F be a face of $P(X)$ with the Baire property. Then F is either a meager or a co-meager subset of $P(X)$.*

Proof. Assume that F is not meager. We will show that in this case F is actually co-meager and this will finish the proof. Since F is not meager, there exist $\mu \in P(X)$ and $\lambda > 0$ such that F is co-meager in $B(\mu, \lambda)$. We select $a, r > 0$ small enough such that $a + r < \lambda/2$. Set $Z := P(X) \times B(\mu, r)$ and $Y := B(\mu, \lambda)$. Consider the restriction of T_a on Z . Denote the restriction by f . Clearly f remains open and continuous. By Lemma 1, we have $f(Z) \subseteq Y$. We set $W := F \cap Y$. By Lemma 4, we obtain that $f^{-1}(W)$ is co-meager in Z .

Set $G := F \cap B(\mu, r)$ (note that G is co-meager in $(B(\mu, r))$). Since F is an extreme convex subset of $P(X)$, for every $\nu \in P(X)$ and every $m \in B(\mu, r)$ we have $T_a(\nu, m) \in F$ if and only if $\nu, m \in F$. It follows that

$$f^{-1}(W) = F \times G,$$

whence F must be co-meager and the proof is completed. \square

3. THE TEST MEASURES OF AN ANALYTIC HAAR-NULL SET

Let X be an abelian Polish group (locally or non-locally compact). For every universally measurable set $A \subseteq X$ we set

$$T(A) := \{\mu \in P(X) : \mu(x + A) = 0 \text{ for every } x \in X\}.$$

Namely, $T(A)$ is the set of test measures of A . We have the following estimate for the complexity of $T(A)$ (see [D, Lemma 4]).

Lemma 6. *If $A \subseteq X$ is analytic, then $T(A)$ is a co-analytic subset of $P(X)$.*

Note that for every universally measurable set A , the set $T(A)$ is a face of $P(X)$. Therefore, by Theorem 5 and Lemma 6, we obtain the following topological dichotomy for the set of test measures of an analytic Haar-null set.

Corollary 7. *Let X be an abelian Polish group. Then for every analytic Haar-null set $A \subseteq X$ the set $T(A)$ is either a meager or a co-meager subset of $P(X)$.*

The following definition is motivated by Corollary 7.

Definition 8. *An analytic Haar-null set $A \subseteq X$ is called strongly Haar-null if $T(A)$ is co-meager. Otherwise, it is called weakly Haar-null.*

Under the terminology of the above definition, Corollary 7 states that every analytic weakly Haar-null set is necessarily tested by meager many measures. We should point out, however, that even if $T(A)$ may be small (in the topological sense), it is always dense in $P(X)$. Indeed, as has been indicated by Christensen (see [C]), for every $x \in X$ and $r > 0$ there exists $\mu \in T(A)$ such that $\rho(\mu, \delta_x) < r$. But the set of convex combinations of Dirac measures is dense in $P(X)$. So, using the properties of the Lévy metric, we can easily verify the density of $T(A)$.

As it has been shown in [D], every analytic and non-meager Haar-null set is necessarily weakly Haar-null (and so strongly Haar-null sets are necessarily meager). On the other hand, we have the following proposition (see [D, Proposition 5]).

Proposition 9. *Let X be an abelian Polish group and $A \subseteq X$ a σ -compact Haar-null set. Then A is strongly Haar-null.*

By the above proposition, it follows that in locally-compact groups closed Haar-null sets are strongly Haar-null.

The situation in non-locally-compact groups is quite different. In particular, we will show that in every non-locally-compact abelian Polish group X , there exists a closed weakly Haar-null set $F \subseteq X$. To this end we mention that the collection $F(X)$ of all closed subsets of X equipped with the Effros–Borel structure is a standard Borel space (see [Ke, page 75]). Denote by \mathcal{H} the collection of all closed Haar-null sets and by \mathcal{H}_s the collection of all closed strongly Haar-null sets. Clearly $\mathcal{H} \supseteq \mathcal{H}_s$. We claim that the inclusion is strict. Indeed, observe that, as noted by Solecki in [S2, page 211], the set

$$H = \{(F, \mu) \in F(X) \times P(X) : \mu(x + F) = 0 \ \forall x \in X\}$$

is Π_1^1 . It follows that the set

$$\mathcal{H}_s = \{F \in F(X) : \{\mu \in P(X) : (F, \mu) \in H\} \text{ is co-meager}\}$$

is Π_1^1 too (see [Ke, page 244]). But as it has been proved by Solecki, \mathcal{H} is Σ_1^1 -hard, hence not Π_1^1 (for the definition of Σ_1^1 -hard sets see [Ke] or [S2]). So the inclusion is strict. Summarizing, we obtain the following corollary which provides another characterization of non-locally-compact abelian Polish groups via properties of Haar-null sets.

Corollary 10. *Let X be an abelian Polish group. Then X is non-locally-compact if and only if there exists a closed weakly Haar-null set $F \subseteq X$.*

Remark 2. Clearly the crucial property of $T(A)$ we have used in order to prove Corollary 7, is that $T(A)$ has the Baire property. If $A \subseteq X$ is co-analytic, then we can easily verify that $T(A)$ is Π_2^1 . Under the standard axioms of set theory, Π_2^1 sets do not necessarily have the property of Baire. However, under any other additional axiom, capable of establishing that Π_2^1 have the Baire property (such as Σ_1^1 -determinacy or Martin’s Axiom and the negation of Continuum Hypothesis), Corollary 7 is valid for co-analytic sets. Moreover, it can be easily checked that under Projective Determinacy the dichotomy is valid for every projective set.

4. SETS INVARIANT UNDER K_σ SUBGROUPS

In [MZ], Matoušková and Zelený (using methods introduced by Solecki [S1]) proved the following theorem.

Theorem 11. *Let X be a non-locally-compact abelian Polish group. Then there exist closed non-Haar-null sets $A, B \subseteq X$ such that the set $(x + A) \cap B$ is compact for every $x \in X$.*

In what follows, by A and B we denote the sets obtained from Theorem 11. We will use the following notation. For every set $S \subseteq X$ we set

$$I(S) := \bigcup_{x \in S} ((x + A) \cap B).$$

We have the following lemma.

Lemma 12. *If $K \subseteq X$ is compact, then $I(K)$ is compact.*

Proof. We first observe that $I(K)$ is closed. Indeed, note that $I(K) = (K + A) \cap B$. Then, as K is compact and A is closed, we see that $K + A$ is closed; hence so is $I(K) = (K + A) \cap B$.

So it is enough to show that $I(K)$ is totally bounded. We will need certain facts from the construction of A and B made in [MZ]. By d we denote a compatible complete translation invariant metric of X .

Fix a dense countable subset $S = (s_n)$ of X and an increasing sequence (Q_k) of finite subsets of X such that $\bigcup_k Q_k$ is dense in X . Also fix sequences (ε_m) and (δ_m) or real numbers such that $\sum_{i>m} \varepsilon_i < \delta_m/2$ and which satisfy, for every m , the following fact proved in [MZ]: *for any pair F_1, F_2 of finite sets there exists $g \in B(0, \varepsilon)$ such that $\text{dist}(F_1, g + F_2) \geq \delta$* . Note that $\delta_m \rightarrow 0$.

Matoušková and Zelený, by induction, constructed sequences (g_k^m) and (\tilde{g}_k^m) which, among other things, satisfy the following (crucial for our considerations) property:

$$(P) \quad \forall n \text{ and } \forall i, j \text{ with } i, j \geq n \text{ we have } \text{dist}(s_n + g_i^m + Q_i, \tilde{g}_j^m + Q_j) \geq 3\delta_m.$$

Then they defined

$$A := \bigcap_{m \geq 1} \bigcup_{k \geq 1} \overline{B(g_k^m + Q_k, \delta_m)} \quad \text{and} \quad B := \bigcap_{m \geq 1} \bigcup_{k \geq 1} \overline{B(\tilde{g}_k^m + Q_k, \delta_m)}.$$

Let $r > 0$ be arbitrary. We select m such that $\delta_m < r$. From the compactness of K , we select l and $(s_{n_i})_{i=1}^l \subset S$ such that

$$\bigcup_{i=1}^l B(s_{n_i}, \delta_m/2) \supseteq K.$$

If $x \in X$ and $s_n \in S$ with $d(s_n, x) < \delta_m/2$, then, as it is shown in [MZ], from property (P) we have

$$(x + A) \cap B \subseteq \left(\bigcup_{k=1}^{n-1} \overline{B(s_n + g_k^m + Q_k, 3\delta_m/2)} \right) \cup \left(\bigcup_{k=1}^{n-1} \overline{B(\tilde{g}_k^m + Q_k, \delta_m)} \right).$$

It follows that

$$I(K) \subseteq \bigcup_{i=1}^l \left(\left(\bigcup_{k=1}^{n-1} \overline{B(s_{n_i} + g_k^m + Q_k, 3\delta_m/2)} \right) \cup \left(\bigcup_{k=1}^{n-1} \overline{B(\tilde{g}_k^m + Q_k, \delta_m)} \right) \right).$$

Therefore, the set $I(K)$ can be covered by finitely many balls of radii $2\delta_m < 2r$. As r was arbitrary, we see that $I(K)$ is totally bounded and the proof is completed. \square

Proposition 13. *Let X be non-locally-compact abelian Polish group and let G be a σ -compact subgroup of X . Then there exists a G -invariant F_σ subset F of X such that F is neither prevalent nor Haar-null. In particular, this holds if G is a countable dense subgroup of X .*

Proof. Write $G = \bigcup_n K_n$ where each K_n is compact. Let A and B be as in Theorem 11. Define

$$C_1 := \bigcup_{x \in G} (x + A) \quad \text{and} \quad C_2 := \bigcup_{x \in G} (x + B).$$

Note that for every n the sets $K_n + A$ and $K_n + B$ are closed. Also observe that $C_1 = \bigcup_n K_n + A$ and $C_2 = \bigcup_n K_n + B$. Therefore, both C_1 and C_2 are F_σ . In addition, from the fact that G is a subgroup of X , we see that both C_1 and C_2 are G -invariant.

As any possible translate of a non-Haar-null set is non-Haar-null, we have that C_1 and C_2 are non-Haar-null. We claim that at least one of them is not prevalent. Indeed, suppose that both C_1 and C_2 were prevalent. So $C_1 \cap C_2$ would be prevalent too. But observe that

$$\begin{aligned} C_1 \cap C_2 &= \left(\bigcup_{x \in G} (x + A) \right) \cap \left(\bigcup_{y \in G} (y + B) \right) \\ &= \bigcup_{n,m} \bigcup_{x \in K_n} \bigcup_{y \in K_m} (x + A) \cap (y + B) \\ &= \bigcup_{n,m} \bigcup_{x \in K_n} \bigcup_{y \in K_m} y + ((x - y) + A) \cap B \\ &\subseteq \bigcup_{n,m} K_m + I(K_n - K_m). \end{aligned}$$

Clearly, the set $K_n - K_m$ is compact. By Lemma 12, the set $I(K_n - K_m)$ is compact too, whence so is the set $K_m + I(K_n - K_m)$. But in non-locally-compact groups, compact sets are Haar-null. It follows that $C_1 \cap C_2$ is Haar-null and we derive a contradiction. Hence, at least one of them is not prevalent, as claimed. \square

Remark 3. Note that if G is countable, then the proof of Proposition 13 is considerably simpler (in particular, Lemma 12 is superfluous). However, Proposition 13 shows that invariance under bigger subgroups is not sufficient to establish the desired dichotomy. For instance, if X is a separable Banach space and (X_n) an increasing sequence of finite-dimensional subspaces with $\bigcup_n X_n$ dense in X , then we may let $G = \bigcup_n X_n$ and still provide a counterexample for the dichotomy.

5. ON THE MEASURE-THEORETIC STRUCTURE OF $T(A)$

By Lemma 6, for every $A \subseteq X$ analytic, the set $T(A)$ is a universally measurable subset of $P(X)$. Therefore, it is natural to wonder about the measure-theoretic structure of $T(A)$. If X is locally-compact, then it is easy to see that there exists $M \in P(P(X))$ (namely, M is a measure on measures) such that $M(T(A)) = 1$ for every $A \subseteq X$ universally measurable. Indeed, let h be the Haar measure and let H be the subset of $P(X)$ consisting of all probability measures μ absolutely continuous with respect to h . If $M \in P(P(X))$ is any measure on measures supported in H , then $M(T(A)) = 1$ for every universally measurable Haar-null set $A \subseteq X$.

In the non-locally-compact case the situation is different. As Stinchcombe observe in [St], for every $M \in P(P(X))$ there exists a σ -compact set $K \subseteq X$ such that $M(\{\mu : \mu(K) = 0\}) = 0$. Hence, for every $M \in P(P(X))$ there exists a Haar-null set $K \subseteq X$ for which $M(T(A)) = 0$. Nevertheless, we will show that in the non-locally-compact case there exists a measure-theoretic analogue of Corollary 7. So, in what follows we will assume that X is non-locally-compact. As before, by d we denote a translation invariant metric of X .

Let us introduce some notation. For every $x \in X$ and every $\mu \in P(X)$ define the probability measure μ_x by $\mu_x(A) = \mu(A - x)$ for every Borel set $A \subseteq X$. That is, $\mu_x = \mu * \delta_x$ (note that $\text{supp}(\mu_x) = x + \text{supp}(\mu)$). So every $x \in X$ gives rise to the function $g_x : P(X) \rightarrow P(X)$ defined by $g_x(\mu) = \mu_x$. Observe that the family $(g_x)_{x \in X}$ is an uncountable group of homeomorphisms of $P(X)$ (actually, G is a group of isometries of $(P(X), \rho)$). Also note that for every universally measurable $A \subseteq X$ the set $T(A)$ is G -invariant.

Lemma 14. *Let $\mu \in P(X)$ be such that $\text{supp}(\mu)$ is compact. Then the orbit $G\mu$ of μ is uncountable.*

Proof. Assume, on the contrary, that $G\mu$ is countable. Write $G\mu = (\nu_i)$ and for every i set

$$C_i := \{x \in X : \mu_x = \nu_i\}.$$

It is easy to verify that every C_i is closed. As $X = \bigcup_i C_i$, by the Baire category theorem, there exists k such that $\text{Int}(C_k) \neq \emptyset$. Therefore, there exist $z \in X$ and $r > 0$ such that $\mu_x = \mu_y = \nu_k$ for every $x, y \in B(z, r)$. But this implies that $x + \text{supp}(\mu) = y + \text{supp}(\mu)$ for every $x, y \in B(z, r)$, and so

$$B(0, r) \subseteq B(z, r) - B(z, r) \subseteq \text{supp}(\mu) - \text{supp}(\mu),$$

which is impossible as $\text{supp}(\mu)$ is compact and X is non-locally-compact. \square

If μ is any probability measure, then the orbit $G\mu$ of μ is Borel. To see this, observe that the group X acts continuously on $P(X)$ under the action $x \cdot \mu = \delta_x * \mu$ and that this action is precisely the action of G . It follows by a result of Miller (see [Ke, Mi]) that the orbit $G\mu$ of μ is Borel. Note, however, that if μ is a Dirac measure, then its orbit is the set of all Dirac measures. But, as it is well-known, this is a closed subset of $P(X)$. This fact is generalized in the following proposition.

Proposition 15. *Let $\mu \in P(X)$ be such that $\text{supp}(\mu)$ is compact. Then the orbit $G\mu$ of μ is closed.*

In order to prove Proposition 15 we will need a certain consequence of the classical Ramsey theorem [Ra] for doubletons of \mathbb{N} . As it is common in Ramsey theory, for every $I \subseteq \mathbb{N}$ by $[I]^2$ we denote the collection of all doubletons (i, j) such that $i, j \in I$ and $i < j$.

Lemma 16. *Let $r > 0$ and let (x_n) be a sequence such that $d(x_n, x_m) \geq r$ for every $n \neq m$. Then for every $K \subseteq X$ compact, there exists a subsequence (y_n) of (x_n) such that*

$$B(y_i - y_j, r/8) \cap K = \emptyset$$

for every i, j with $i < j$.

Proof. Let

$$A = \{(i, j) \in [\mathbb{N}]^2 : B(x_i - x_j, r/8) \cap K = \emptyset\}$$

and

$$B = \{(i, j) \in [\mathbb{N}]^2 : B(x_i - x_j, r/8) \cap K \neq \emptyset\}.$$

Then $[\mathbb{N}]^2 = A \cup B$ and $A \cap B = \emptyset$. By Ramsey's theorem, there exists an infinite subset M of \mathbb{N} such that either $[M]^2 \subseteq A$ or $[M]^2 \subseteq B$.

We claim that the $[M]^2 \subseteq A$. Indeed, assume on the contrary that $[M]^2 \subseteq B$. Set $m := \min\{i : i \in M\}$ and $M' := M \setminus \{m\}$. Then for every $n \in M'$ we have

$$B(x_m - x_n, r/8) \cap K \neq \emptyset.$$

So, for every $n \in M'$ there exists $w_n \in K$ such that $d(x_m - x_n, w_n) < r/8$. Since K is compact, we select a finite $r/8$ -net $(z_l)_{l=1}^k$ of K . By cardinality arguments, there exist an infinite set $I \subseteq M'$ and $l \in \{1, \dots, k\}$ such that $d(w_i, z_l) < r/8$ for every $i \in I$. Therefore, $d(w_i, w_j) < r/4$ for every $i, j \in I$. But if $i, j \in I$ with $i \neq j$, then we have

$$\begin{aligned} d(x_i, x_j) &= d(-x_i, -x_j) = d(x_m - x_i, x_m - x_j) \\ &\leq d(x_m - x_i, w_i) + d(w_i, w_j) + d(w_j, x_m - x_j) < r/2, \end{aligned}$$

which contradicts our assumption on (x_n) . Hence, $[M]^2 \subseteq A$.

Now let (m_n) be the increasing enumeration of M and set $y_n = x_{m_n}$ for every n . Clearly, (y_n) is the desired sequence. \square

We continue with the proof of Proposition 15.

Proof of Proposition 15. We first observe that we may assume that $0 \in \text{supp}(\mu)$. Indeed, if $y \in \text{supp}(\mu)$, then set $\nu := \mu_{-y}$ and notice that $0 \in \text{supp}(\nu)$ and $G\nu = G\mu$. So, in what follows we will assume that $0 \in \text{supp}(\mu)$.

Let (x_n) in X and $\nu \in P(X)$ such that $g_{x_n}(\mu) = \mu_{x_n} \rightarrow \nu$. Note that it suffices to prove that there exist $x \in X$ and a subsequence (x_{n_k}) of (x_n) such that $x_{n_k} \rightarrow x$. Indeed, in this case observe that $\delta_{x_{n_k}} \rightarrow \delta_x$. Hence, $\mu_{x_{n_k}} = \mu * \delta_{x_{n_k}} \rightarrow \mu * \delta_x = \mu_x$. On the other hand, since $(\mu_{x_{n_k}})$ is a subsequence of (μ_{x_n}) , we still have $\mu_{x_{n_k}} \rightarrow \nu$. This implies that $\nu = \mu_x$ as desired.

Assume, towards a contradiction, that the sequence (x_n) does not contain any convergent subsequence. So, there exist $r > 0$ and a subsequence (x_{n_k}) of (x_n) such that $d(x_{n_i}, x_{n_j}) \geq r$ for every $i \neq j$.

Applying Lemma 16 to the sequence (x_{n_k}) for $K = \text{supp}(\mu)$, we obtain a subsequence (y_n) of (x_{n_k}) —and so of (x_n) —such that for every i, j with $j > i$ we have

$$B(y_i - y_j, r/8) \cap \text{supp}(\mu) = \emptyset.$$

The subsequence (μ_{y_n}) of (μ_{x_n}) determined by (y_n) , still converges to ν . From the properties of (y_n) we have that if $n > i$, then

$$\mu_{y_n}(B(y_i, r/8)) = \mu(B(y_i, r/8) - y_n) = \mu(B(y_i - y_n, r/8)) = 0$$

as $B(y_i - y_n, r/8) \cap \text{supp}(\mu) = \emptyset$. Since for every open $U \subseteq X$ the function $\mu \mapsto \mu(U)$ is lower semicontinuous, we obtain that

$$\nu(B(y_i, r/8)) \leq \liminf \mu_{y_n}(B(y_i, r/8)) = 0$$

for every i . Therefore, the set $V := \bigcup_i B(y_i, r/8)$ is ν -null.

Next, set $F := \bigcup_i \overline{B(y_i, r/9)}$. Note that F is closed as $d(y_i, y_j) \geq r$ for $i \neq j$, and that $V \supseteq F$. Then observe that for every n we have

$$\mu_{y_n}(F) = \mu(F - y_n) = \mu\left(\bigcup_i \overline{B(y_i - y_n, r/9)}\right) \geq \mu(B(0, r/9)) > 0$$

where the last inequality holds from the fact that $0 \in \text{supp}(\mu)$. By the upper semicontinuity of the function $\mu \mapsto \mu(F)$, we obtain that

$$\nu(F) \geq \limsup \mu_{y_n}(F) \geq \mu(B(0, r/9)) > 0.$$

But this implies that

$$0 = \nu(V) \geq \nu(F) \geq \mu(B(0, r/9)) > 0$$

and we derive the contradiction. \square

Corollary 17. *Let X be a non-locally-compact abelian Polish group and let $A \subseteq X$ be an analytic Haar-null set. Then there exists a continuous Borel probability measure M on $P(X)$ with the following properties.*

- (i) *We have $M(T(A)) = 1$.*
- (ii) *If $B \subseteq X$ is any other analytic Haar-null set, then either $M(T(B)) = 1$ or $M(T(B)) = 0$.*

Proof. We select any compactly supported probability measure $\mu \in T(A)$. By Lemma 14 and Proposition 15, the orbit $G\mu$ of μ is an uncountable closed subset of $P(X)$. Therefore, if M is any continuous measure on measures supported in $G\mu$, then $M(T(A)) = 1$. Finally, observe that if $B \subseteq X$ is any other analytic Haar-null set, then as $T(B)$ is G -invariant either $G\mu \subseteq T(B)$ or $G\mu \cap T(B) = \emptyset$. This clearly establishes property (ii). \square

Acknowledgments. I am indebted to Professor S. Papadopoulou for providing me with references [E1] and [E2]. I would also like to thank Professor A. S. Kechris for his interest on the paper.

REFERENCES

- [BL] Y. Benyamini and J. Lindenstrauss, *Geometric Nonlinear Functional Analysis*, American Mathematical Society Colloquium Publications, 2000.
- [C] J. P. R. Christensen, *On sets of Haar measure zero in abelian Polish groups*, Israel J. Math. 13 (1972), 255–260.
- [D] P. Dodos, *On certain regularity properties of Haar-null sets*, Fund. Math. 181 (2004), 97–109.
- [Dou] R. Dougherty, *Examples of non shy sets*, Fund. Math. 144 (1994), 73–88.
- [E1] L. Q. Eifler, *Open mapping theorems for probability measures on metric spaces*, Pacific J. Math. 66 (1976), 89–97.
- [E2] L. Q. Eifler, *Convex averaging of Radon probability measures*, Glasnik Mat. Ser. III 12(32) (1977), 21–24.
- [HSY] B. Hunt, T. Sauer and J. Yorke, *Prevalence: a translation-invariant “almost every” on infinite-dimensional spaces*, Bull. Amer. Math. Soc. 27 (1992), 217–238.
- [Ke] A. S. Kechris, *Classical Descriptive Set Theory*, Grad. Texts in Math., Vol. 156, Springer-Verlag, 1995.
- [MZ] E. Matoušková and M. Zelený, *A note on intersection of non-Haar null sets*, Coll. Math. 96 (2003), 1–4.
- [Mi] B. D. Miller, *On the measurability of orbits in Borel actions*, Proc. Amer. Math. Soc. 63 (1977), 165–170.
- [My] J. Mycielski, *Some unsolved problems on the prevalence of ergodicity, instability and algebraic independence*, Ulam Quart. 1 (1992), 30–37.
- [Pa] K. R. Parthasarathy, *Probability Measures on Metric Spaces*, Academic Press, New York, 1967.
- [Ra] F. P. Ramsey, *On a problem of formal logic*, Proc. London Math. Soc. 30 (1930), 264–286.
- [S1] S. Solecki, *On Haar null sets*, Fund. Math. 149 (1996), 205–210.
- [S2] S. Solecki, *Haar null and non-dominating sets*, Fund. Math. 170 (2001), 197–217.
- [St] M. B. Stinchcombe, *The gap between probability and prevalence: loneliness in vector spaces*, Proc. Amer. Math. Soc. 129 (2001), 451–457.

NATIONAL TECHNICAL UNIVERSITY OF ATHENS, FACULTY OF APPLIED SCIENCES, DEPARTMENT OF MATHEMATICS, ZOGRAFOU CAMPUS, 157 80, ATHENS, GREECE.

E-mail address: `pdodos@math.ntua.gr`