

CODINGS OF SEPARABLE COMPACT SUBSETS OF THE FIRST BAIRE CLASS

PANDELIS DODOS

ABSTRACT. Let X be a Polish space and let \mathcal{K} be a separable compact subset of the first Baire class on X . For every sequence $\mathbf{f} = (f_n)$ dense in \mathcal{K} , the descriptive set-theoretic properties of the set

$$\mathcal{L}_{\mathbf{f}} := \{L \in [\mathbb{N}]^{\infty} : (f_n)_{n \in L} \text{ is pointwise convergent}\}$$

are analyzed. We show that if \mathcal{K} is not first countable, then $\mathcal{L}_{\mathbf{f}}$ is Π_1^1 -complete. This can also happen even if \mathcal{K} is a pre-metric compactum of degree at most two, in the sense of Todorćević. However, if \mathcal{K} is of degree exactly two, then $\mathcal{L}_{\mathbf{f}}$ is always Borel. A deep result of Debs implies that $\mathcal{L}_{\mathbf{f}}$ contains a Borel cofinal set, and this gives a tree-representation of \mathcal{K} . We show that classical ordinal assignments of Baire-1 functions are actually Π_1^1 -ranks on \mathcal{K} . We also provide an example of a Σ_1^1 Ramsey-null subset A of $[\mathbb{N}]^{\infty}$ for which there does not exist a Borel set $B \supseteq A$ such that the difference $B \setminus A$ is Ramsey-null.

1. INTRODUCTION

Let X be a Polish space. A Rosenthal compact on X is a subset of real-valued Baire-1 functions on X , compact in the pointwise topology. Standard examples of such compacta include the Helly space (the space of all non-decreasing functions from the unit interval into itself), the split interval (the lexicographical ordered product of the unit interval and the two-element ordering), and the ball of the double dual of a separable Banach space not containing ℓ_1 . The fact that the latter space is indeed a compact subset of the first Baire class follows from the famous Odell–Rosenthal theorem [OR] which states that the ball of the double dual of a separable Banach space with the weak* topology consists only of Baire-1 functions if and only if the space does not contain ℓ_1 . Actually this result motivated Rosenthal to initiate the study of compact subsets of the first Baire class in [Ro1]. He showed that all such compacta are sequentially compact. Bourgain, Fremlin and Talagrand proved that Rosenthal compacta are Fréchet spaces [BFT]. We refer to [AGR, P1, Ro2] for thorough introductions to the theory as well as its applications in analysis.

2000 *Mathematics Subject Classification*: 03E15, 26A21, 54H05, 05D10.

Research supported by a grant of EPEAEK program “Pythagoras”.

Separability is the crucial property that divides this class in two. As Todorćević has pointed out in [To] while non-separable Rosenthal compacta can be quite pathological, the separable ones are all “definable”. This is supported by the work of many researchers, including Godefroy [Go], Krawczyk [Kr], Marciszewski [Ma], Pol [P2] and is highlighted in the remarkable dichotomies and trichotomies of [To].

Our starting point of view is how we can code separable compact subsets of the first Baire class by members of a standard Borel space. Specifically, by a *code* of a separable Rosenthal compact \mathcal{K} on a Polish space X , we mean a standard Borel space C and a surjection $C \ni c \mapsto f_c \in \mathcal{K}$ such that for every $a \in \mathbb{R}$ the relation

$$(c, x) \in R_a \Leftrightarrow f_c(x) > a$$

is Borel in $C \times X$. In other words, inverse images of sub-basic open subsets of \mathcal{K} are Borel in C uniformly in X .

There is a natural object one associates to every separable Rosenthal compact \mathcal{K} and can serve as a coding of \mathcal{K} . More precisely, for every dense sequence $\mathbf{f} = (f_n)$ in \mathcal{K} we define

$$\mathcal{L}_{\mathbf{f}} := \{L \in [\mathbb{N}]^{\infty} : (f_n)_{n \in L} \text{ is pointwise convergent}\}.$$

The Bourgain–Fremlin–Talagrand theorem [BFT] implies that $\mathcal{L}_{\mathbf{f}}$ totally describes the members of \mathcal{K} , in the sense that for every accumulation point f of \mathcal{K} there exists $L \in \mathcal{L}_{\mathbf{f}}$ such that f is the pointwise limit of the sequence $(f_n)_{n \in L}$. Moreover, for every $f \in \mathcal{K}$ we define

$$\mathcal{L}_{\mathbf{f},f} := \{L \in [\mathbb{N}]^{\infty} : (f_n)_{n \in L} \text{ is pointwise convergent to } f\}.$$

Both $\mathcal{L}_{\mathbf{f}}$ and $\mathcal{L}_{\mathbf{f},f}$ have been studied in the literature. In [Kr], Krawczyk proved that $\mathcal{L}_{\mathbf{f},f}$ is Borel if and only if f is a G_{δ} point of \mathcal{K} . The set $\mathcal{L}_{\mathbf{f}}$ (more precisely, the set $\mathcal{L}_{\mathbf{f}} \setminus \mathcal{L}_{\mathbf{f},f}$) has been also considered by Todorćević [To] in his solution of characters of points in separable Rosenthal compacta.

There is an awkward fact concerning $\mathcal{L}_{\mathbf{f}}$, namely that $\mathcal{L}_{\mathbf{f}}$ can be non-Borel. However, a deep result of Debs [De] implies that $\mathcal{L}_{\mathbf{f}}$ always contains a Borel cofinal set and this subset of $\mathcal{L}_{\mathbf{f}}$ can serve as a coding. This leads to the following tree-representation of separable Rosenthal compacta.

Proposition A. *Let \mathcal{K} be a separable Rosenthal compact. Then there exist a countable tree T and a sequence $(g_t)_{t \in T}$ in \mathcal{K} such that the following hold.*

- (1) *For every $\sigma \in [T]$ the sequence $(g_{\sigma|n})$ is pointwise convergent.*
- (2) *For every $f \in \mathcal{K}$ there exists $\sigma \in [T]$ such that f is the pointwise limit of the sequence $(g_{\sigma|n})$.*

It is natural to ask when the set $\mathcal{L}_{\mathbf{f}}$ is Borel or, equivalently, when $\mathcal{L}_{\mathbf{f}}$ can serve itself as a coding (it is easy to see that $\mathcal{L}_{\mathbf{f}}$ and $\mathcal{L}_{\mathbf{f},f}$ are always Π_1^1). In this direction, we show the following theorem.

Theorem B. *Let \mathcal{K} be a separable Rosenthal compact.*

- (1) If \mathcal{K} is not first countable, then for every dense sequence $\mathbf{f} = (f_n)$ in \mathcal{K} the set $\mathcal{L}_{\mathbf{f}}$ is Π_1^1 -complete.
- (2) If \mathcal{K} is pre-metric of degree exactly two, then for every dense sequence $\mathbf{f} = (f_n)$ in \mathcal{K} the set $\mathcal{L}_{\mathbf{f}}$ is Borel.

Part (1) above is based on a result of Krawczyk. In part (2), \mathcal{K} is said to be a pre-metric compactum of degree exactly two if there exist a countable subset D of X and a countable subset \mathcal{D} of \mathcal{K} such that at most two functions in \mathcal{K} coincide on D and moreover for every $f \in \mathcal{K} \setminus \mathcal{D}$ there exists $g \in \mathcal{K}$ with $f \neq g$ and such that g coincides with f on D . This is a subclass of the class of pre-metric compacta of degree at most two, as it was defined by Todorćević in [To]. We notice that part (2) of Theorem B cannot be lifted to all pre-metric compacta of degree at most two, as there are examples of such compacta for which the set $\mathcal{L}_{\mathbf{f}}$ is Π_1^1 -complete.

We proceed now to discuss some applications of the above approach. It is well-known that to every real-valued Baire-1 function on a Polish space X one associates several (equivalent) ordinal rankings measuring the discontinuities of the function. An extensive study of them is done by Kechris and Louveau [KL]. An important example is the separation rank α . We have the following boundedness result concerning this index.

Theorem C. *Let X be a Polish space and let $\mathbf{f} = (f_n)$ be a sequence of Borel real-valued functions on X . Set*

$$\mathcal{L}_{\mathbf{f}}^1 := \{L \in [\mathbb{N}]^\infty : (f_n)_{n \in L} \text{ is pointwise convergent to a Baire-1 function}\}.$$

Then for every Borel set $C \subseteq \mathcal{L}_{\mathbf{f}}^1$ we have

$$\sup\{\alpha(f_L) : L \in C\} < \omega_1$$

where, for every $L \in C$, f_L denotes the pointwise limit of the sequence $(f_n)_{n \in L}$.

The proof of Theorem C is based on the fact that the separation rank is a parameterized Π_1^1 -rank. Theorem C, combined with the aforementioned result of Debs, gives a proof of the boundedness result of [ADK]. Historically the first result of this form is due to Bourgain [Bo]. We should point out that in order to give a descriptive set-theoretic proof of Bourgain's result one does not need to invoke Debs' theorem.

Theorem C can also be used to provide natural counterexamples to the following approximation question in Ramsey theory. Namely, given a Σ_1^1 subset A of $[\mathbb{N}]^\infty$ can we always find a Borel set $B \supseteq A$ such that the difference $B \setminus A$ is Ramsey-null? Miller had also asked whether there exists an analytic set which is not equal to Borel modulo Ramsey-null (see [Mi, Problem 1.6*]). We show the following proposition.

Proposition D. *There exists a Σ_1^1 Ramsey-null subset A of $[\mathbb{N}]^\infty$ for which there does not exist a Borel set $B \supseteq A$ such that the difference $B \setminus A$ is Ramsey-null.*

Acknowledgments. Part of this work was done when I visited the Department of Mathematics at Caltech. I would like to thank the department for the hospitality and the financial support. I would also like to thank the anonymous referee for his thorough report which improved, substantially, the presentation of the paper.

2. PRELIMINARIES

For any Polish space X by $K(X)$ we denote the hyperspace of all compact subsets of X equipped with the Vietoris topology. By $\mathcal{B}_1(X)$ (respectively, $\mathcal{B}(X)$) we denote the space of all real-valued Baire-1 (respectively, Borel) functions on X . By $\mathbb{N} = \{0, 1, 2, \dots\}$ we denote the natural numbers while by $[\mathbb{N}]^\infty$ the set of all infinite subsets of \mathbb{N} (which is clearly a Polish subspace of $2^\mathbb{N}$). For every $L \in [\mathbb{N}]^\infty$ by $[L]^\infty$ we denote the set of all infinite subsets of L . For every function $f: X \rightarrow \mathbb{R}$ and every $a \in \mathbb{R}$ by $[f > a]$ we denote the set $\{x : f(x) > a\}$. The set $[f < a]$ has the obvious meaning.

Our descriptive set-theoretic notation and terminology follows [Ke]. So Σ_1^1 stands for the analytic sets, while Π_1^1 for the co-analytic. A set is said to be Π_1^1 -true if it is co-analytic non-Borel. If X, Y are Polish spaces, $A \subseteq X$ and $B \subseteq Y$, we say that A is Wadge (respectively, Borel) reducible to B if there exists a continuous (respectively, Borel) map $f: X \rightarrow Y$ such that $f^{-1}(B) = A$. A set A is said to be Π_1^1 -complete if it is Π_1^1 and any other co-analytic set is Borel reducible to A . Clearly any Π_1^1 -complete set is Π_1^1 -true. The converse is also true under large cardinal hypotheses (see [MK] or [Mo]). If A is Π_1^1 , then a map $\phi: A \rightarrow \omega_1$ is said to be a Π_1^1 -rank on A if there are relations \leq_Σ , \leq_Π in Σ_1^1 and Π_1^1 respectively, such that for any $y \in A$

$$\phi(x) \leq \phi(y) \Leftrightarrow x \leq_\Sigma y \Leftrightarrow x \leq_\Pi y.$$

Notice that if A is Borel reducible to B via a Borel map f and ϕ is a Π_1^1 -rank on B , then the map $\psi: A \rightarrow \omega_1$ defined by $\psi(x) = \phi(f(x))$ is a Π_1^1 -rank on A .

Trees. If A is a nonempty set, then by $A^{<\mathbb{N}}$ we denote the set of all finite sequences of A . We view $A^{<\mathbb{N}}$ as a tree equipped with the (strict) partial order \sqsubset of extension. If $s \in A^{<\mathbb{N}}$, then the length $|s|$ of s is defined to be the cardinality of the set $\{t : t \sqsubset s\}$. If $s, t \in A^{<\mathbb{N}}$, then by $s \frown t$ we denote their concatenation. If $L \in [\mathbb{N}]^\infty$, then by $[L]^{<\mathbb{N}}$ we denote the increasing finite sequences in L . For every $x \in A^\mathbb{N}$ and every $n \geq 1$ we set $x|n = (x(0), \dots, x(n-1)) \in A^{<\mathbb{N}}$ while $x|0 = \emptyset$. A *tree* T on A is a downwards closed subset of $A^{<\mathbb{N}}$. The set of all trees on A is denoted by $\text{Tr}(A)$. Therefore,

$$T \in \text{Tr}(A) \Leftrightarrow \forall s, t \in A^{<\mathbb{N}} (t \sqsubset s \wedge s \in T \Rightarrow t \in T).$$

For a tree T on A the *body* $[T]$ of T is defined by $\{x \in A^\mathbb{N} : x|n \in T \text{ for every } n \in \mathbb{N}\}$. A tree T is called *pruned* if for every $s \in T$ there exists $t \in T$ with $s \sqsubset t$. It is called *well-founded* if for every $x \in A^\mathbb{N}$ there exists n such that $x|n \notin T$, equivalently if

$[T] = \emptyset$. The set of well-founded trees on A is denoted by $\text{WF}(A)$. If T is a well-founded tree, then we set $T' := \{t : \exists s \in T \text{ with } t \sqsubset s\}$. By transfinite recursion, one defines the iterated derivatives $T^{(\xi)}$ of T . The *order* $o(T)$ of T is defined to be the least ordinal ξ such that $T^{(\xi)} = \emptyset$. If S, T are well-founded trees, then a map $\phi: S \rightarrow T$ is called monotone if $s_1 \sqsubset s_2$ in S implies that $\phi(s_1) \sqsubset \phi(s_2)$ in T ; notice that in this case we have $o(S) \leq o(T)$. If A, B are sets, then we identify every tree T on $A \times B$ with the set of all pairs $(s, t) \in A^{<\mathbb{N}} \times B^{<\mathbb{N}}$ such that $|s| = |t| = k$ and $((s(0), t(0)), \dots, (s(k-1), t(k-1))) \in T$. If $A = \mathbb{N}$, then we shall simply denote by Tr and WF the sets of all trees and well-founded trees on \mathbb{N} respectively. The set WF is Π_1^1 -complete and the map $T \mapsto o(T)$ is a Π_1^1 -rank on WF . The same holds for $\text{WF}(A)$ for every countable set A .

The separation rank. Let X be a Polish space. Given $A, B \subseteq X$ one associates with them a derivative on closed sets, by $F'_{A,B} = \overline{F \cap A} \cap \overline{F \cap B}$. By transfinite recursion, we define the iterated derivatives $F_{A,B}^{(\xi)}$ of F and we set $\alpha(F, A, B)$ to be the least ordinal ξ with $F_{A,B}^{(\xi)} = \emptyset$ if such an ordinal exists; otherwise we set $\alpha(F, A, B) = \omega_1$. Now let $f: X \rightarrow \mathbb{R}$ be a function. For each pair $a, b \in \mathbb{R}$ with $a < b$ let $A = [f < a]$ and $B = [f > b]$. For every closed set $F \subseteq X$ let $F_{f,a,b}^{(\xi)} = F_{A,B}^{(\xi)}$ and $\alpha(f, F, a, b) = \alpha(F, A, B)$. Also let $\alpha(f, a, b) = \alpha(f, X, a, b)$. The separation rank of f is defined by

$$\alpha(f) = \sup\{\alpha(f, a, b) : a, b \in \mathbb{Q}, a < b\}.$$

The basic fact is the following (see [KL]).

Proposition 1. *A function f is Baire-1 if and only if $\alpha(f) < \omega_1$.*

3. CODINGS OF SEPARABLE ROSENTHAL COMPACTA

Let X be a Polish space and let $\mathbf{f} = (f_n)$ be a sequence of Borel real-valued functions on X . Assume that the closure \mathcal{K} of $\{f_n\}$ in \mathbb{R}^X is a compact subset of $\mathcal{B}(X)$. Consider the set

$$\mathcal{L}_{\mathbf{f}} = \{L \in [\mathbb{N}]^\infty : (f_n)_{n \in L} \text{ is pointwise convergent}\}.$$

For every $L \in \mathcal{L}_{\mathbf{f}}$ by f_L we shall denote the pointwise limit of the sequence $(f_n)_{n \in L}$. Notice that $\mathcal{L}_{\mathbf{f}}$ is Π_1^1 . As the pointwise topology is not effected by the topology on X , we may (and we will) assume that each f_n is continuous (and so, \mathcal{K} is a separable Rosenthal compact). By a result of Rosenthal [Ro1], we see that $\mathcal{L}_{\mathbf{f}}$ is cofinal; that is, for every $M \in [\mathbb{N}]^\infty$ there exists $L \in [M]^\infty$ such that $L \in \mathcal{L}_{\mathbf{f}}$. Also the celebrated Bourgain–Fremlin–Talagrand theorem [BFT] implies that $\mathcal{L}_{\mathbf{f}}$ totally describes \mathcal{K} . However, most important for our purposes is the fact that $\mathcal{L}_{\mathbf{f}}$ contains a Borel cofinal set. This is a consequence of the following theorem of Debs [De] (which itself is the classical interpretation of the effective version of the Bourgain–Fremlin–Talagrand theorem proved by Debs in [De]).

Theorem 2. *Let Y, X be Polish spaces and let (g_n) be a sequence of Borel functions on $Y \times X$ such that for every $y \in Y$ the sequence $(g_n(y, \cdot))$ is a sequence of continuous functions relatively compact in $\mathcal{B}(X)$. Then there exists a Borel map $\sigma: Y \rightarrow [\mathbb{N}]^\infty$ such that for every $y \in Y$ the sequence $(g_n(y, \cdot))_{n \in \sigma(y)}$ is pointwise convergent.*

Let us show how Theorem 2 implies the existence of a Borel cofinal subset of \mathcal{L}_f . Given $L, M \in [\mathbb{N}]^\infty$ with $L = \{l_0 < l_1 < \dots\}$ and $M = \{m_0 < m_1 < \dots\}$ their increasing enumerations, we set $L * M = \{l_{m_0} < l_{m_1} < \dots\}$. Clearly $L * M \in [L]^\infty$ and, moreover, the function $(L, M) \mapsto L * M$ is continuous. Let (f_n) be as in the beginning of the section and set $Y = [\mathbb{N}]^\infty$. For every $n \in \mathbb{N}$ we define the function $g_n: [\mathbb{N}]^\infty \times X \rightarrow \mathbb{R}$ by setting

$$g_n(L, x) = f_{l_n}(x)$$

where l_n is the n -th element of the increasing enumeration of L . The sequence (g_n) satisfies the hypotheses of Theorem 2. Let $\sigma: [\mathbb{N}]^\infty \rightarrow [\mathbb{N}]^\infty$ be the Borel function such that for every $L \in [\mathbb{N}]^\infty$ the sequence

$$(g_n(L, \cdot))_{n \in \sigma(L)} = (f_n)_{n \in L * \sigma(L)}$$

is pointwise convergent. The function $L \mapsto L * \sigma(L)$ is Borel and so the set

$$A = \{L * \sigma(L) : L \in [\mathbb{N}]^\infty\}$$

is an analytic cofinal subset of \mathcal{L}_f . By separation, we conclude that there exists a Borel cofinal subset of \mathcal{L}_f . The cofinality of this set in conjunction with the Bourgain–Fremlin–Talagrand theorem yield the following corollary.

Corollary 3. *Let X be a Polish space and let (f_n) be a sequence of Borel functions on X which is relatively compact in $\mathcal{B}(X)$. Then there exists a Borel set $C \subseteq [\mathbb{N}]^\infty$ such that for every $c \in C$ the sequence $(f_n)_{n \in c}$ is pointwise convergent, and for every accumulation point f of (f_n) there exists $c \in C$ with $f = \lim_{n \in c} f_n$.*

In the sequel we will say that the set C obtained by Corollary 3 is a *code* of (f_n) . If \mathcal{K} is a separable Rosenthal compact and (f_n) is a dense sequence in \mathcal{K} , then we will say that C is the code of \mathcal{K} . Notice that the codes depend on the dense sequence. If $c \in C$, then by f_c we shall denote the function coded by c . That is, f_c is the pointwise limit of the sequence $(f_n)_{n \in c}$.

In the following lemma we present some basic definability properties of the set of codes. Its easy proof is left to the reader.

Lemma 4. *Let X and (f_n) be as in Corollary 3, and let C be a code of (f_n) . Then for every $a \in \mathbb{R}$ the following relations*

- (i) $(c, x) \in R_a \Leftrightarrow f_c(x) > a$,
- (ii) $(c, x) \in R'_a \Leftrightarrow f_c(x) \geq a$,
- (iii) $(c_1, c_2, x) \in D_a \Leftrightarrow |f_{c_1}(x) - f_{c_2}(x)| > a$

are Borel.

The existence of codings of separable Rosenthal compacta gives us the following tree-representation of them.

Proposition 5. *Let \mathcal{K} be a separable Rosenthal compact. Then there exist a countable tree T and a sequence $(g_t)_{t \in T}$ in \mathcal{K} such that the following hold.*

- (1) *For every $\sigma \in [T]$ the sequence $(g_{\sigma|n})$ is pointwise convergent.*
- (2) *For every $f \in \mathcal{K}$ there exists $\sigma \in [T]$ such that f is the pointwise limit of the sequence $(g_{\sigma|n})$.*

Proof. Let (f_n) be a dense sequence in \mathcal{K} . We may assume that for every $n \in \mathbb{N}$ the set $\{m : f_m = f_n\}$ is infinite. This extra condition guarantees that for every $f \in \mathcal{K}$ there exists $L \in \mathcal{L}_f$ such that $f = f_L$. Let C be the codes of (f_n) . Now we shall use a common unfolding trick. Since C is Borel in $2^{\mathbb{N}}$, there exists $F \subseteq 2^{\mathbb{N}} \times \mathbb{N}^{\mathbb{N}}$ closed such that $C = \text{proj}_{2^{\mathbb{N}}} F$. Let T be the unique (downwards closed) pruned tree on $2 \times \mathbb{N}$ such that $F = [T]$. This will be the desired tree. It remains to define the sequence $(g_t)_{t \in T}$. Set $g_{(\emptyset, \emptyset)} = f_0$. Let $t = (s, w) \in T$ and $k \geq 1$ with $s \in 2^{<\mathbb{N}}$, $w \in \mathbb{N}^{<\mathbb{N}}$ and $|s| = |w| = k$. Define $n_t \in \mathbb{N}$ by setting $n_t := \max\{n < k : s(n) = 1\}$, if the set $\{n < k : s(n) = 1\}$ is nonempty; otherwise, set $n_t := 0$. Finally, set $g_t := f_{n_t}$. It is easy to check that for every $\sigma \in [T]$ the sequence $(g_{\sigma|n})$ is pointwise convergent, and so (1) is satisfied. The fact that (2) is also satisfied follows from the fact that for every $f \in \mathcal{K}$ there exists $L \in \mathcal{L}_f$ with $f = f_L$ and the fact that C is cofinal. \square

Remark 1. (1) We should point out that Corollary 3, combined with Silver's theorem (see [MK] or [S2]) on the number of equivalence classes of co-analytic equivalence relations, gives an answer to the cardinality problem of separable Rosenthal compacta, a well-known fact that can also be derived by the results of [To] (see also [ADK, Remark 3]). Indeed, let \mathcal{K} be a separable Rosenthal compact and let C be the set of codes of \mathcal{K} . Define the following equivalence relation on C by

$$c_1 \sim c_2 \Leftrightarrow f_{c_1} = f_{c_2} \Leftrightarrow \forall x \ f_{c_1}(x) = f_{c_2}(x).$$

Then \sim is a Π_1^1 equivalence relation. Hence, by Silver's dichotomy, either the equivalence classes are countable or perfectly many. The first case implies that $|\mathcal{K}| = \aleph_0$ while the second one yields that $|\mathcal{K}| = 2^{\aleph_0}$.

(2) Although the set C of codes of a separable Rosenthal compact \mathcal{K} is considered to be a Borel set which describes \mathcal{K} efficiently, when it is considered as a subset of $[\mathbb{N}]^{\infty}$ it can be chosen to have rich structural properties. In particular, it can be chosen to be hereditary (that is, if $c \in C$ and $c' \in [c]$, then $c' \in C$) and invariant under finite changes. To see this, start with a code C_1 of \mathcal{K} , that is, a Borel cofinal

subset of \mathcal{L}_f . Let

$$\begin{aligned} \Phi = \{ (F, G) : (F \subseteq \mathcal{L}_f) \wedge (G \cap C_1 = \emptyset) \wedge \\ [\forall L, M (L \in F \wedge M \subseteq L \Rightarrow M \notin G)] \wedge \\ [\forall L, M, s (L \in F \wedge (L \triangle M = s) \Rightarrow M \notin G)] \}. \end{aligned}$$

Also let $A = \{N : \exists L \in C_1 \exists s \in [\mathbb{N}]^{<\mathbb{N}} \exists M \in [L]^\infty \text{ with } N \triangle M = s\}$. Then A is Σ_1^1 and clearly $\Phi(A, \sim A)$. Since Φ is Π_1^1 on Σ_1^1 , hereditary and continuous upward in the second variable, by the dual form of the second reflection theorem (see [Ke, Theorem 35.16]), there exists $C \supseteq A$ Borel with $\Phi(C, \sim C)$. Then C is as desired.

(3) We notice that the idea of coding subsets of function spaces by converging sequences appears also in [Be], where a representation result of Σ_2^1 subsets of $C([0, 1])$ is proved.

4. A BOUNDEDNESS RESULT

4.1. Determining $\alpha(f)$ by compact sets. Let X be a Polish space and let $f: X \rightarrow \mathbb{R}$ be a Baire-1 function. The aim of this subsection is to show that the value $\alpha(f)$ is completely determined by the derivatives taken over compact subsets of X (notice that this is trivial if X is compact metrizable). Specifically, we have the following proposition.

Proposition 6. *Let X be a Polish space, $f: X \rightarrow \mathbb{R}$ Baire-1 and $a < b$ reals. Then $\alpha(f, a, b) = \sup\{\alpha(f, K, a, b) : K \subseteq X \text{ compact}\}$.*

The proof of Proposition 6 is an immediate consequence of the following lemmas. In what follows, all balls in X are taken with respect to some compatible complete metric ρ of X .

Lemma 7. *Let X , f and $a < b$ be as in Proposition 6. Also let $F \subseteq X$ be closed, let $x \in X$ and let $\xi < \omega_1$ such that $x \in F_{f,a,b}^{(\xi)}$. Then for every $\varepsilon > 0$, setting $C = F \cap \overline{B(x, \varepsilon)}$, we have $x \in C_{f,a,b}^{(\xi)}$.*

Proof. Fix F and ε as above. For notational simplicity we set $U := B(x, \varepsilon)$ and $C := F \cap \overline{B(x, \varepsilon)}$. By induction, we shall show that

$$F^{(\xi)} \cap U \subseteq C^{(\xi)}$$

where $F^{(\xi)} = F_{f,a,b}^{(\xi)}$ and similarly for C . This clearly implies the lemma. For $\xi = 0$ is straightforward. Suppose that the lemma is true for every $\xi < \zeta$. Assume that $\zeta = \xi + 1$ is a successor ordinal. Let $y \in F^{(\xi+1)} \cap U$. Since U is open, we have

$$y \in \overline{F^{(\xi)} \cap U \cap [f < a]} \cap \overline{F^{(\xi)} \cap U \cap [f > b]}.$$

By the inductive assumption, we obtain that

$$y \in \overline{C^{(\xi)} \cap [f < a]} \cap \overline{C^{(\xi)} \cap [f > b]} = C^{(\xi+1)}$$

which proves the case of successor ordinals. If ζ is limit, then

$$F^{(\zeta)} \cap U = \bigcap_{\xi < \zeta} F^{(\xi)} \cap U \subseteq \bigcap_{\xi < \zeta} C^{(\xi)} = C^{(\zeta)}$$

and the lemma is proved. \square

Lemma 8. *Let X , f and $a < b$ be as in Proposition 6. Also let $F \subseteq X$ be closed, let $x \in X$ and let $\xi < \omega_1$ such that $x \in F_{f,a,b}^{(\xi)}$. Then there exists countable compact $K \subseteq F$ such that $x \in K_{f,a,b}^{(\xi)}$.*

Proof. Again, for notational simplicity, for every closed set $C \subseteq X$ and every $\xi < \omega_1$ we set $C^{(\xi)} = C_{f,a,b}^{(\xi)}$. The proof is by induction on countable ordinals, as before. For $\xi = 0$ the lemma is obviously true. Suppose that the lemma has been proved for every $\xi < \zeta$. Let $F \subseteq X$ be closed and let $x \in F^{(\zeta)}$. Notice that one of the following alternatives must occur.

- (A1) $f(x) < a$ and there exists a sequence (y_n) such that $y_n \neq y_m$ if $n \neq m$, $f(y_n) > b$, $y_n \in F^{(\xi_n)}$ and $y_n \rightarrow x$;
- (A2) $f(x) > b$ and there exists a sequence (z_n) such that $z_n \neq z_m$ if $n \neq m$, $f(z_n) < a$, $z_n \in F^{(\xi_n)}$ and $z_n \rightarrow x$;
- (A3) there exist two distinct sequences (y_n) and (z_n) such that $y_n \neq y_m$ and $z_n \neq z_m$ if $n \neq m$, $f(y_n) < a$, $f(z_n) > b$, $y_n, z_n \in F^{(\xi_n)}$ and $y_n \rightarrow x$, $z_n \rightarrow x$,

where above the sequence (ξ_n) of countable ordinals is as follows.

- (C1) If $\zeta = \xi + 1$, then $\xi_n = \xi$ for every n .
- (C2) If ζ is limit, then (ξ_n) is an increasing sequence of successor ordinals with $\xi_n \nearrow \zeta$.

We shall treat alternative (A1) (the other ones are similar). Let (r_n) be a sequence of positive reals such that $\overline{B(y_n, r_n)} \cap \overline{B(y_m, r_m)} = \emptyset$ if $n \neq m$ and $x \notin \overline{B(y_n, r_n)}$ for every n . Set $C_n := F \cap \overline{B(y_n, r_n)}$. By Lemma 7, we obtain that $y_n \in C_n^{(\xi_n)}$. By the inductive assumption, there exists $K_n \subseteq C_n \subseteq F_n$ countable compact such that $y_n \in K_n^{(\xi_n)}$. Finally, set $K := \{x\} \cup (\bigcup_n K_n)$. Then K is countable compact and it is easy to see that $x \in K^{(\zeta)}$. \square

Remark 2. Notice that the proof of Lemma 8 actually shows that

$$\alpha(f, a, b) = \sup\{\alpha(f, K, a, b) : K \subseteq X \text{ countable compact}\}.$$

Moreover observe that if $\alpha(f, a, b)$ is a successor ordinal, then the above supremum is attained.

4.2. The main result. This subsection is devoted to the proof of the following theorem.

Theorem 9. *Let X be a Polish space and let $\mathbf{f} = (f_n)$ be a sequence of Borel real-valued functions on X . Set*

$$\mathcal{L}_{\mathbf{f}}^1 := \{L \in [\mathbb{N}]^\infty : (f_n)_{n \in L} \text{ is pointwise convergent to a Baire-1 function}\}.$$

Then for every Borel set $C \subseteq \mathcal{L}_{\mathbf{f}}^1$ we have

$$\sup\{\alpha(f_L) : L \in C\} < \omega_1$$

where, for every $L \in C$, f_L denotes the pointwise limit of the sequence $(f_n)_{n \in L}$.

For the proof of Theorem 9 we will need the following theorem which gives us a way of defining parameterized Π_1^1 -ranks (see [Ke, page 275]).

Theorem 10. *Let Y be a standard Borel space, let X be a Polish space and let $\mathbb{D} : Y \times K(X) \rightarrow K(X)$ be a Borel map such that for every $y \in Y$ the map \mathbb{D}_y is a derivative on $K(X)$. Then the set*

$$\Omega_{\mathbb{D}} := \{(y, K) : \mathbb{D}_y^{(\infty)}(K) = \emptyset\}$$

is Π_1^1 and the map $(y, K) \rightarrow |K|_{\mathbb{D}_y}$ is a Π_1^1 -rank on $\Omega_{\mathbb{D}}$.

We proceed to the proof of Theorem 9.

Proof of Theorem 9. Let $C \subseteq \mathcal{L}_{\mathbf{f}}^1$ be Borel. Fix $a, b \in \mathbb{R}$ with $a < b$. Define $\mathbb{D} : C \times K(X) \rightarrow K(X)$ by

$$\mathbb{D}(L, K) = \overline{K \cap [f_L < a]} \cap \overline{K \cap [f_L > b]}$$

where f_L is the pointwise limit of the sequence $(f_n)_{n \in L}$. It is clear that for every $L \in C$ the map $K \mapsto \mathbb{D}(L, K)$ is a derivative on $K(X)$ and $\alpha(f_L, K, a, b) = |K|_{\mathbb{D}_L}$. We will show that \mathbb{D} is Borel. Define $A, B \in C \times K(X) \times X$ by

$$(L, K, x) \in A \Leftrightarrow (x \in K) \wedge (f_L(x) < a)$$

and

$$(L, K, x) \in B \Leftrightarrow (x \in K) \wedge (f_L(x) > b).$$

It is easy to check that both A and B are Borel. Also let $\tilde{A}, \tilde{B} \subseteq C \times K(X) \times X$ be defined by

$$(L, K, x) \in \tilde{A} \Leftrightarrow x \in \overline{A_{(L, K)}}$$

and

$$(L, K, x) \in \tilde{B} \Leftrightarrow x \in \overline{B_{(L, K)}},$$

where $A_{(L, K)} = \{x : (L, K, x) \in A\}$ is the section of A (and, similarly, for B). Notice that for every $(L, K) \in C \times K(X)$ we have $\mathbb{D}(L, K) = \tilde{A}_{(L, K)} \cap \tilde{B}_{(L, K)}$. Since $\tilde{A}_{(L, K)}$ and $\tilde{B}_{(L, K)}$ are compact (they are subsets of K), by Theorem 28.8 in [Ke], it is enough to show that the sets \tilde{A} and \tilde{B} are Borel. We will need the following easy consequence of the Arsenin–Kunugui theorem (the proof is left to the reader).

Lemma 11. *Let Z be a standard Borel space, let X be a Polish space and let $F \subseteq Z \times X$ be Borel with K_σ sections. Then the set \tilde{F} defined by*

$$(z, x) \in \tilde{F} \Leftrightarrow x \in \overline{F_z}$$

is a Borel subset of $Z \times X$.

By our assumptions, for every $L \in C$ the function f_L is Baire-1 and so for every $(L, K) \in C \times K(X)$ the sections $A_{(L,K)}$ and $B_{(L,K)}$ of A and B respectively, are K_σ . Hence, by Lemma 11, we see that \tilde{A} and \tilde{B} are Borel.

By the above discussion, we conclude that \mathbb{D} is a Borel map. By Theorem 10, the map $(L, K) \rightarrow |K|_{\mathbb{D}_L}$ is a Π_1^1 -rank on $\Omega_{\mathbb{D}}$. By Proposition 1 and the fact that $C \subseteq \mathcal{L}_{\mathbf{f}}^1$, we obtain that for every $(L, K) \in C \times K(X)$ the transfinite sequence $(\mathbb{D}_L^{(\xi)}(K))_{\xi < \omega_1}$ must be stabilized at \emptyset and so $\Omega_{\mathbb{D}} = C \times K(X)$. Since $\Omega_{\mathbb{D}}$ is Borel, by boundedness, we have

$$\sup\{|K|_{\mathbb{D}_L} : (L, K) \in C \times K(X)\} < \omega_1.$$

It follows that

$$\sup\{\alpha(f_L, K, a, b) : (L, K) \in C \times K(X)\} < \omega_1.$$

By Proposition 6, we obtain that

$$\sup\{\alpha(f_L, a, b) : L \in C\} < \omega_1.$$

This completes the proof of the theorem. \square

4.3. Consequences. We recall some definitions from [ADK]. Let X be a Polish space, let (f_n) a sequence of real-valued functions on X and let \mathcal{K} be the closure of $\{f_n\}$ in \mathbb{R}^X . We say that \mathcal{K} is a (separable) *quasi-Rosenthal* if every accumulation point of \mathcal{K} is a Baire-1 function and, moreover, we say that \mathcal{K} is *Borel separable* if the sequence (f_n) consists of Borel functions. Combining Theorem 9 with Corollary 3, we obtain the following result from [ADK].

Theorem 12. *Let X be a Polish space and let \mathcal{K} be a Borel separable quasi-Rosenthal compact on X . Then*

$$\sup\{\alpha(f) : f \in \text{Acc}(\mathcal{K})\} < \omega_1$$

where $\text{Acc}(\mathcal{K})$ denotes the accumulation points of \mathcal{K} . In particular, if \mathcal{K} is a separable Rosenthal compact on X , then

$$\sup\{\alpha(f) : f \in \mathcal{K}\} < \omega_1.$$

Besides boundedness, the implications of Theorem 9 and the relation between the separation rank and the Borelness of $\mathcal{L}_{\mathbf{f}}$ are more transparently seen when X is a compact metrizable space. In particular, we have the following proposition.

Proposition 13. *Let X be a compact metrizable space and let \mathcal{K} be a separable Rosenthal compact on X . Let $\mathbf{f} = (f_n)$ be a dense sequence in \mathcal{K} and $a < b$ reals. Then the map $L \mapsto \alpha(f_L, a, b)$ is a Π_1^1 -rank on $\mathcal{L}_{\mathbf{f}}$ if and only if the set $\mathcal{L}_{\mathbf{f}}$ is Borel.*

Proof. First assume that $\mathcal{L}_{\mathbf{f}}$ is not Borel. By Theorem 12, we have

$$\sup\{\alpha(f_L, a, b) : L \in \mathcal{L}_{\mathbf{f}}\} < \omega_1$$

and so the map $L \mapsto \alpha(f_L, a, b)$ cannot be a Π_1^1 -rank on $\mathcal{L}_{\mathbf{f}}$, since $\mathcal{L}_{\mathbf{f}}$ is Π_1^1 -true. Conversely, assume that $\mathcal{L}_{\mathbf{f}}$ is Borel. By the proof of Theorem 9, we have that the map $(L, K) \mapsto |K|_{\mathbb{D}_L}$ is a Π_1^1 -rank on $\mathcal{L}_{\mathbf{f}} \times K(X)$. It follows that the relation

$$L_1 \preceq L_2 \Leftrightarrow \alpha(f_{L_1}, a, b) \leq \alpha(f_{L_2}, a, b) \Leftrightarrow |X|_{\mathbb{D}_{L_1}} \leq |X|_{\mathbb{D}_{L_2}}$$

is Borel in $\mathcal{L}_{\mathbf{f}} \times \mathcal{L}_{\mathbf{f}}$. This implies that the map $L \mapsto \alpha(f_L, a, b)$ is a Π_1^1 -rank on $\mathcal{L}_{\mathbf{f}}$ as desired. \square

Remark 3. Although the map $L \mapsto \alpha(f_L, a, b)$ is not always a Π_1^1 -rank on $\mathcal{L}_{\mathbf{f}}$, it is easy to see that it is a Π_1^1 -rank on the codes C of \mathcal{K} , since the relation

$$c_1 \preceq c_2 \Leftrightarrow \alpha(f_{c_1}, a, b) \leq \alpha(f_{c_2}, a, b) \Leftrightarrow |X|_{\mathbb{D}_{c_1}} \leq |X|_{\mathbb{D}_{c_2}}$$

is Borel in $C \times C$ for every pair $a < b$ of reals. Hence, when X is compact metrizable space, we could say that the separation rank is a Π_1^1 -rank “in the codes”.

We proceed to discuss another application of Theorem 9 which deals with the following approximation question in Ramsey theory. Recall that a set $N \subseteq [\mathbb{N}]^\infty$ is called *Ramsey-null* if for every $s \in [\mathbb{N}]^{<\mathbb{N}}$ and every $L \in [\mathbb{N}]^\infty$ with $s < L$, there exists $L' \in [L]^\infty$ such that $[s, L'] \cap N = \emptyset$. As every analytic set is Ramsey [S1], it is natural to ask the following. Is it true that for every analytic set $A \subseteq [\mathbb{N}]^\infty$ there exists $B \supseteq A$ Borel such that $B \setminus A$ is Ramsey-null? As we will show the answer is no and a counterexample can be found which is in addition Ramsey-null.

To this end we will need some pieces of notation from [AGR]. Let X be a separable Banach space. By $X_{\mathcal{B}_1}^{**}$ we denote the set of all Baire-1 elements of the ball of the second dual X^{**} of X . We say that X is α -universal if

$$\sup\{\alpha(x^{**}) : x^{**} \in X_{\mathcal{B}_1}^{**}\} = \omega_1.$$

We should point out that there exist non-universal (in the classical sense) separable Banach spaces which are α -universal (see [AD]). We have the following proposition.

Proposition 14. *There exists a Σ_1^1 Ramsey-null subset A of $[\mathbb{N}]^\infty$ for which there does not exist a Borel set $B \supseteq A$ such that the difference $B \setminus A$ is Ramsey-null.*

Proof. Let X be a separable α -universal Banach space and fix a norm dense sequence $\mathbf{f} = (x_n)$ in the ball of X (it will be convenient to assume that $x_n \neq x_m$ if $n \neq m$). Let

$$\mathcal{L}_{\mathbf{f}} := \{L \in [\mathbb{N}]^\infty : (x_n)_{n \in L} \text{ is weak}^* \text{ convergent}\}.$$

Clearly \mathcal{L}_f is Π_1^1 . Moreover, notice that $\mathcal{L}_f = \mathcal{L}_f^1$ according to the notation of Theorem 9.

Let $x^{**} \in X_{\mathcal{B}_1}^{**}$ be arbitrary. By the Odell–Rosenthal theorem (see [AGR], or [OR]), there exists $L \in \mathcal{L}_f$ such that $x^{**} = w^* - \lim_{n \in L} x_n$. It follows that

$$\sup\{\alpha(x^{**}) : x^{**} \in X_{\mathcal{B}_1}^{**}\} = \sup\{\alpha(x_L) : L \in \mathcal{L}_f\}$$

where x_L denotes the weak* limit of the sequence $(x_n)_{n \in L}$. Denote by (e_n) the standard basis of ℓ_1 and set

$$\Lambda := \{L \in [\mathbb{N}]^\infty : \exists k \text{ such that } (x_n)_{n \in L} \text{ is } (k+1)\text{-equivalent to } (e_n)\}$$

where if $L \in [\mathbb{N}]^\infty$ with $L = \{l_0 < l_1 < \dots\}$ its increasing enumeration, then $(x_n)_{n \in L}$ is $(k+1)$ -equivalent to (e_n) if for every $m \in \mathbb{N}$ and every $a_0, \dots, a_m \in \mathbb{R}$ we have

$$\frac{1}{k+1} \sum_{n=0}^m |a_n| \leq \left\| \sum_{n=0}^m a_n x_{l_n} \right\| \leq (k+1) \sum_{n=0}^m |a_n|.$$

Notice that Λ is Σ_2^0 . We also remark that, by Bourgain's result [Bo] and our assumptions on the space X , the set Λ is nonempty. Also let

$$\Lambda_1 := \{N \in [\mathbb{N}]^\infty : \exists L \in \Lambda \exists s \in [\mathbb{N}]^{<\mathbb{N}} \text{ such that } N \triangle L = s\}.$$

Clearly Λ_1 is Σ_2^0 too. Observe that both \mathcal{L}_f and Λ_1 are hereditary and invariant under finite changes. Moreover the set $\mathcal{L}_f \cup \Lambda_1$ is cofinal. This is essentially a consequence of Rosenthal's dichotomy (see, for instance, [LT]). It follows that the set $A = [\mathbb{N}]^\infty \setminus (\mathcal{L}_f \cup \Lambda_1)$ is Σ_1^1 and Ramsey-null.

We claim that A is the desired set. Assume not, that is, there exists a Borel set $B \supseteq A$ such that the difference $B \setminus A$ is Ramsey-null. We set $C = [\mathbb{N}]^\infty \setminus (B \cup \Lambda_1)$. Then $C \subseteq \mathcal{L}_f$ is Borel and, moreover, $\mathcal{L}_f \setminus C$ is Ramsey-null. It follows that for every $x^{**} \in X_{\mathcal{B}_1}^{**}$ there exists $L \in C$ such that $x^{**} = x_L$. Since C is Borel, by Theorem 9, we have

$$\sup\{\alpha(x^{**}) : x^{**} \in X_{\mathcal{B}_1}^{**}\} = \sup\{\alpha(x_L) : L \in C\} < \omega_1$$

which contradicts the fact that X is α -universal. The proof is completed. \square

Remark 4. (1) An example as in Proposition 14 can also be given using the convergence rank γ studied by Kechris and Louveau [KL]. As the reasoning is the same, we shall briefly describe the argument. Let (f_n) be a sequence of continuous function on $2^\mathbb{N}$ with $\|f_n\|_\infty \leq 1$ for every $n \in \mathbb{N}$ and such that the set $\{f_n : n \in \mathbb{N}\}$ is norm dense in the ball of $C(2^\mathbb{N})$. As in Proposition 14, consider the sets \mathcal{L}_f , Λ_1 and $A = [\mathbb{N}]^\infty \setminus (\mathcal{L}_f \cup \Lambda_1)$. Then the set A is Σ_1^1 and Ramsey-null. The fact that the set A cannot be covered by a Borel set B such that the difference $B \setminus A$ is Ramsey-null follows essentially from the following properties.

- (P1) The map $(g_n) \mapsto \gamma((g_n))$ is a Π_1^1 -rank on the set $CN = \{(g_n) \in C(2^{\mathbb{N}})^{\mathbb{N}} : (g_n) \text{ is pointwise convergent}\}$; see [Ke, page 279]. Hence, the map

$$\mathcal{L}_{\mathbf{f}} \ni L = \{l_0 < l_1 < \dots\} \mapsto \gamma((f_{l_n}))$$

is a Π_1^1 -rank on $\mathcal{L}_{\mathbf{f}}$.

- (P2) For every $\Delta \in \Delta_2^0$, there exists $L \in \mathcal{L}_{\mathbf{f}}$ such that the sequence $(f_n)_{n \in L}$ is pointwise convergent to $\mathbf{1}_{\Delta}$. By Proposition 1 in [KL], we obtain that $\alpha(\mathbf{1}_{\Delta}) \leq \gamma((f_n)_{n \in L})$. It follows that

$$\sup\{\gamma((f_n)_{n \in L}) : L \in \mathcal{L}_{\mathbf{f}}\} \geq \sup\{\alpha(\mathbf{1}_{\Delta}) : \Delta \in \Delta_2^0\} = \omega_1.$$

(2) For the important special case of a separable Rosenthal compact \mathcal{K} defined on a compact metrizable space X and having a dense set of continuous functions, Theorem 12 has originally been proved by Bourgain [Bo]. We should point out that in this case one does not need Corollary 3 in order to carry out the proof. Let us briefly explain how this can be done. So assume that X is compact metrizable and $\mathbf{f} = (f_n)$ is a sequence of continuous functions dense in \mathcal{K} . Fix $a, b \in \mathbb{Q}$ with $a < b$ and for every $n \in \mathbb{N}$ let $A_n = [f_n \leq a]$ and $B_n = [f_n \geq b]$. For every $M \in [\mathbb{N}]^{\infty}$ set

$$\liminf_{n \in M} A_n = \bigcup_{n \in M} \bigcap_{k \geq n, k \in M} A_k$$

and similarly for $\liminf_{n \in M} B_n$. Observe the following.

- (O1) For every $M \in [\mathbb{N}]^{\infty}$ the sets $\liminf_{n \in M} A_n$ and $\liminf_{n \in M} B_n$ are Σ_2^0 .
- (O2) If $L, M \in [\mathbb{N}]^{\infty}$ are such that $L \subseteq M$, then $\liminf_{n \in M} A_n \subseteq \liminf_{n \in L} A_n$, and similarly for B_n .
- (O3) If $L \in \mathcal{L}_{\mathbf{f}}$, then $[f_L < a] \subseteq \liminf_{n \in L} A_n \subseteq [f_L \leq a]$ and, respectively, $[f_L > b] \subseteq \liminf_{n \in L} B_n \subseteq [f_L \geq b]$.

Define $\mathbb{D}: [\mathbb{N}]^{\infty} \times K(X) \rightarrow K(X)$ by

$$\mathbb{D}(M, K) = \overline{K \cap \liminf_{n \in M} A_n} \cap \overline{K \cap \liminf_{n \in M} B_n}.$$

By (O1) and using the same arguments as in the proof of Theorem 9, we can easily verify that \mathbb{D} is Borel. Since $\mathcal{L}_{\mathbf{f}}$ is cofinal, by (O2) and (O3), we can also easily verify that $\Omega_{\mathbb{D}} = [\mathbb{N}]^{\infty} \times K(X)$. Therefore, by boundedness, we obtain that $\sup\{|K|_{\mathbb{D}_M} : (M, K) \in [\mathbb{N}]^{\infty} \times K(X)\} < \omega_1$. Using (O3) again, we finally obtain that $\sup\{\alpha(f, a, b) : f \in \mathcal{K}\} < \omega_1$, as desired.

5. ON THE DESCRIPTIVE SET-THEORETIC PROPERTIES OF $\mathcal{L}_{\mathbf{f}}$

In this section we will show that certain topological properties of a separable Rosenthal compact \mathcal{K} imply the Borelness of the set $\mathcal{L}_{\mathbf{f}}$. To this end, we recall that \mathcal{K} is said to be a *pre-metric compactum of degree at most two* if there exists a countable subset D of X such that at most two functions in \mathcal{K} agree on D (see [To]). Let us consider the following subclass.

Defintion 15. We say that \mathcal{K} is a pre-metric compactum of degree exactly two if there exist a countable subset D of X and a countable subset \mathcal{D} of \mathcal{K} such that at most two functions in \mathcal{K} coincide on D and, moreover, for every $f \in \mathcal{K} \setminus \mathcal{D}$ there exists $g \in \mathcal{K}$ with $g \neq f$ and such that g coincides with f on D .

An important example of such a compact is the split interval (but it is not the only important one—see Remark 5 below). Under the above terminology we have the following theorem.

Theorem 16. Let X be a Polish space and let \mathcal{K} a separable Rosenthal compact on X . If \mathcal{K} is pre-metric of degree exactly two, then for every dense sequence $\mathbf{f} = (f_n)$ in \mathcal{K} the set $\mathcal{L}_{\mathbf{f}}$ is Borel.

Proof. Let $\mathbf{f} = (f_n)$ be a dense sequence in \mathcal{K} and let C be the set of codes of (f_n) . Also let $D \subseteq X$ be countable and let $\mathcal{D} \subseteq \mathcal{K}$ be the countable set verifying that \mathcal{K} is pre-metric of degree exactly two.

Claim. There exists $\mathcal{D}' \subseteq \mathcal{K}$ countable with $\mathcal{D} \subseteq \mathcal{D}'$ such that for every $c \in C$ with $f_c \in \mathcal{K} \setminus \mathcal{D}'$ there exists $c' \in C$ such that $f_{c'} \neq f_c$ and $f_{c'}$ coincides with f_c on D .

Proof of the claim. Let $c \in C$ be such that $f_c \in \mathcal{K} \setminus \mathcal{D}$. Let g be the (unique) function in \mathcal{K} with $g \neq f_c$ and such that g coincides with f_c on D . If there does not exist $c' \in C$ with $g = f_{c'}$, then g is an isolated point of \mathcal{K} . We set

$$\mathcal{D}' := \mathcal{D} \cup \{f \in \mathcal{K} : \exists g \in \mathcal{K} \text{ isolated such that } f(x) = g(x) \forall x \in D\}.$$

Since the isolated points of \mathcal{K} are countable and \mathcal{K} is pre-metric of degree at most two, we obtain that \mathcal{D}' is countable. Clearly \mathcal{D}' is as desired. \square

Let \mathcal{D}' be the set obtained above and set

$$\mathcal{L}_{\mathcal{D}'} := \bigcup_{f \in \mathcal{D}'} \mathcal{L}_{\mathbf{f}, f} = \bigcup_{f \in \mathcal{D}'} \{L \in [\mathbb{N}]^\infty : (f_n)_{n \in L} \text{ is pointwise convergent to } f\}.$$

Since every point in \mathcal{K} is G_δ , we see that $\mathcal{L}_{\mathcal{D}'}$ is Borel (actually, it is Σ_4^0). Consider the following equivalence relation \sim on C defined by the rule

$$c_1 \sim c_2 \Leftrightarrow \forall x \in D \ f_{c_1}(x) = f_{c_2}(x).$$

By Lemma 4, the equivalence relation \sim is Borel. Next, consider the relation P on $C \times C \times K(X) \times X$ defined by

$$(c_1, c_2, K, x) \in P \Leftrightarrow (c_1 \sim c_2) \wedge (x \in K) \wedge (|f_{c_1}(x) - f_{c_2}(x)| > 0).$$

Again we see that P is Borel. Moreover, notice that for every $c_1, c_2 \in C$ the function $x \mapsto |f_{c_1}(x) - f_{c_2}(x)|$ is Baire-1, and so, for every $(c_1, c_2, K) \in C \times C \times K(X)$ the section $P_{(c_1, c_2, K)} = \{x \in X : (c_1, c_2, K, x) \in P\}$ of P is K_σ . By Theorem 35.46 in [Ke], the set $S \subseteq C \times C \times K(X)$ defined by

$$(c_1, c_2, K) \in S \Leftrightarrow \exists x \ (c_1, c_2, K, x) \in P$$

is Borel and there exists a Borel map $\phi: S \rightarrow X$ such that for every $(c_1, c_2, K) \in S$ we have $(c_1, c_2, K, \phi(c_1, c_2, K)) \in P$. By the above claim, we have that for every $c \in C \setminus \mathcal{L}_{\mathcal{D}'}$ there exist $c' \in C$ and $K \in K(X)$ such that $(c, c', K) \in S$. Moreover, observe that the set $D \cup \{\phi(c, c', K)\}$ determines the neighborhood basis of f_c . The crucial fact is that this can be done in a Borel way.

Now we claim that

$$\begin{aligned} L \in \mathcal{L}_{\mathbf{f}} \Leftrightarrow & (L \in \mathcal{L}_{\mathcal{D}'}) \vee \left[(\forall x \in D (f_n(x))_{n \in L} \text{ converges}) \wedge \right. \\ & \left. \{ \exists s \in S \text{ with } s = (c_1, c_2, K) \text{ such that} \right. \\ & \left. [\forall x \in D f_{c_1}(x) = \lim_{n \in L} f_n(x)] \wedge \right. \\ & \left. [(f_n(\phi(s)))_{n \in L} \text{ converges}] \wedge \right. \\ & \left. [f_{c_1}(\phi(s)) = \lim_{n \in L} f_n(\phi(s))] \} \right]. \end{aligned}$$

Granting this, the proof is completed since the above expression gives a Σ_1^1 definition of $\mathcal{L}_{\mathbf{f}}$. As $\mathcal{L}_{\mathbf{f}}$ is also Π_1^1 , this implies that $\mathcal{L}_{\mathbf{f}}$ is Borel, as desired.

It remains to prove the above equivalence. First assume that $L \in \mathcal{L}_{\mathbf{f}}$. We need to show that L satisfies the expression on the right. If $L \in \mathcal{L}_{\mathcal{D}'}$ this is clearly true. If $L \notin \mathcal{L}_{\mathcal{D}'}$, then pick a code $c \in C \setminus \mathcal{L}_{\mathcal{D}'}$ such that $f_c = f_L$. By the above claim and the remarks of the previous paragraph, we can easily verify that L satisfies the expression on the right. Conversely, let L fulfill the right side of the equivalence. If $L \in \mathcal{L}_{\mathcal{D}'}$ we are done. If not, then by the Bourgain–Fremlin–Talagrand theorem, it suffices to show that all convergent subsequences of $(f_n)_{n \in L}$ have the same limit. The first two conjuncts enclosed in the square brackets on the right side of the equivalence guarantee that each such convergent subsequence of $(f_n)_{n \in L}$ converges either to f_{c_1} or to f_{c_2} . The last two conjuncts guarantee that it is not f_{c_2} , so it is always f_{c_1} . Thus, $L \in \mathcal{L}_{\mathbf{f}}$ and the proof is completed. \square

Remark 5. (1) Let \mathcal{K} be a pre-metric compactum of degree at most two and let $D \subseteq X$ countable such that at most two functions in \mathcal{K} coincide on D . Notice that the set C of codes of \mathcal{K} is naturally divided into two parts, namely

$$C_2 := \{c \in C : \exists c' \in C \text{ with } f_c \neq f_{c'} \text{ and } f_c(x) = f_{c'}(x) \forall x \in D\}$$

and its complement $C_1 = C \setminus C_2$. The assumption that \mathcal{K} is pre-metric of degree exactly two, simply reduces to the assumption that the functions coded by C_1 are at most countable. We could say that C_1 is the set of metrizable codes, since it is immediate that the set $\{f_c : c \in C_1\}$ is a metrizable subspace of \mathcal{K} . It is easy to check, using the set S defined in the proof of Theorem 16, that C_2 is always Σ_1^1 . As we shall see, it might happen that C_1 is Π_1^1 -true. However, if the set C_1 of metrizable codes is Borel, or equivalently if C_2 is Borel, then the set $\mathcal{L}_{\mathbf{f}}$ is Borel too. Indeed, let Φ be the second part of the disjunction of the expression in the proof of Theorem 16. Then it is easy to see, using the same arguments as in the

proof of Theorem 16, that

$$L \in \mathcal{L}_f \Leftrightarrow (L \in \Phi) \vee (\exists c \in C_1 \forall x \in D f_c(x) = \lim_{n \in L} f_n(x)).$$

Clearly the above formula gives a Σ_1^1 definition of \mathcal{L}_f , provided that C_1 is Borel.

(2) Besides the split interval, there exists another important example of a separable Rosenthal compact which is pre-metric of degree exactly two. This is the separable companion of the Alexandroff duplicate of the Cantor set $D(2^\mathbb{N})$ (see [To] for more details). An interesting feature of this compact is that it is *not* hereditarily separable.

Example 1. We proceed to give examples of pre-metric compacta of degree at most two for which Theorem 16 is not valid. Let us recall first the split Cantor set $S(2^\mathbb{N})$ which is simply the combinatorial analogue of the split interval. In the sequel by \leq_{lex} we shall denote the lexicographical ordering on $2^\mathbb{N}$ and by $<_{\text{lex}}$ its strict part. For every $x \in 2^\mathbb{N}$ we set $f_x^+ := \mathbf{1}_{\{y: x \leq_{\text{lex}} y\}}$ and $f_x^- := \mathbf{1}_{\{y: x <_{\text{lex}} y\}}$. The split Cantor set $S(2^\mathbb{N})$ is the set $\{f_x^+ : x \in 2^\mathbb{N}\} \cup \{f_x^- : x \in 2^\mathbb{N}\}$. Clearly, $S(2^\mathbb{N})$ is a hereditarily separable Rosenthal compact and it is a fundamental example of a pre-metric compactum of degree at most two (see [To]). There is a canonical dense sequence in $S(2^\mathbb{N})$ defined as follows. Fix a bijection $h: 2^{<\mathbb{N}} \rightarrow \mathbb{N}$ such that $h(s) < h(t)$ if $|s| < |t|$, and enumerate the nodes of Cantor tree as (s_n) according to h . For every $s \in 2^{<\mathbb{N}}$ let $x_s^0 = s \smallfrown 0^\infty \in 2^\mathbb{N}$ and $x_s^1 = s \smallfrown 1^\infty \in 2^\mathbb{N}$. For every $n \in \mathbb{N}$ let $f_{4n} = f_{x_{s_n}^0}^+$, $f_{4n+1} = f_{x_{s_n}^1}^+$, $f_{4n+2} = f_{x_{s_n}^0}^-$ and $f_{4n+3} = f_{x_{s_n}^1}^-$. The sequence (f_n) is a dense sequence in $S(2^\mathbb{N})$.

Let A be a subset of $2^\mathbb{N}$ and assume that A does not contain the eventually constant sequences. With every such A we associate, naturally, a subset of \mathbb{R}^A , which we will denote by $S(A)$, simply by restricting every function of $S(2^\mathbb{N})$ on A . Clearly, if A is Σ_1^1 , then $S(A)$ is again a hereditarily separable Rosenthal compact. Notice, however, that if $2^\mathbb{N} \setminus A$ is uncountable, then $S(A)$ is not of degree exactly two. The dense sequence (f_n) of $S(2^\mathbb{N})$ still remains a dense sequence in $S(A)$. Viewing (f_n) as a dense sequence in $S(A)$, we set

$$\mathcal{L}_A := \{L \in [\mathbb{N}]^\infty : (f_n|_A)_{n \in L} \text{ is pointwise convergent on } A\}.$$

We have the following proposition.

Proposition 17. *Let $A \subseteq 2^\mathbb{N}$ be Σ_1^1 . Assume that A does not contain the eventually constant sequences. Then $2^\mathbb{N} \setminus A$ is Wadge reducible to \mathcal{L}_A . In particular, if A is Σ_1^1 -complete, then \mathcal{L}_A is Π_1^1 -complete. Moreover, if A is Borel, then \mathcal{L}_A is Borel.*

Proof. Consider the map $\Phi: 2^\mathbb{N} \rightarrow 2^{2^{<\mathbb{N}}}$ defined by

$$\Phi(x) = \{x(0) + 1, x(0) \smallfrown (x(1) + 1), x(0) \smallfrown x(1) \smallfrown (x(2) + 1), \dots\}$$

where the above addition is taken modulo 2; Φ is continuous. Let $h: 2^{<\mathbb{N}} \rightarrow \mathbb{N}$ be the fixed enumeration of the nodes of the Cantor tree. For every $x \in 2^\mathbb{N}$ we set

$L_x := \{h(t) : t \in \Phi(x)\} \in [\mathbb{N}]^\infty$, and for every $t \in \Phi(x)$ let $x_t^0 := t \smallfrown 0^\infty \in 2^\mathbb{N}$. Notice that $(t_n)_{n \in L_x}$ is the enumeration of $\Phi(x)$ according to h and that x is the limit of the sequence $(x_{t_n}^0)_{n \in L_x}$. However, as one easily observes, if x is not eventually constant, then there exist infinitely many $n \in L_x$ such that $x_{t_n}^0 < x$ and infinitely many $n \in L_x$ such that $x < x_{t_n}^0$.

Now define $H : 2^\mathbb{N} \rightarrow [\mathbb{N}]^\infty$ by

$$H(x) = \{4h(t) : t \in \Phi(x)\} = \{4n : n \in L_x\}.$$

Clearly H is continuous. We claim that

$$x \notin A \Leftrightarrow H(x) \in \mathcal{L}_A.$$

Indeed, first assume that $x \notin A$. As we have already remarked, we have that $x = \lim_{n \in L_x} x_{t_n}^0$. Notice that the sequence $(f_n)_{n \in H(x)}$ is simply the sequence $(f_{x_t^0}^+)_{t \in \Phi(x)}$. Observe that for every $y \neq x$ the sequence $(f_n(y))_{n \in H(x)}$ converges to 0 if $y < x$ and to 1 if $x < y$. Since $x \notin A$, this implies that $(f_n)_{n \in H(x)}$ is pointwise convergent, and so $H(x) \in \mathcal{L}_A$. Conversely, assume that $x \in A$. Since A does not contain the eventually constant sequences, by the remarks after the definition of Φ , we see that there exist infinitely many $n \in H(x)$ such that $f_n(x) = 0$ and infinitely many $n \in H(x)$ such that $f_n(x) = 1$. Hence, the sequence $(f_n(x))_{n \in H(x)}$ does not converge, and since $x \in A$, we conclude that $H(x) \notin \mathcal{L}_A$. As H is continuous, this completes the proof the proof that $2^\mathbb{N} \setminus A$ is Wadge reducible to \mathcal{L}_A . Finally, the fact that if A is Borel, then \mathcal{L}_A is Borel too follows by straightforward descriptive set-theoretic computation and we prefer not to bother the reader with it. \square

Remark 6. Besides the fact that Theorem 16 cannot be lifted to all pre-metric compacta of degree at most two, Proposition 17 has another consequence. Namely, the fact that we cannot bound the Borel complexity of \mathcal{L}_f for a dense sequence $f = (f_n)$ in \mathcal{K} . This is in contrast with the situation with $\mathcal{L}_{f,f}$ for some $f \in \mathcal{K}$, which when it is Borel (equivalently, when f is a G_δ point), it is always Π_3^0 .

Concerning the class of not first countable separable Rosenthal compacta we have the following proposition.

Proposition 18. *Let \mathcal{K} be a separable Rosenthal compact. If there exists a non- G_δ point f in \mathcal{K} , then for every dense sequence $f = (f_n)$ in \mathcal{K} the set \mathcal{L}_f is Π_1^1 -complete.*

The proof of Proposition 18 is essentially based on a result of Krawczyk [Kr]. To state it, we need to recall some pieces of notation and some definitions. For every $a, b \in [\mathbb{N}]^\infty$ we write $a \subseteq^* b$ if $a \setminus b$ is finite, while we write $a \perp b$ if the set $a \cap b$ is finite. If \mathcal{A} is a subset of $[\mathbb{N}]^\infty$ we set $\mathcal{A}^\perp := \{b : b \perp a \ \forall a \in \mathcal{A}\}$ and $\mathcal{A}^* := \{\mathbb{N} \setminus a : a \in \mathcal{A}\}$. For every $\mathcal{A}, \mathcal{B} \subseteq [\mathbb{N}]^\infty$ we say that \mathcal{A} is *countably \mathcal{B} -generated* if there exists $\{b_n : n \in \mathbb{N}\} \subseteq \mathcal{B}$ such that for every $a \in \mathcal{A}$ there exists $k \in \mathbb{N}$ with $a \subseteq b_0 \cup \dots \cup b_k$. An ideal \mathcal{I} on \mathbb{N} is said to be *bisequential* if for every $p \in \beta\mathbb{N}$ with $\mathcal{I} \subseteq p^*$ we have that \mathcal{I} is countably p^* -generated. Finally, for every

$t \in \mathbb{N}^{<\mathbb{N}}$ set $\hat{t} := \{s : t \sqsubset s\}$. We will need the following result of Krawczyk (see [Kr, Lemma 2]).

Proposition 19. *Let \mathcal{I} be Σ_1^1 , bisquential and not countably \mathcal{I} -generated ideal on \mathbb{N} . Then there exists a 1-1 map $\psi : \mathbb{N}^{<\mathbb{N}} \rightarrow \mathbb{N}$ such that, setting*

$$\mathcal{J} := \{\psi^{-1}(a) : a \in \mathcal{I}\},$$

the following hold.

- (P1) *For every $\sigma \in \mathbb{N}^{\mathbb{N}}$ we have $\{\sigma|n : n \in \mathbb{N}\} \in \mathcal{J}$.*
- (P2) *For every $b \in \mathcal{J}$ and every $n \in \mathbb{N}$ there exist $t_0, \dots, t_k \in \mathbb{N}^n$ with $b \subseteq^* \hat{t}_0 \cup \dots \cup \hat{t}_k$.*

We proceed to the proof of Proposition 18.

Proof of Proposition 18. Let $\mathbf{f} = (f_n)$ be a dense sequence in \mathcal{K} and let $f \in \mathcal{K}$ be a non- G_δ point. Consider the ideal

$$\mathcal{I} := \{L \in [\mathbb{N}]^\infty : f \notin \overline{\{f_n\}_{n \in L}}^p\}.$$

In [Kr, page 1099], it is shown that \mathcal{I} is a Σ_1^1 , bisquential ideal on \mathbb{N} which is not countably \mathcal{I} -generated (the bisquentiality of \mathcal{I} can be derived either by a result of Pol [P3], or by the non-effective version of Debs' theorem [AGR]). Also, by the Bourgain–Fremlin–Talagrand theorem, we have that $\mathcal{I}^\perp = \mathcal{L}_{\mathbf{f},f}$. We apply Proposition 19 and we obtain a 1-1 map $\psi : \mathbb{N}^{<\mathbb{N}} \rightarrow \mathbb{N}$ satisfying (P1) and (P2).

Claim. *For every infinite $T \in \text{WF}$ we have $T \in \mathcal{J}^\perp$.*

Proof of the claim. Assume not. Then there exist $T \in \text{WF}$ infinite and $b \in \mathcal{J}$ with $b \subseteq T$. For every $s \in T$ let $T_s = \{t \in T : s \sqsubseteq t\}$. Set $S := \{s \in T : T_s \cap b \text{ is infinite}\}$ and notice that S is downwards closed subtree of T . Moreover, by (P2) in Proposition 19, we see that S is finitely splitting. Finally, observe that for every $s \in S$ there exists $n \in \mathbb{N}$ with $s \frown n \in S$. Indeed, let $s \in S$ and set $b_s := T_s \cap b \in \mathcal{J}$. Let $N_s = \{n \in \mathbb{N} : s \frown n \in T\}$ and observe that $b_s \setminus \{s\} = \bigcup_{n \in N_s} (T_{s \frown n} \cap b_s)$. By (P2) in Proposition 19 again, we see that there exists $n_0 \in N_s$ with $T_{s \frown n_0} \cap b_s$ infinite. Thus $s \frown n_0 \in S$. It follows that S is a finitely splitting, infinite tree. By König's lemma, we conclude that $S \notin \text{WF}$. But then $T \notin \text{WF}$, a contradiction. The claim is proved. \square

Fix $T_0 \in \text{WF}$ infinite. The map $\Psi : \text{Tr} \rightarrow [\mathbb{N}]^\infty$ defined by the rule $\Psi(T) = \{\psi(t) : t \in T \cup T_0\}$ is clearly continuous. If $T \in \text{WF}$, then $T \cup T_0 \in \text{WF}$. By the above claim, we see that $T \cup T_0 \in \mathcal{J}^\perp$, and so, $\Psi(T) \in \mathcal{I}^\perp = \mathcal{L}_{\mathbf{f},f} \subseteq \mathcal{L}_{\mathbf{f}}$. On the other hand, if $T \notin \text{WF}$, then by (P1) in Proposition 19 and the above claim, there exist $L \in \mathcal{L}_{\mathbf{f}} \setminus \mathcal{L}_{\mathbf{f},f}$ and $M \in \mathcal{L}_{\mathbf{f},f}$ with $L \cup M \subseteq \Psi(T)$. Therefore, $\Psi(T) \notin \mathcal{L}_{\mathbf{f}}$. It follows that WF is Wadge reducible to $\mathcal{L}_{\mathbf{f}}$ and the proof is completed. \square

The last part of this section is devoted to the construction of canonical Π_1^1 -ranks on the sets $\mathcal{L}_{\mathbf{f}}$ and $\mathcal{L}_{\mathbf{f},f}$. So let X be a Polish space, let $\mathbf{f} = (f_n)$ be a sequence relatively compact in $\mathcal{B}_1(X)$ and let f be an accumulation point of (f_n) . Since the sets $\mathcal{L}_{\mathbf{f}}$ and $\mathcal{L}_{\mathbf{f},f}$ do not depend on the topology on X , we may assume, by enlarging the topology of X if necessary, that the functions (f_n) and the function f are continuous (see [Ke]). We need to deal with decreasing sequences of closed subsets of X à la Cantor. We fix a countable dense subset D of X . Let (B_n) be an enumeration of all closed balls in X (taken with respect to some compatible complete metric) with centers in D and rational radii. If X happens to be locally compact, we will assume that every ball B_n is compact. We say that a finite sequence $w = (l_0, \dots, l_k) \in \mathbb{N}^{<\mathbb{N}}$ is acceptable if

- (i) $B_{l_0} \supseteq B_{l_1} \supseteq \dots \supseteq B_{l_k}$, and
- (ii) $\text{diam}(B_{l_i}) \leq \frac{1}{i+1}$ for every $i \in \{0, \dots, k\}$.

By convention \emptyset is acceptable. Notice that if $w_1 \sqsubset w_2$ and w_2 is acceptable, then w_1 is acceptable too. We also need to introduce some pieces of notation.

Notation. By Fin we denote the set of all finite subsets of \mathbb{N} . For every $F, G \in \text{Fin}$ we write $F < G$ if $\max\{n : n \in F\} < \min\{n : n \in G\}$. For every $L \in [\mathbb{N}]^\infty$ by $\text{Fin}(L)$ we denote the set of all finite subsets of L . Finally, by $[\text{Fin}(L)]^{<\mathbb{N}}$ we denote the set of all finite sequences $t = (F_0, \dots, F_k) \in (\text{Fin}(L))^{<\mathbb{N}}$ which are increasing, that is, $F_0 < F_1 < \dots < F_k$.

We will construct the Π_1^1 -ranks on $\mathcal{L}_{\mathbf{f}}$ and $\mathcal{L}_{\mathbf{f},f}$ by finding appropriate reductions of the sets in question to well-founded trees. In particular, we shall construct

- (C1) a continuous map $[\mathbb{N}]^\infty \ni L \mapsto T_L \in \text{Tr}(\mathbb{N} \times \text{Fin} \times \mathbb{N})$, and
- (C2) a continuous map $[\mathbb{N}]^\infty \ni L \mapsto S_L \in \text{Tr}(\mathbb{N} \times \mathbb{N})$

such that

- (C3) $L \in \mathcal{L}_{\mathbf{f}}$ if and only if $T_L \in \text{WF}(\mathbb{N} \times \text{Fin} \times \mathbb{N})$, and
- (C4) $L \in \mathcal{L}_{\mathbf{f},f}$ if and only if $S_L \in \text{WF}(\mathbb{N} \times \mathbb{N})$.

By (C1)–(C4), it follows that the maps $L \mapsto o(T_L)$ and $L \mapsto o(S_L)$ are Π_1^1 -ranks on $\mathcal{L}_{\mathbf{f}}$ and $\mathcal{L}_{\mathbf{f},f}$.

5.1. The reduction of $\mathcal{L}_{\mathbf{f}}$ to $\text{WF}(\mathbb{N} \times \text{Fin} \times \mathbb{N})$. Let $d \in \mathbb{N}$. For every $L \in [\mathbb{N}]^\infty$ we define $T_L^d \in \text{Tr}(\mathbb{N} \times \text{Fin} \times \mathbb{N})$ by the rule

$$\begin{aligned}
 T_L^d = \{ (s, t, w) \quad & : \quad \exists k \text{ with } |s| = |t| = |w| = k, \\
 & s = (n_0 < \dots < n_{k-1}) \in [L]^{<\mathbb{N}}, \\
 & t = (F_0 < \dots < F_{k-1}) \in [\text{Fin}(L)]^{<\mathbb{N}}, \\
 & w = (l_0, \dots, l_{k-1}) \in \mathbb{N}^{<\mathbb{N}} \text{ is acceptable, and} \\
 & \forall 0 \leq i \leq k-1 \quad \forall z \in B_{l_i} \text{ there exists } m_i \in F_i \text{ with} \\
 & |f_{n_i}(z) - f_{m_i}(z)| > \frac{1}{d+1} \}.
 \end{aligned}$$

Next we “glue” the sequence of trees $\langle T_L^d : d \in \mathbb{N} \rangle$ in a natural way and we define a tree $T_L \in \text{Tr}(\mathbb{N} \times \text{Fin} \times \mathbb{N})$ by

$$(s, t, w) \in T_L \iff \exists d \exists (s', t', w') \text{ such that } (s', t', w') \in T_L^d \text{ and } s = d^\frown s', t = \{d\}^\frown t', w = d^\frown w'.$$

It is clear that the map $[\mathbb{N}]^\infty \ni L \mapsto T_L \in \text{Tr}(\mathbb{N} \times \text{Fin} \times \mathbb{N})$ is continuous. Moreover, the following holds.

Lemma 20. *Let $L \in [\mathbb{N}]^\infty$. Then $L \in \mathcal{L}_f$ if and only if $T_L \in \text{WF}(\mathbb{N} \times \text{Fin} \times \mathbb{N})$.*

Proof. Notice, first, that if $L \notin \mathcal{L}_f$, then there exist $L_1, L_2 \in [L]^\infty$ with $L_1 \cap L_2 = \emptyset$, $L_1, L_2 \in \mathcal{L}_f$ and $f_{L_1} \neq f_{L_2}$ where f_{L_1} and f_{L_2} are the pointwise limits of the sequences $(f_n)_{n \in L_1}$ and $(f_n)_{n \in L_2}$ respectively. We select $x \in X$ and $d \in \mathbb{N}$ such that $|f_{L_1}(x) - f_{L_2}(x)| > \frac{1}{d+1}$. Clearly, we may assume that $|f_n(x) - f_m(x)| > \frac{1}{d+1}$ for every $n \in L_1$ and $m \in L_2$. Let $L_1 = \{n_0 < n_1 < \dots\}$ and $L_2 = \{m_0 < m_1 < \dots\}$ be the increasing enumerations of L_1 and L_2 . Using the continuity of the functions (f_n) we select $w = (l_0, \dots, l_k, \dots) \in \mathbb{N}^\mathbb{N}$ such that: (i) $w|k$ is acceptable for every $k \in \mathbb{N}$, (ii) $\bigcap_k B_{l_k} = \{x\}$, and (iii) $|f_{n_k}(z) - f_{m_k}(z)| > \frac{1}{d+1}$ for every $k \in \mathbb{N}$ and every $z \in B_{l_k}$. Therefore, $((n_0, \dots, n_k), (\{m_0\}, \dots, \{m_k\}), w|k) \in T_L^d$ for every $k \in \mathbb{N}$ which implies that $T_L \notin \text{WF}(\mathbb{N} \times \text{Fin} \times \mathbb{N})$.

Conversely, assume that T_L is not well-founded and let $d \in \mathbb{N}$ such that T_L^d is not well-founded. Let $\langle (s_k, t_k, w_k) : k \in \mathbb{N} \rangle$ be an infinite branch of T_L^d . We set $N := \bigcup_k s_k = \{n_0 < \dots\} \in [L]^\infty$, $\mathcal{F} := \bigcup_k t_k = (F_0 < \dots) \in \text{Fin}(L)^\mathbb{N}$ and $w := \bigcup_k w_k = (l_0, \dots) \in \mathbb{N}^\mathbb{N}$. By the definition of T_L^d , we have $\bigcap_k B_{l_k} = \{x\} \in X$; moreover, for every $k \in \mathbb{N}$ there exists $m_k \in F_k \subseteq L$ with $|f_{n_k}(x) - f_{m_k}(x)| > \frac{1}{d+1}$. Since $F_i < F_j$ for every $i < j$, we see that $m_i < m_j$ if $i < j$. It follows that $M = \{m_0 < \dots < m_k < \dots\} \in [L]^\infty$. Thus, the sequence $(f_n(x))_{n \in L}$ is not Cauchy and so $L \notin \mathcal{L}_f$, as desired. \square

By Lemma 20, the reduction of \mathcal{L}_f to $\text{WF}(\mathbb{N} \times \text{Fin} \times \mathbb{N})$ is constructed. Notice that for every $L \in \mathcal{L}_f$ and every $d_1 \leq d_2$ we have $o(T_L^{d_1}) \leq o(T_L^{d_2})$ and, moreover, $o(T_L) = \sup\{o(T_L^d) : d \in \mathbb{N}\} + 1$.

Remark 7. We should point out that the reason why in the definition of T_L^d the node t is a finite sequence of finite sets rather than natural numbers, is to obtain the estimate in Proposition 22 below. Having natural numbers instead of finite sets would also lead to a canonical rank.

5.2. The reduction of $\mathcal{L}_{f,f}$ to $\text{WF}(\mathbb{N} \times \mathbb{N})$. The reduction is similar to that of the previous step, and so, we shall indicate only the necessary changes. Let $d \in \mathbb{N}$.

As before, with every $L \in [\mathbb{N}]^\infty$ we associate a tree $S_L^d \in \text{Tr}(\mathbb{N} \times \mathbb{N})$ by setting

$$\begin{aligned} S_L^d = \{ (s, w) : & \exists k \in \mathbb{N} \text{ with } |s| = |w| = k, \\ & s = (n_0 < \dots < n_{k-1}) \in [L]^{<\mathbb{N}}, \\ & w = (l_0, \dots, l_{k-1}) \in \mathbb{N}^{<\mathbb{N}} \text{ is acceptable and} \\ & \forall 0 \leq i \leq k-1 \forall z \in B_{l_i} \text{ we have } |f_{n_i}(z) - f(z)| > \frac{1}{d+1} \}. \end{aligned}$$

Next, we “glue” the sequence of trees $\langle S_L^d : d \in \mathbb{N} \rangle$ as we did with the sequence $\langle T_L^d : d \in \mathbb{N} \rangle$ and we define a tree $S_L \in \text{Tr}(\mathbb{N} \times \mathbb{N})$ by the rule

$$\begin{aligned} (s, w) \in S_L & \Leftrightarrow \exists d \exists (s', w') \text{ such that } (s', w') \in T_L^d \text{ and} \\ & s = d \smallfrown s', w = d \smallfrown w'. \end{aligned}$$

Again, it is easy to check that the map $[\mathbb{N}]^\infty \ni L \mapsto S_L \in \text{Tr}(\mathbb{N} \times \mathbb{N})$ is continuous. Moreover, we have the following analogue of Lemma 20. The proof is identical and is left to the reader.

Lemma 21. *Let $L \in [\mathbb{N}]^\infty$. Then $L \in \mathcal{L}_{\mathbf{f},f}$ if and only if $S_L \in \text{WF}(\mathbb{N} \times \mathbb{N})$.*

This gives us the reduction of $\mathcal{L}_{\mathbf{f},f}$ to $\text{WF}(\mathbb{N} \times \mathbb{N})$. As before, we have that $o(S_L) = \sup\{o(S_L^d) : d \in \mathbb{N}\} + 1$ for every $L \in \mathcal{L}_{\mathbf{f},f}$.

5.3. We proceed to discuss the question whether for a given $L \in \mathcal{L}_{\mathbf{f},f}$ we can bound the order of the tree S_L by the order of T_L . The following example shows that this is not possible in general.

Example 2. Let $A(2^\mathbb{N}) := \{\delta_\sigma : \sigma \in 2^\mathbb{N}\} \cup \{0\}$ denote the one point compactification of $2^\mathbb{N}$. This is not a separable Rosenthal compact, but it can be supplemented to one in a standard way (see [P1, Ma, To]). Specifically, let (s_n) be the enumeration of the Cantor tree $2^{<\mathbb{N}}$ as in Example 1. For every $n \in \mathbb{N}$ set $f_n := \mathbf{1}_{V_{s_n}}$ where $V_{s_n} = \{\sigma \in 2^\mathbb{N} : s_n \sqsubset \sigma\}$. Then $A(2^\mathbb{N}) \cup \{f_n\}$ is a separable Rosenthal compact. Now let A be a Σ_1^1 non-Borel subset of $2^\mathbb{N}$. Following [P2] (see, also, [Ma]), let \mathcal{K}_A be the separable Rosenthal compact obtained by restricting every function in $A(2^\mathbb{N}) \cup \{f_n\}$ on A . The sequence $\mathbf{f}_A = (f_n|_A)$ is a countable dense subset of \mathcal{K}_A consisting of continuous functions and $0 \in \mathcal{K}_A$ is a non- G_δ point (and obviously continuous). Consider the sets

$$\mathcal{L}_{\mathbf{f}}^A := \{L \in [\mathbb{N}]^\infty : (f_n|_A)_{n \in L} \text{ is pointwise convergent on } A\}$$

and

$$\mathcal{L}_{\mathbf{f},0}^A := \{L \in [\mathbb{N}]^\infty : (f_n|_A)_{n \in L} \text{ is pointwise convergent to } 0 \text{ on } A\}.$$

Let ϕ be a Π_1^1 -rank on $\mathcal{L}_{\mathbf{f}}^A$ and let ψ be a Π_1^1 -rank on $\mathcal{L}_{\mathbf{f},0}^A$. We claim that there does not exist a map $\Phi : \omega_1 \rightarrow \omega_1$ such that $\psi(L) \leq \Phi(\phi(L))$ for every $L \in \mathcal{L}_{\mathbf{f},0}^A$. Assume not. Set

$$R := \{L \in [\mathbb{N}]^\infty : \exists \sigma \in 2^\mathbb{N} \text{ with } s_n \sqsubset \sigma \forall n \in L\}.$$

Then R is a closed subset of $\mathcal{L}_{\mathbf{f}}^A$. For every $L \in R$ let $\sigma_L = \bigcup_{n \in L} s_n \in 2^{\mathbb{N}}$. The map $R \ni L \mapsto \sigma_L \in 2^{\mathbb{N}}$ is clearly continuous. Observe that for every $L \in R$ we have that $L \in \mathcal{L}_{\mathbf{f},0}^A$ if and only if $\sigma_L \notin A$. Since R is a Borel subset of $\mathcal{L}_{\mathbf{f}}^A$, by boundedness, we obtain that $\sup\{\phi(L) : L \in R\} = \xi < \omega_1$. Let $\zeta = \sup\{\Phi(\lambda) : \lambda \leq \xi\}$. The set $B := R \cap \{L \in \mathcal{L}_{\mathbf{f},0}^A : \psi(L) \leq \zeta\}$ is Borel and $B = R \cap \mathcal{L}_{\mathbf{f},0}^A$. Hence, the set $\Sigma_B := \{\sigma_L : L \in B\}$ is an analytic subset of $2^{\mathbb{N}} \setminus A$. Since $2^{\mathbb{N}} \setminus A$ is $\mathbf{\Pi}_1^1$ -true, there exists $\sigma_0 \in 2^{\mathbb{N}} \setminus A$ with $\sigma_0 \notin \Sigma_B$. We select $L \in R$ with $\sigma_L = \sigma_0$. Then $L \in B$ yet $\sigma_L \notin \Sigma_B$, a contradiction.

Although we cannot, in general, bound the order of the tree S_L by that of T_L , the following proposition shows that this is possible for an important special case.

Proposition 22. *Let X be a locally compact Polish space, let \mathcal{K} a separable Rosenthal compact on X , let $\mathbf{f} = (f_n)$ a dense sequence in \mathcal{K} consisting of continuous functions and let $f \in \mathcal{K}$. If f is continuous, then $o(S_L) \leq o(T_L)$ for every $L \in \mathcal{L}_{\mathbf{f},f}$.*

In particular, there exist a $\mathbf{\Pi}_1^1$ -rank ϕ on $\mathcal{L}_{\mathbf{f}}$ and a $\mathbf{\Pi}_1^1$ -rank ψ on $\mathcal{L}_{\mathbf{f},f}$ with $\psi(L) \leq \phi(L)$ for every $L \in \mathcal{L}_{\mathbf{f},f}$.

Proof. We will show that for every $d \in \mathbb{N}$ we have $o(S_L^d) \leq o(T_L^d)$ for every $L \in \mathcal{L}_{\mathbf{f},f}$. This clearly completes the proof. So fix $d \in \mathbb{N}$ and $L \in \mathcal{L}_{\mathbf{f},f}$. We shall construct a monotone map $\Phi: S_L^d \rightarrow [\text{Fin}(L)]^{<\mathbb{N}}$ such that for every $(s, w) \in S_L^d$ the following hold.

- (i) We have $|(s, w)| = |\Phi((s, w))|$.
- (ii) If $s = (n_0 < \dots < n_k)$, $w = (l_0, \dots, l_k)$ and $\Phi((s, w)) = (F_0 < \dots < F_k)$, then for every $i \in \{0, \dots, k\}$ and every $z \in B_{l_i}$ there exists $m_i \in F_i$ with $|f_{n_i}(z) - f_{m_i}(z)| > \frac{1}{d+1}$.

Assuming that Φ has been constructed, let $M: S_L^d \rightarrow T_L^d$ be defined by

$$M((s, w)) = (s, \Phi((s, w)), w).$$

It is easy to see that M is a well-defined monotone map, and so, $o(S_L^d) \leq o(T_L^d)$ as desired.

The construction of the map Φ proceeds by recursion on the length of (s, w) . We set $\Phi((\emptyset, \emptyset)) = \emptyset$. Let $k \in \mathbb{N}$ and assume that $\Phi((s, w))$ has been defined for every $(s, w) \in S_L^d$ with $|(s, w)| \leq k$. Let $(s', w') = (s \hat{\ } n_k, w \hat{\ } l_k) \in S_L^d$ with $|s'| = |w'| = k + 1$. By the definition of S_L^d , we have that $|f_{n_k}(z) - f(z)| > \frac{1}{d}$ for every $z \in B_{l_k}$. We set $p = \max\{n : n \in F \text{ and } F \in \Phi((s, w))\} \in \mathbb{N}$. For every $z \in B_{l_k}$ we select $m_z \in L$ with $m_z > p$ and such that $|f_{n_k}(z) - f_{m_z}(z)| > \frac{1}{d+1}$. Since the functions (f_n) are continuous, we may select an open neighborhood U_z of z such that $|f_{n_k}(y) - f_{m_z}(y)| > \frac{1}{d+1}$ for every $y \in U_z$. By the compactness of B_{l_k} , there exist $z_0, \dots, z_{j_k} \in B_{l_k}$ such that $U_{z_0} \cup \dots \cup U_{z_{j_k}} \supseteq B_{l_k}$. Set

$$F_k := \{m_{z_i} : i \in \{0, \dots, j_k\}\} \in \text{Fin}(L)$$

and notice that $F \leq p < F_k$ for every $F \in \Phi((s, w))$. We set

$$\Phi((s', w')) := \Phi((s, w)) \smallfrown F_k \in [\text{Fin}(L)]^{<\mathbb{N}}.$$

It is easy to check that $\Phi((s', w'))$ satisfies (i) and (ii) above. The proof is completed. \square

REFERENCES

- [AD] S. A. Argyros and P. Dodos, *Genericity and amalgamation of classes of Banach spaces*, Adv. Math. 209 (2007), 666–748.
- [ADK] S. A. Argyros, P. Dodos and V. Kanellopoulos, *Tree structures associated to a family of functions*, Journal Symb. Logic 70 (2005), 681–695.
- [AGR] S. A. Argyros, G. Godefroy and H. P. Rosenthal, *Descriptive Set Theory and Banach spaces*, Handbook of the Geometry of Banach Spaces, Vol. 2, edited by W. B. Johnson and J. Lindenstrauss, Elsevier, 2003.
- [Be] H. Becker, *Pointwise limits of subsequences and Σ_2^1 sets*, Fund. Math. 128 (1987), 159–170.
- [Bo] J. Bourgain, *On convergent sequences of continuous functions*, Bull. Soc. Math. Belg. 32 (1980), 235–249.
- [BFT] J. Bourgain, D. H. Fremlin and M. Talagrand, *Pointwise compact sets of Baire-measurable functions*, Amer. J. Math. 100 (1978), 845–886.
- [De] G. Debs, *Effective properties in compact sets of Borel functions*, Mathematika 34 (1987), 64–68.
- [Go] G. Godefroy, *Compacts de Rosenthal*, Pacific J. Math. 91 (1980), 293–306.
- [Ke] A. S. Kechris, *Classical Descriptive Set Theory*, Grad. Texts in Math., Vol. 156, Springer-Verlag, 1995.
- [KL] A. S. Kechris and A. Louveau, *A classification of Baire class 1 functions*, Trans. Amer. Math. Soc. 318 (1990), 209–236.
- [Kr] A. Krawczyk, *Rosenthal compacta and analytic sets*, Proc. Amer. Math. Soc. 115 (1992), 1095–1100.
- [LT] J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces I and II*, Springer, 1996.
- [Ma] W. Marciszewski, *On a classification of pointwise compact sets of the first Baire class*, Fund. Math. 133 (1989), 195–209.
- [MK] D. A. Martin and A. S. Kechris, *Infinite games and effective descriptive set theory*, in “Analytic Sets”, editors C. A. Rogers et al., Academic Press, 1980.
- [Mi] A. W. Miller, *Problem list*, available at <http://www.math.wisc.edu/~miller>.
- [Mo] Y. N. Moschovakis, *Descriptive Set Theory*, North Holland, Amsterdam, 1980.
- [OR] E. Odell and H. P. Rosenthal, *A double-dual characterization of separable Banach spaces not containing ℓ_1* , Israel J. Math. 20 (1975), 375–384.
- [P1] R. Pol, *On pointwise and weak topology in function spaces*, Warsaw University, 1984.
- [P2] R. Pol, *Note on compact sets of first Baire class functions*, Proc. Amer. Math. Soc. 96 (1986), 152–154.
- [P3] R. Pol, *Note on pointwise convergence of sequences of analytic sets*, Mathematika 36 (1989), 290–300.
- [Ro1] H. P. Rosenthal, *Pointwise compact subsets of the first Baire class*, Amer. J. Math. 99 (1977), 362–378.
- [Ro2] H. P. Rosenthal, *Some recent discoveries in the isomorphic theory of Banach spaces*, Bull. Amer. Math. Soc. 84 (1978), 803–831.
- [S1] J. H. Silver, *Every analytic set is Ramsey*, Journal Symb. Logic 35 (1970), 60–64.

- [S2] J. H. Silver, *Counting the number of equivalence classes of Borel and coanalytic equivalence relations*, Ann. Math. Logic 18 (1980), 1–28.
- [To] S. Todorćević, *Compact subsets of the first Baire class*, J. Amer. Math. Soc. 12 (1999), 1179–1212.

NATIONAL TECHNICAL UNIVERSITY OF ATHENS, FACULTY OF APPLIED SCIENCES, DEPARTMENT
OF MATHEMATICS, ZOGRAFOU CAMPUS, 157 80, ATHENS, GREECE.

E-mail address: `pdodos@math.ntua.gr`