

# UNCONDITIONAL FAMILIES IN BANACH SPACES

SPIROS ARGYROS, PANDELIS DODOS AND VASSILIS KANELLOPOULOS

ABSTRACT. It is shown that for every separable Banach space  $X$  with non-separable dual, the space  $X^{**}$  contains an unconditional family of size  $|X^{**}|$ . The proof is based on Ramsey theory for trees and finite products of perfect sets of reals. Among its consequences, it is proved that every dual Banach space has a separable quotient.

## 1. INTRODUCTION

The problem of the existence of an unconditional basic sequence in every, infinite dimensional, Banach space was a central one and remained open for many years. At the beginning of 1990s Gowers and Maurey [GM] settled that problem in the negative. Their celebrated example led to the profound concept, introduced by Johnson, of Hereditarily Indecomposable (HI) spaces, which completely changed our understanding of the structure of Banach spaces. The class of HI spaces stands in the opposite of the class of spaces with an unconditional basis and Gowers' dichotomy [G1], a Ramsey theoretic principle for Banach spaces, yields that every Banach space either is HI saturated, or contains an unconditional basic sequence. Further investigation, by several authors, has shown that HI spaces occur almost everywhere and this indicates the difficulty to obtain positive results concerning the existence of unconditional sequences.

The aim of the present work is to prove a theorem, of unexpected generality, providing unconditional families in the second dual of a separable Banach space, and also, to present some of its consequences. More precisely, the following theorem is proved.

**Theorem 1.** *Let  $X$  be a separable Banach space not containing  $\ell_1$  and such that  $X^*$  is non-separable. Then there exists a bounded bi-orthogonal system  $\{(z_\sigma^*, z_\sigma^{**}) : \sigma \in 2^{\mathbb{N}}\}$  in  $X^* \times X^{**}$  such that the family  $\{z_\sigma^{**} : \sigma \in 2^{\mathbb{N}}\}$  is 1-unconditional, weak\* discrete and has 0 as the unique weak\* accumulation point.*

A rather direct consequence is the following.

*The second dual  $X^{**}$  of a separable space  $X$  with non-separable dual contains an unconditional family of size  $|X^{**}|$ .*

---

2000 *Mathematics Subject Classification*: 05C05, 05D10, 46B03, 46B20, 46B26.

*Keywords*: Ramsey theory, chains and antichains of trees, dual Banach spaces, unconditional bases, quotient of Banach spaces.

We also obtain a trichotomy, answering affirmatively the “unconditionality or reflexivity problem”, which is stated as follows.

*Every separable space  $X$  either is reflexively saturated, or one of its second or third dual contains an unconditional family of cardinality equal to the size of the corresponding dual.*

A third application concerns the classical “separable quotient problem”, posed by Banach, and settles the problem for the class of spaces isomorphic to a dual. In particular, the following is shown.

*Every dual Banach space has a separable quotient.*

Let us point out that Theorem 1, as well as the aforementioned trichotomy, are sharp. Indeed, there are separable spaces with separable dual and non-separable HI second dual [ATo]. Moreover, such a space  $X$  can be chosen so that  $X$ ,  $X^*$  and  $X^{**}$  are all HI and not containing a reflexive subspace [AAT]. Let us also note the stability of the unconditionality constant obtained by Theorem 1 which remains the best possible for any equivalent norm on  $X$ . This could be compared to Maurey’s theorem [Mau] concerning second dual types in separable Banach spaces containing  $\ell_1$ . The Odell–Rosenthal theorem [OR] permits us to lift structure from the 1-unconditional family  $\{z_\sigma^{**} : \sigma \in 2^{\mathbb{N}}\}$  into the space  $X$  itself. This is the content of the following theorem which corresponds to Theorem 18 in the main text.

**Theorem 2.** *Let  $X$  be as in Theorem 1. Then there exists a Schauder tree basis  $(w_t)_{t \in 2^{<\mathbb{N}}}$  in  $X$  such that the following are satisfied.*

- (1) *For every  $n \geq 1$  the finite family  $\{w_t : t \in 2^n\}$  is  $(1 + \frac{1}{n})$ -unconditional.*
- (2) *For every  $n, m \in \mathbb{N}$  with  $1 \leq n < m$  and every  $\{s_t : t \in 2^m\} \subseteq 2^m$  with  $t \sqsubset s_t$  for all  $t \in 2^n$ , the families  $\{w_t : t \in 2^n\}$  and  $\{w_{s_t} : t \in 2^n\}$  are  $(1 + \frac{1}{n})$ -equivalent under the natural correspondence.*
- (3) *For every infinite chain  $(t_n)$  of  $2^{<\mathbb{N}}$  the sequence  $(w_{t_n})$  is weak Cauchy, and for every infinite antichain  $(s_n)$  of  $2^{<\mathbb{N}}$  the sequence  $(w_{s_n})$  is weakly-null.*

This result reveals the generic character of the basis of the James Tree space  $JT$ , the first example of a separable Banach space not containing  $\ell_1$  and with non-separable dual ([J]). For further applications of Theorem 1 we refer to [AAK].

The ingredients for proving Theorem 1 are mainly Ramsey theoretical. In particular, we use results concerning definable partitions of certain classes of antichains of the dyadic tree, which we call increasing and decreasing, as well as, definable partitions of finite products of perfect sets. Theorem 12, extracted from Stegall’s fundamental construction [St] for separable Banach spaces with non-separable dual, also plays a key role. More precisely, using the Ramsey properties of increasing and decreasing antichains, proved in [ADK], we obtain the following extension of Stern’s theorem [Ste] (see §2 for unexplained terminology).

**Theorem 3.** *Let  $X$  be a separable Banach space and let  $\Delta = \{x_t : t \in 2^{<\mathbb{N}}\}$  be a bounded family in  $X$ . Then there exists a regular dyadic subtree  $T$  of  $2^{<\mathbb{N}}$  such that the following are satisfied.*

- (1) *Either, (i) there exists  $C > 0$  such that for every infinite chain  $(t_n)$  of  $T$  the sequence  $(x_{t_n})$  is  $C$ -equivalent to the standard basis of  $\ell_1$ , or (ii) for every infinite chain  $(t_n)$  of  $T$  the sequence  $(x_{t_n})$  is weak Cauchy.*
- (2) *Either, (i) there exists  $C > 0$  such that for every increasing antichain  $(t_n)$  of  $T$  the sequence  $(x_{t_n})$  is  $C$ -equivalent to the standard basis of  $\ell_1$ , or (ii) for every increasing antichain  $(t_n)$  of  $T$  the sequence  $(x_{t_n})$  is weak Cauchy. Moreover, for every pair  $(t_n)$  and  $(s_n)$  of increasing antichains of  $T$  with the same limit point in  $2^{\mathbb{N}}$ , the sequences  $(x_{t_n})$  and  $(x_{s_n})$  are both weak\* convergent to the same element of  $X^{**}$ .*
- (3) *Similar to (2) for the decreasing antichains.*

We should point out that part (1.i) of Theorem 3 does not necessarily imply part (2.i), or conversely (see Remark 1 in the main text). Theorem 3 incorporates all the machinery of Ramsey theory for trees needed for the proof of Theorem 1, which proceeds as follows. For a separable Banach space  $X$  with non-separable dual, Theorem 12 yields that there exist a bounded family  $\{x_t\}_{t \in 2^{<\mathbb{N}}}$  in  $X$  and a bounded family  $\{x_\sigma^* : \sigma \in 2^{\mathbb{N}}\}$  in  $X^*$  such that for every  $\sigma, \tau \in 2^{\mathbb{N}}$  and every weak\* accumulation point  $x_\sigma^{**}$  of  $(x_{\sigma|n})$  we have  $x_\sigma^{**}(x_\tau^*) = \delta_{\sigma\tau}$ . Next, applying Theorem 3 and taking into account that  $\ell_1$  does not embed into  $X$ , we obtain a regular dyadic subtree  $T$  of  $2^{<\mathbb{N}}$  and to each  $\sigma$  in the body  $[\hat{T}]$  of  $T$  a triplet  $\{x_\sigma^0, x_\sigma^+, x_\sigma^-\}$  in  $X^{**}$  associated to the unique weak\* limit points along subsequences of  $\{x_t\}_{t \in T}$  determined by chains, increasing and decreasing antichains. The key observation is that the family  $\{z_\sigma^{**} = x_\sigma^0 - x_\sigma^+ : \sigma \in [\hat{T}]\}$  is weak\* discrete having 0 as the unique weak\* accumulation point. Moreover, for every  $\sigma, \tau \in [\hat{T}]$  we have  $z_\sigma^{**}(x_\tau^*) = \delta_{\sigma\tau}$ . The final step in the proof of Theorem 1 is the perfect unconditionality theorem, stated as follows.

**Theorem 4.** *Let  $X$  be a separable Banach space. Also let  $Q$  be a perfect subset of  $2^{\mathbb{N}}$  and let  $\mathcal{D} = \{z_\sigma^{**} : \sigma \in Q\}$  be a bounded family in  $X^{**}$  which is weak\* discrete and has 0 as the unique weak\* accumulation point. Assume that the map  $\Phi: Q \times (B_{X^*}, w^*) \rightarrow \mathbb{R}$  defined by  $\Phi(\sigma, x^*) = z_\sigma^{**}(x^*)$  is Borel. Then there exists a perfect subset  $R$  of  $Q$  such that the family  $\{z_\sigma^{**} : \sigma \in R\}$  is 1-unconditional.*

The construction of the perfect subset  $R$  in Theorem 4 is done by induction and by repeated applications of a partition theorem due to Galvin. Note that the Borelness of the function  $\Phi$  is crucial for the proof, as it is used to show that certain partitions are definable. One could not expect a similar result for an arbitrary uncountable family as above. Indeed, there exists an uncountable weakly discrete family accumulating to 0 in a reflexive and HI saturated space  $X$  (see [AT]).

## 2. THE RAMSEY THEORETICAL BACKGROUND

The aim of this section is to review the Ramsey theoretical background needed in the sequel. There is a long history on the interaction between infinite dimensional Ramsey theory and Banach space theory, going back to Farahat's proof [F] of Rosenthal's  $\ell_1$  theorem [Ro]. We refer the reader to the survey papers [Od] and [G2] for an account of related results.

The component of Ramsey theory we will use, concerns partitions of infinite subsets of the dyadic tree and in particular partitions of chains and antichains. As it is well-known there is a complete Ramsey theory for partitions of infinite subsets of  $\mathbb{N}$  as long as the colors are sufficiently definable (see [E]). On the other hand, the corresponding result for partitions of infinite dyadic subtrees of the Cantor tree fails in the sense that if we color all dyadic subtrees of the Cantor tree into finitely many, say, open colors, then we cannot expect to find a dyadic subtree all of whose dyadic subtrees are monochromatic. This has been recognized quite early by Galvin. His conjecture about partitions of  $k$ -tuples of reals, settled in the affirmative by Blass [B], reflects this phenomenon.

So, it is necessary, in order to have Ramsey theorems for trees, to consider not all subsets of the dyadic tree but only those which are of a fixed "shape". By now, there are several partition theorems along this line, obtained in [Ste] for chains, in [Mil] for strong subtrees, in [LSV] for strongly increasing sequences of reals, and in [Ka] for rapidly increasing subtrees.

It is well-known, and it is incorporated in the abstract Ramsey theory due to Carlson [C], that in order to obtain an infinite dimensional Ramsey result, one needs a pigeon-hole principle that corresponds to the finite dimensional case. In the case of partitions of infinite subsets of  $\mathbb{N}$ , this is the classical pigeon-hole principle. In the case of trees, this is the deep and fundamental Halpern–Läuchli partition theorem [HL]. The original proof was using metamathematical arguments. The proof avoiding metamathematics was given in [AFK].

For a presentation of some of the partition theorems we use, we refer the reader to [AT]. Applications of Ramsey theory for trees in analysis and topology can be found in [ADK] and [To1].

It is a standard fact that once one is willing to present results about trees one has to set up a, rather large, notational system. Below, we gather all the notations we need. We follow the conventions from [ADK] which are, more or less, standard.

**2.1. Notation.** Let  $\mathbb{N} = \{0, 1, 2, \dots\}$  denote the set of natural numbers. By  $[\mathbb{N}]^\infty$  we denote the set of all infinite subsets of  $\mathbb{N}$ , and for every  $L \in [\mathbb{N}]^\infty$  by  $[L]^\infty$  we denote the set of all infinite subsets of  $L$ .

**2.1.1.** By  $2^{<\mathbb{N}}$  we denote the set of all finite sequences of 0's and 1's (the empty sequence is included). We view  $2^{<\mathbb{N}}$  as a tree equipped with the (strict) partial order  $\sqsubset$  of extension. If  $t \in 2^{<\mathbb{N}}$ , then the length  $|t|$  of  $t$  is defined to be the cardinality

of the set  $\{s : s \sqsubset t\}$ . If  $s, t \in 2^{<\mathbb{N}}$ , then by  $s \hat{\ } t$  we denote their concatenation. Two nodes  $s, t$  are said to be incomparable if neither  $s \sqsubseteq t$  nor  $t \sqsubseteq s$ . A subset of  $2^{<\mathbb{N}}$  consisting of pairwise incomparable nodes is said to be an *antichain*, while a subset of  $2^{<\mathbb{N}}$  consisting of pairwise comparable nodes is called a *chain*. For every  $x \in 2^{\mathbb{N}}$  and every  $n \geq 1$  we set  $x|n = (x(0), \dots, x(n-1)) \in 2^{<\mathbb{N}}$  while  $x|0 = \emptyset$ . For  $x, y \in (2^{<\mathbb{N}} \cup 2^{\mathbb{N}})$  with  $x \neq y$  we denote by  $x \wedge y$  the  $\sqsubseteq$ -maximal node  $t$  of  $2^{<\mathbb{N}}$  with  $t \sqsubseteq x$  and  $t \sqsubseteq y$ . Moreover, we write  $x \prec y$  if  $w \hat{\ } 0 \sqsubseteq x$  and  $w \hat{\ } 1 \sqsubseteq y$ , where  $w = x \wedge y$ . We also write  $x \preceq y$  if either  $x = y$  or  $x \prec y$ . The ordering  $\prec$  restricted on  $2^{\mathbb{N}}$  is the usual lexicographical ordering of the Cantor set.

2.1.2. We view every subset of  $2^{<\mathbb{N}}$  as a *subtree* with the induced partial ordering. A subtree  $T$  of  $2^{<\mathbb{N}}$  is said to be *pruned* if for every  $t \in T$  there exists  $s \in T$  with  $t \sqsubset s$ . It is said to be *downwards closed* if for every  $t \in T$  and every  $s \sqsubset t$  we have that  $s \in T$ . For a subtree  $T$  of  $2^{<\mathbb{N}}$  (not necessarily downwards closed), by  $\hat{T}$  we denote the *downwards closure* of  $T$ , that is, the set  $\hat{T} := \{s : \exists t \in T \text{ with } s \sqsubseteq t\}$ . If  $T$  is downwards closed, then the *body*  $[T]$  of  $T$  is the set  $\{x \in 2^{\mathbb{N}} : x|n \in T \ \forall n\}$ .

2.1.3. Let  $T$  be a (not necessarily downwards closed) subtree of  $2^{<\mathbb{N}}$ . For every  $t \in T$  by  $|t|_T$  we denote the cardinality of the set  $\{s \in T : s \sqsubset t\}$  and for every  $n \in \mathbb{N}$  we set  $T(n) := \{t \in T : |t|_T = n\}$ . Moreover, for every  $t_1, t_2 \in T$  by  $t_1 \wedge_T t_2$  we denote the  $\sqsubseteq$ -maximal node  $w$  of  $T$  such that  $w \sqsubseteq t_1$  and  $w \sqsubseteq t_2$ . Notice that  $t_1 \wedge_T t_2 \sqsubseteq t_1 \wedge t_2$ . Given two subtrees  $S$  and  $T$  of  $2^{<\mathbb{N}}$ , we say that  $S$  is a *regular* subtree of  $T$  if  $S \subseteq T$  and for every  $n \in \mathbb{N}$  there exists  $m \in \mathbb{N}$  such that  $S(n) \subseteq T(m)$ . For a regular subtree  $T$  of  $2^{<\mathbb{N}}$ , the *level set*  $L_T$  of  $T$  is the set  $\{l_n : T(n) \subseteq 2^{l_n}\} \subseteq \mathbb{N}$ . Notice that the infinite chains of  $T$  are naturally identified with the product  $[\hat{T}] \times [L_T]^\infty$ . A pruned subtree  $T$  of  $2^{<\mathbb{N}}$  is said to be *dyadic* if every  $t \in T$  has exactly two immediate successors in  $T$ . We observe that a subtree  $T$  of the Cantor tree is a regular dyadic subtree of  $2^{<\mathbb{N}}$  if and only if there exists a (necessarily unique) bijection  $i_T : 2^{<\mathbb{N}} \rightarrow T$  such that the following are satisfied.

- (1) For every  $t_1, t_2 \in 2^{<\mathbb{N}}$  we have  $|t_1| = |t_2|$  if and only if  $|i_T(t_1)|_T = |i_T(t_2)|_T$ .
- (2) For every  $t_1, t_2 \in 2^{<\mathbb{N}}$  we have  $t_1 \sqsubset t_2$  (respectively,  $t_1 \prec t_2$ ) if and only if  $i_T(t_1) \sqsubset i_T(t_2)$  (respectively,  $i_T(t_1) \prec i_T(t_2)$ ).

When we write  $T = (s_t)_{t \in 2^{<\mathbb{N}}}$ , where  $T$  is a regular dyadic subtree of  $2^{<\mathbb{N}}$ , we mean that  $s_t = i_T(t)$  for all  $t \in 2^{<\mathbb{N}}$ . Finally we notice the following. If  $T$  is a regular dyadic subtree of  $2^{<\mathbb{N}}$  and  $R$  is a regular dyadic subtree of  $T$ , then  $R$  is a regular dyadic subtree of  $2^{<\mathbb{N}}$  too.

2.1.4. Let  $L$  be an infinite subset of  $2^{<\mathbb{N}}$  and  $\sigma \in 2^{\mathbb{N}}$ . We say that  $L$  *converges* to  $\sigma$  if for every  $k \in \mathbb{N}$  the set  $L \setminus \{t \in 2^{<\mathbb{N}} : \sigma|k \sqsubseteq t\}$  is finite. The element  $\sigma$  will be called the *limit* of the set  $L$ . We write  $L \rightarrow \sigma$  to denote that  $L$  converges to  $\sigma$ .

2.1.5. For every infinite  $L \subseteq 2^{<\mathbb{N}}$  and every  $\sigma \in 2^{\mathbb{N}}$  we write  $L \prec \sigma$  (respectively,  $\sigma \prec L$ ) to denote the fact that for every  $t \in L$  we have  $t \prec \sigma$  (respectively, for every  $t \in L$  we have  $\sigma \prec t$ ).

2.2. **Chains.** For a regular dyadic subtree  $T$  of  $2^{<\mathbb{N}}$ , denote by  $[T]_{\text{chains}}$  the set of all infinite chains of  $T$ . By identifying every infinite chain of  $T$  with its characteristic function (that is, an element of  $2^T$ ), it is easy to see that the set  $[T]_{\text{chains}}$  is a  $G_\delta$  (hence Polish) subspace of  $2^T$ . The following result, essentially due to Stern [Ste] (see also [Mi, Pa]), includes the Ramsey property of  $[T]_{\text{chains}}$  needed in the sequel.

**Theorem 5.** *Let  $T$  be a regular dyadic subtree of  $2^{<\mathbb{N}}$  and let  $A$  be an analytic subset of  $[T]_{\text{chains}}$ . Then there exists a regular dyadic subtree  $R$  of  $T$  such that either  $[R]_{\text{chains}} \subseteq A$ , or  $[R]_{\text{chains}} \cap A = \emptyset$ .*

2.3. **Increasing and decreasing antichains.** This subsection is devoted to the presentation of an analogue of Theorem 5 for infinite antichains of the Cantor tree. It not difficult to find an open partition of all infinite antichains  $(t_n)$  of  $2^{<\mathbb{N}}$  satisfying  $t_n \prec t_{n+1}$  and  $|t_n| < |t_{n+1}|$  for every  $n \in \mathbb{N}$  and such that there is no dyadic subtree of  $2^{<\mathbb{N}}$  for which all of its antichains of the above form are monochromatic. This explains the necessity of condition (2) in the following definition.

**Definition 6.** *Let  $T$  be a regular dyadic subtree of the Cantor tree  $2^{<\mathbb{N}}$ . An infinite antichain  $(t_n)$  of  $T$  will be called increasing if the following conditions are satisfied.*

- (1) *For every  $n, m \in \mathbb{N}$  with  $n < m$  we have  $|t_n|_T < |t_m|_T$ .*
- (2) *For every  $n, m, l \in \mathbb{N}$  with  $n < m < l$  we have  $|t_n|_T \leq |t_m \wedge_T t_l|_T$ .*
- (3I) *For every  $n, m \in \mathbb{N}$  with  $n < m$  have  $t_n \prec t_m$ .*

*The set of all increasing antichains of  $T$  will be denoted by  $\text{Incr}(T)$ . Similarly, an infinite antichain  $(t_n)$  of  $T$  will be called decreasing if (1) and (2) above are satisfied and (3I) is replaced by the following.*

- (3D) *For every  $n, m \in \mathbb{N}$  with  $n < m$  we have  $t_m \prec t_n$ .*

*The set of all decreasing antichains of  $T$  will be denoted by  $\text{Decr}(T)$ .*

Below we collect some basic properties of increasing and decreasing antichains.

**Proposition 7.** *The following hold.*

- (P1) *Let  $(t_n) \in \text{Incr}(T)$  and let  $L = \{l_0 < l_1 < \dots\}$  be an infinite subset of  $\mathbb{N}$ . Then  $(t_{l_n}) \in \text{Incr}(T)$ . Similarly, if  $(t_n) \in \text{Decr}(T)$ , then  $(t_{l_n}) \in \text{Decr}(T)$ .*
- (P2) *Let  $(t_n)$  be an infinite antichain of  $T$ . Then there exists  $L = (l_n) \in [\mathbb{N}]^\infty$  such that either  $(t_{l_n}) \in \text{Incr}(T)$  or  $(t_{l_n}) \in \text{Decr}(T)$ .*
- (P3) *We have  $\text{Incr}(T) = \text{Incr}(2^{<\mathbb{N}}) \cap 2^T$ , and similarly for the decreasing antichains.*
- (P4) *Let  $(t_n)$  be an increasing (respectively, decreasing) antichain of  $2^{<\mathbb{N}}$ . Then  $(t_n)$  converges to  $\sigma$ , where  $\sigma$  is the unique element of  $2^{\mathbb{N}}$  determined by the chain  $(c_n)$  with  $c_n = t_n \wedge t_{n+1}$ .*

- (P5) If  $L$  is an infinite subset of  $2^{<\mathbb{N}}$  and  $\sigma \in 2^{\mathbb{N}}$  are such that  $L \rightarrow \sigma$  and  $L \prec \sigma$  (respectively,  $\sigma \prec L$ ), then every infinite subset of  $L$  contains an increasing (respectively, decreasing) antichain converging to  $\sigma$ .
- (P6) Let  $A_1 = (t_n^1)$  and  $A_2 = (t_n^2)$  be two increasing (respectively, decreasing) antichains of  $2^{<\mathbb{N}}$  converging to the same  $\sigma \in 2^{\mathbb{N}}$ . Then there exists an increasing (respectively, decreasing) antichain  $(t_n)$  of  $2^{<\mathbb{N}}$  converging to  $\sigma$  such that  $t_{2n} \in A_1$  and  $t_{2n+1} \in A_2$  for all  $n \in \mathbb{N}$ .
- (P7) Let  $(\sigma_n)$  be a sequence in  $2^{\mathbb{N}}$  converging to  $\sigma \in 2^{\mathbb{N}}$ . For every  $n \in \mathbb{N}$  let  $N_n = (t_k^n)$  be a sequence in  $2^{<\mathbb{N}}$  converging to  $\sigma_n$ . If  $\sigma_n \prec \sigma$  (respectively,  $\sigma_n \succ \sigma$ ) for all  $n$ , then there exist an increasing (respectively, decreasing) antichain  $(t_m)$  and  $L = \{n_m : m \in \mathbb{N}\}$  such that  $(t_m)$  converges to  $\sigma$  and  $t_m \in N_{n_m}$  for all  $m \in \mathbb{N}$ .

Most of the above properties are easily verified. We refer the reader to [ADK] for more information.

By property (P4) of the above proposition, we see that for every regular dyadic subtree  $T$  of  $2^{<\mathbb{N}}$  and every increasing (respectively, decreasing) antichain  $(t_n)$  of  $T$  there exists a unique  $\sigma \in [\hat{T}]$  such that the sequence  $(t_n)$  converges to  $\sigma$ . We call this  $\sigma$  as the *limit point* of  $(t_n)$ .

Let  $T$  be a regular dyadic subtree of  $2^{<\mathbb{N}}$ . As in the case of chains and by identifying every increasing antichain of  $T$  with its characteristic function, we see that the set  $\text{Incr}(T)$  is a  $G_\delta$  subspace of  $2^T$ . Respectively, the set  $\text{Decr}(T)$  is also a  $G_\delta$  subspace of  $2^T$ . The Ramsey properties of increasing and decreasing antichains are included in the following theorem.

**Theorem 8.** *Let  $T$  be a regular dyadic subtree of  $2^{<\mathbb{N}}$  and let  $A$  be an analytic subset of  $\text{Incr}(T)$  (respectively, of  $\text{Decr}(T)$ ). Then there exists a regular dyadic subtree  $R$  of  $T$  such that either  $\text{Incr}(R) \subseteq A$ , or  $\text{Incr}(R) \cap A = \emptyset$  (respectively, either  $\text{Decr}(R) \subseteq A$ , or  $\text{Decr}(R) \cap A = \emptyset$ ).*

We will briefly comment on the proof, referring to [ADK] for a more detailed presentation. The method is to reduce the coloring of  $\text{Incr}(T)$  (respectively,  $\text{Decr}(T)$ ) to a coloring of a certain class of subtrees of the dyadic tree, for which it is known that it is Ramsey.

We argue for the case of increasing antichains as the case of decreasing antichains is similar. For every regular dyadic subtree  $T$  of  $2^{<\mathbb{N}}$  we define a class  $[T]_{\text{Incr}}$  of regular subtrees of  $T$  as follows. For notational convenience, let us assume that  $T = 2^{<\mathbb{N}}$ . Let  $\sigma \in 2^{\mathbb{N}}$  not eventually zero. We select a sequence  $(s_n)$  in  $2^{<\mathbb{N}}$  such that  $s_n \sqsubset s_n \hat{\ } 1 \sqsubseteq s_{n+1} \sqsubset \sigma$  for every  $n \in \mathbb{N}$  (this can be done since  $\sigma$  is not eventually zero). Next, we select a sequence  $(\sigma_n)$  in  $2^{\mathbb{N}}$  such that  $s_n \hat{\ } 0 \sqsubset \sigma_n$  for every  $n \in \mathbb{N}$ . Let  $L = \{l_0 < l_1 < \dots\} \in [\mathbb{N}]^\infty$  where  $l_n = |s_n|$  for all  $n \in \mathbb{N}$ . A tree  $S$  belongs to  $[2^{<\mathbb{N}}]_{\text{Incr}}$  if there exist  $\sigma \in 2^{\mathbb{N}}$ , a sequence  $(s_n)$  in  $2^{<\mathbb{N}}$  and a sequence

$(\sigma_n)$  in  $2^{\mathbb{N}}$  as described above, such that

$$S = \bigcup_{k \in \mathbb{N}} \{\sigma_n |_{l_k} : n \leq k\}.$$

It is easy to see that  $S$  is a regular subtree and  $[\hat{S}] = \{\sigma_n : n \in \mathbb{N}\} \cup \{\sigma\}$ . Moreover, observe that the sequence  $I_S = (\sigma_n |_{l_{n+1}})$  is an increasing antichain of  $2^{<\mathbb{N}}$  which converges to  $\sigma$ . The map  $\Phi: [T]_{\text{Incr}} \rightarrow \text{Incr}(T)$  defined by  $\Phi(S) = I_S$  is easily seen to be continuous and onto. By the results in [Ka], the family  $[T]_{\text{Incr}}$  is Ramsey, that is, for every analytic subset  $B$  of  $[T]_{\text{Incr}}$  there exists a regular dyadic subtree  $R$  of  $T$  such that either  $[R]_{\text{Incr}} \subseteq B$ , or  $[R]_{\text{Incr}} \cap B = \emptyset$ .

Now let  $T$  and  $A$  be as in Theorem 8 and consider the coloring  $B = \Phi^{-1}(A)$  of  $[T]_{\text{Incr}}$ . If  $R$  is any regular dyadic subtree of  $T$  such that  $[R]_{\text{Incr}}$  is monochromatic with respect to  $B$ , then it is easy to see that so is  $\text{Incr}(R)$  with respect to  $A$ .

We notice that Theorem 8 has been obtained independently by Todorćević with a different proof based on Milliken's theorem ([To2]).

**2.4. Partitions of perfect sets of reals.** Recall that a subset  $M$  of a Polish space  $X$  is said to be meager (or of first category) if  $M$  is covered by a countable union of closed nowhere dense sets. A subset  $C$  of  $X$  is said to be co-meager if its complement is meager. Finally, a subset  $A$  of  $X$  is said to have the Baire property if there exist an open subset  $U$  of  $X$  and meager set  $M$  such that  $A \triangle U = M$ . It is classical fact that the family of all sets with the Baire property contains the  $\sigma$ -algebra generated by the analytic sets (see [Ke]). We will need the following partition theorem due to Galvin (see, e.g., [Ke, Theorem 19.6]).

**Theorem 9.** *Let  $X_1, \dots, X_n$  be perfect Polish spaces. Also let  $A$  be a subset of  $X_1 \times \dots \times X_n$  with the Baire property. If  $A$  is non-meager, then for every  $i \in \{1, \dots, n\}$  there exists a perfect set  $P_i \subseteq X_i$  such that  $P_1 \times \dots \times P_n \subseteq A$ .*

### 3. AN EXTENSION OF STERN'S THEOREM

Let us start with the proof of Theorem 3 stated in the introduction, which is implicitly contained in [ADK].

*Proof of Theorem 3.* Denote by  $(e_n)$  the standard basis of  $\ell_1$ . First, we argue as in [Ste] to homogenize the behavior of all subsequences of  $\{x_t : t \in 2^{<\mathbb{N}}\}$  determined by chains. In particular, consider the following subsets of  $[2^{<\mathbb{N}}]_{\text{chains}}$  defined by

$$\mathcal{X}_1 := \{(t_n) \in [2^{<\mathbb{N}}]_{\text{chains}} : (x_{t_n}) \text{ is equivalent to } (e_n)\},$$

$$\mathcal{X}_2 := \{(t_n) \in [2^{<\mathbb{N}}]_{\text{chains}} : (x_{t_n}) \text{ is weak Cauchy}\}, \text{ and}$$

$$\mathcal{X}_3 := [2^{<\mathbb{N}}]_{\text{chains}} \setminus (\mathcal{X}_1 \cup \mathcal{X}_2).$$

It is easy to see that the set  $\mathcal{X}_1$  is  $F_\sigma$ . On the other hand, the set  $\mathcal{X}_2$  is co-analytic (see [Ste] for a detailed explanation of this fact). Applying Theorem 5 successively three times, we obtain a regular dyadic subtree  $T_1$  of  $2^{<\mathbb{N}}$  such that



for every  $i \in \{1, 2, 3\}$  we have that either  $[T_1]_{\text{chains}} \subseteq \mathcal{X}_i$  or  $[T_1]_{\text{chains}} \cap \mathcal{X}_i = \emptyset$ . By Rosenthal's  $\ell_1$  theorem [Ro], we see that for every regular dyadic subtree  $R$  of  $2^{<\mathbb{N}}$  we have that either  $[R]_{\text{chains}} \cap \mathcal{X}_1 \neq \emptyset$ , or  $[R]_{\text{chains}} \cap \mathcal{X}_2 \neq \emptyset$ . It follows that there exists  $i \in \{1, 2\}$  such that  $[T]_{\text{chains}} \subseteq \mathcal{X}_i$ , that is, either for every infinite chain  $(t_n)$  of  $T_1$  the sequence  $(x_{t_n})$  is equivalent to the standard basis of  $\ell_1$ , or for every infinite chain  $(t_n)$  of  $T_1$  the sequence  $(x_{t_n})$  is weak Cauchy.

Now consider the following subsets of  $\text{Incr}(T_1)$  defined by

$$\begin{aligned} \mathcal{C}_1 &:= \{(t_n) \in \text{Incr}(T_1) : (x_{t_n}) \text{ is equivalent to } (e_n)\}, \\ \mathcal{C}_2 &:= \{(t_n) \in \text{Incr}(T_1) : (x_{t_n}) \text{ is weak Cauchy}\}, \text{ and} \\ \mathcal{C}_3 &:= \text{Incr}(T_1) \setminus (\mathcal{C}_1 \cup \mathcal{C}_2). \end{aligned}$$

Again we see that  $\mathcal{C}_1$  is  $F_\sigma$  while the set  $\mathcal{C}_2$  is co-analytic (this can be checked by similar arguments as in [Ste]). Applying Theorem 8 three times and arguing as before, we obtain a regular dyadic subtree  $T_2$  of  $T_1$  and  $j \in \{1, 2\}$  such that  $\text{Incr}(T_2) \subseteq \mathcal{C}_j$ .

Finally, applying Theorem 8 for the decreasing antichains of  $T_2$  and the colors

$$\begin{aligned} \mathcal{K}_1 &:= \{(t_n) \in \text{Decr}(T_2) : (x_{t_n}) \text{ is equivalent to } (e_n)\}, \\ \mathcal{K}_2 &:= \{(t_n) \in \text{Decr}(T_2) : (x_{t_n}) \text{ is weak Cauchy}\}, \text{ and} \\ \mathcal{K}_3 &:= \text{Decr}(T_2) \setminus (\mathcal{K}_1 \cup \mathcal{K}_2) \end{aligned}$$

we find a regular dyadic subtree  $T_3$  of  $T_2$  and  $l \in \{1, 2\}$  such that  $\text{Decr}(T_3) \subseteq \mathcal{K}_l$ .

If  $[T_3]_{\text{chains}}$ ,  $\text{Incr}(T_3)$  and  $\text{Decr}(T_3)$  avoid the colors  $\mathcal{X}_1$ ,  $\mathcal{C}_1$  and  $\mathcal{K}_1$  respectively, then the tree  $T_3$  is the desired one. If not, then we will pass to a further dyadic subtree  $T$  of  $T_3$  in order to achieve uniformity. So, assume that  $\text{Incr}(T_3)$  is included in  $\mathcal{C}_1$  (the other cases are similar). For every  $k \in \mathbb{N}$  set

$$\mathcal{F}_k := \{(t_n) \in \text{Incr}(T_3) : (x_{t_n}) \text{ is } k\text{-equivalent to } (e_n)\}.$$

Clearly  $\mathcal{F}_k$  is a closed subset  $\text{Incr}(T_3)$ . Moreover,  $\text{Incr}(T_3) = \bigcup_k \mathcal{F}_k$ . It follows that there exists  $k_0 \in \mathbb{N}$  such that the set  $\mathcal{F}_{k_0}$  has nonempty interior in  $\text{Incr}(T_3)$ . Let  $(t_n) \in \text{Int}(\mathcal{F}_{k_0})$ . There exists  $n_0 \in \mathbb{N}$  such that if  $(s_n) \in \text{Incr}(T_3)$  and  $s_n = t_n$  for every  $n \leq n_0$ , then  $(s_n) \in \mathcal{F}_{k_0}$ . We set  $w := t_{n_0+1} \wedge_T t_{n_0+2}$  and we define  $T := \{t \in T_3 : w \sqsubseteq t\}$ . Clearly  $T$  is a regular dyadic subtree of  $T_3$ . Moreover, it is easy to see that  $\text{Incr}(T) \subseteq \mathcal{F}_{k_0}$ . That is, for every increasing antichain  $(r_n)$  of  $T$  the sequence  $(x_{r_n})$  is  $k_0$ -equivalent to  $(e_n)$ . Thus, we have achieved the desired uniformity.

Finally, we notice that if  $\text{Incr}(T) \subseteq \mathcal{C}_2$ , then for every  $(t_n)$  and  $(s_n)$  in  $\text{Incr}(T)$  with the same limit point in  $[\hat{T}]$ , the sequences  $(x_{t_n})$  and  $(x_{s_n})$  are both weak\* convergent to the same element of  $X^{**}$ . For if not, then by property (P6) of Proposition 7, we would be able to construct an increasing antichain  $(r_n)$  of  $T$  such that the sequence  $(x_{r_n})$  is not weak Cauchy, contradicting in particular the fact

that  $\text{Incr}(T) \subseteq \mathcal{C}_2$ . The case of decreasing antichains is similarly treated. The proof is completed.  $\square$

**Remark 1.** We notice that the behavior of the sequence  $(x_t)_{t \in T}$  along chains of  $T$  is independent of the corresponding one along increasing antichains (and decreasing antichains, respectively). In particular, all subsequences of  $(x_t)_{t \in T}$  determined by chains and increasing antichains can be weak\* convergent while all subsequences determined by decreasing antichains are equivalent to the standard basis of  $\ell_1$ . For example, let  $X$  be the completion of  $c_{00}(2^{<\mathbb{N}})$  under the norm

$$\|x\| := \sup \left\{ \sum_{n \in \mathbb{N}} |x(t_n)| : (t_n) \in \text{Decr}(2^{<\mathbb{N}}) \right\}.$$

Consider the standard Hamel basis  $(e_t)_{t \in 2^{<\mathbb{N}}}$  of  $c_{00}(2^{<\mathbb{N}})$ . It is easy to see that for every sequence  $(t_n)$  in  $2^{<\mathbb{N}}$  which is either a chain or an increasing antichain, the sequence  $(e_{t_n})$  is 1-equivalent to the standard basis of  $c_0$ . In particular, it is weakly-null. On the other hand, if  $(t_n)$  is a decreasing antichain, then the sequence  $(e_{t_n})$  is 1-equivalent to the standard basis of  $\ell_1$ .

We will also need the following result which is based on Theorem 3 and on the properties of increasing and decreasing antichains described in Proposition 7.

**Theorem 10.** *Let  $X$  be a separable Banach space not containing  $\ell_1$ . Also let  $\Delta = \{x_t : t \in 2^{<\mathbb{N}}\}$  be a bounded family in  $X$ . Then there exist a regular dyadic subtree  $T$  of  $2^{<\mathbb{N}}$  and a family  $\{y_\sigma^0, y_\sigma^+, y_\sigma^- : \sigma \in P\} \subseteq X^{**}$ , where  $P = [\hat{T}]$ , such that for every  $\sigma \in P$  the following are satisfied.*

- (1) *The sequence  $(x_{\sigma|n})_{n \in L_T}$  is weak\* convergent to  $y_\sigma^0$  (recall that  $L_T$  stands for the level set of  $T$ ).*
- (2) *For every sequence  $(\sigma_n)$  in  $P$  converging to  $\sigma$  such that  $\sigma_n \prec \sigma$  for all  $n \in \mathbb{N}$ , the sequence  $(y_{\sigma_n}^{\varepsilon_n})$  is weak\* convergent to  $y_\sigma^+$  for any choice of  $\varepsilon_n \in \{0, +, -\}$ . If such a sequence  $(\sigma_n)$  does not exist, then  $y_\sigma^+ = y_\sigma^0$ .*
- (3) *For every sequence  $(\sigma_n)$  in  $P$  converging to  $\sigma$  such that  $\sigma \prec \sigma_n$  for all  $n \in \mathbb{N}$ , the sequence  $(y_{\sigma_n}^{\varepsilon_n})$  is weak\* convergent to  $y_\sigma^-$  for any choice of  $\varepsilon_n \in \{0, +, -\}$ . If such a sequence  $(\sigma_n)$  does not exist, then  $y_\sigma^- = y_\sigma^0$ .*
- (4) *For every infinite subset  $L$  of  $T$  converging to  $\sigma$  with  $L \prec \sigma$ , the sequence  $(x_t)_{t \in L}$  is weak\* convergent to  $y_\sigma^+$ .*
- (5) *For every infinite subset  $L$  of  $T$  converging to  $\sigma$  with  $\sigma \prec L$ , the sequence  $(x_t)_{t \in L}$  is weak\* convergent to  $y_\sigma^-$ .*

Moreover, the functions  $0, +, - : P \times (B_{X^*}, w^*) \rightarrow \mathbb{R}$  defined by

$$0(\sigma, x^*) = y_\sigma^0(x^*), \quad +(\sigma, x^*) = y_\sigma^+(x^*), \quad -(\sigma, x^*) = y_\sigma^-(x^*)$$

are all Borel.

The family  $\{y_\sigma^0, y_\sigma^+, y_\sigma^- : \sigma \in P\}$  obtained by Theorem 10, determines the weak\* closure of the family  $\{x_t : t \in T\}$ . Theorem 10 appears in [ADK] where it is stated

and proved in the broader frame of separable Rosenthal compacta. It is part of a finer analysis of the topological behavior of the family  $\{x_t : t \in 2^{<\mathbb{N}}\}$  yielding a complete canonization of any family as above.

*Proof.* Applying Theorem 3 and invoking our hypotheses on the space  $X$ , we obtain a regular dyadic subtree  $T$  of  $2^{<\mathbb{N}}$  such that, setting  $P = [\hat{T}]$ , the following are satisfied.

- (i) For every  $(t_n) \in \text{Incr}(T)$  the sequence  $(x_{t_n})$  is weak Cauchy.
- (ii) For every  $(t_n) \in \text{Decr}(T)$  the sequence  $(x_{t_n})$  is weak Cauchy.
- (iii) For every  $\sigma \in P$  the sequence  $(x_{\sigma|n})_{n \in L_T}$  is weak Cauchy.

For every  $\sigma \in P$  we define  $y_\sigma^0, y_\sigma^+$  and  $y_\sigma^-$  in  $X^{**}$  as follows. First, let  $y_\sigma^0$  be the weak\* limit of the sequence  $(x_{\sigma|n})_{n \in L_T}$ . If there exists an increasing antichain  $(t_n)$  of  $T$  converging to  $\sigma$ , then let  $y_\sigma^+$  be the weak\* limit of the sequence  $(x_{t_n})$ . By Theorem 3,  $y_\sigma^+$  is well-defined and independent of the choice of  $(t_n)$ . Otherwise, we set  $y_\sigma^+ = y_\sigma^0$ . Similarly, we define  $y_\sigma^-$  as the weak\* limit of the sequence  $(x_{t_n})$  with  $(t_n)$  a decreasing antichain of  $T$  converging to  $\sigma$ , if such an antichain exists. Otherwise, we set  $y_\sigma^- = y_\sigma^0$ .

We claim that the tree  $T$  and the family  $\{y_\sigma^0, y_\sigma^+, y_\sigma^- : \sigma \in P\}$  are as desired. First we notice that, by property (P5) in Proposition 7, properties (i) and (ii) above are strengthened as follows.

- (iv) For every  $\sigma \in P$  and every infinite subset  $L$  of  $T$  with  $L \rightarrow \sigma$  and  $L \prec \sigma$ , the sequence  $(x_t)_{t \in L}$  is weak\* convergent to  $y_\sigma^+$ .
- (v) For every  $\sigma \in P$  and every infinite subset  $L$  of  $T$  with  $L \rightarrow \sigma$  and  $\sigma \prec L$ , the sequence  $(x_t)_{t \in L}$  is weak\* convergent to  $y_\sigma^-$ .

Hence, by (iii), (iv) and (v), we see that properties (1), (4) and (5) in the statement of the theorem are satisfied. We will only check that property (2) is satisfied (the argument for (3) is symmetric). We argue by contradiction. So assume that there exist a sequence  $(\sigma_n)$  in  $P$ ,  $\sigma \in P$  and a sequence  $(\varepsilon_n)$  in  $\{0, +, -\}$  such that  $\sigma_n \prec \sigma$  for all  $n \in \mathbb{N}$ ,  $\sigma_n \rightarrow \sigma$  while  $(y_{\sigma_n}^{\varepsilon_n})$  is not weak\* convergent to  $y_\sigma^+$ . Hence, there exist  $L \in [\mathbb{N}]^\infty$ , a weak\* open neighborhood  $V$  of  $y_\sigma^+$  such that  $y_{\sigma_n}^{\varepsilon_n} \notin \bar{V}^{w*}$  for every  $n \in L$ . For every  $n \in L$  we select a sequence  $(t_k^n)$  in  $T$  such that the following are satisfied.

- (a) The sequence  $N_n = (t_k^n)$  converges (as a subset of  $T$ ) to  $\sigma_n$ .
- (b) The sequence  $(x_{t_k^n})$  is weak\* convergent to  $y_{\sigma_n}^{\varepsilon_n}$ .
- (c) For every  $k \in \mathbb{N}$  we have  $x_{t_k^n} \notin \bar{V}^{w*}$ .

By property (P7) in Proposition 7, there exists a diagonal increasing antichain  $(t_m)$  converging to  $\sigma$ . By (c) above, we see that  $(x_{t_m})$  is not weak\* convergent to  $y_\sigma^+$ , which is a contradiction by the definition of  $y_\sigma^+$ .

Finally, we will check the Borelness of the maps  $0, +$  and  $-$ . Let  $\{l_0 < l_1 < \dots\}$  be the increasing enumeration of the level set  $L_T$  of  $T$ . For every  $n \in \mathbb{N}$  define  $h_n : P \times (B_{X^*}, w^*) \rightarrow \mathbb{R}$  by  $h_n(\sigma, x^*) = x^*(x_{\sigma|l_n})$ . Clearly  $h_n$  is continuous. Notice

that for every  $(\sigma, x^*) \in P \times B_{X^*}$  we have

$$0(\sigma, x^*) = y_\sigma^0(x^*) = \lim h_n(\sigma, x^*).$$

Hence 0 is Borel (actually, it is Baire class one). We will only check the Borelness of the function  $+$  (the argument for the map  $-$  is symmetric). For every  $n \in \mathbb{N}$  and every  $\sigma \in P$  let  $l_n(\sigma)$  be the lexicographically minimum of the closed set  $\{\tau \in P : \sigma|_{l_n} \sqsubset \tau\}$ . Clearly  $l_n(\sigma) \in P$ . Moreover, observe that the function  $P \ni \sigma \mapsto l_n(\sigma) \in P$  is continuous. Invoking the definition of  $y_\sigma^+$  and property (2) in the statement of the theorem, we see that for all  $(\sigma, x^*) \in P \times B_{X^*}$  we have

$$+(\sigma, x^*) = y_\sigma^+(x^*) = \lim y_{l_n(\sigma)}^0(x^*) = \lim 0(l_n(\sigma), x^*).$$

Thus,  $+$  is a Borel map and the proof is completed.  $\square$

#### 4. PERFECT UNCONDITIONAL FAMILIES

This section is devoted to the proof of Theorem 4 stated in the introduction. Let us recall that a family  $\{x_i : i \in I\}$  in a Banach space  $X$  is said to be *1-unconditional* if for every  $F \subseteq G \subseteq I$  and every  $(a_i)_{i \in G}$  in  $\mathbb{R}^G$  we have

$$\left\| \sum_{i \in F} a_i x_i \right\| \leq \left\| \sum_{i \in G} a_i x_i \right\|.$$

As we have already mentioned, the construction of the perfect subset  $R$  in Theorem 4 is done by induction. The basic step for accomplishing the construction is described in the following lemma. Its proof is based on the partition theorem of Galvin (Theorem 9 above).

**Lemma 11.** *Let  $X, Q$  and  $\mathcal{D}$  be as in Theorem 4. Let  $n \in \mathbb{N}$  and let  $Q_0, \dots, Q_n$  be pairwise disjoint perfect subsets of  $Q$ . Then for every  $i \in \{0, \dots, n\}$  there exists a perfect subset  $R_i$  of  $Q_i$  such that for every  $(\sigma_0, \dots, \sigma_n) \in R_0 \times \dots \times R_n$  the family  $\{z_{\sigma_0}^{**}, \dots, z_{\sigma_n}^{**}\}$  is 1-unconditional.*

*Proof.* For every  $k \in \mathbb{N}$  and every tuple  $P_0, \dots, P_k$  of pairwise disjoint perfect subsets of  $Q$  we set

$$U(P_0, \dots, P_k) := \{(\sigma_0, \dots, \sigma_k) \in P_0 \times \dots \times P_k : \{z_{\sigma_0}^{**}, \dots, z_{\sigma_k}^{**}\} \text{ is 1-unconditional}\}.$$

Let  $n \in \mathbb{N}$  and let  $Q_0, \dots, Q_n$  be as in the statement of the lemma. For every nonempty  $F \subseteq \{0, \dots, n\}$ , every rational  $\varepsilon > 0$  and every  $(a_i)_{i=0}^n \in \mathbb{Q}^{n+1}$  we define  $D := D(F, \varepsilon, (a_i)_{i=0}^n)$  by

$$D = \left\{ (\sigma_0, \dots, \sigma_n) \in Q_0 \times \dots \times Q_n : \left\| \sum_{i \in F} a_i z_{\sigma_i}^{**} \right\| < (1 + \varepsilon) \left\| \sum_{i=0}^n a_i z_{\sigma_i}^{**} \right\| \right\}.$$

Clearly we have

$$(1) \quad U(Q_0, \dots, Q_n) = \bigcap_{F, \varepsilon, (a_i)_{i=0}^n} D(F, \varepsilon, (a_i)_{i=0}^n).$$

**Claim 1.** *The set  $D = D(F, \varepsilon, (a_i)_{i=0}^n)$  has the Baire property in  $Q_0 \times \dots \times Q_n$ .*

*Proof of the claim.* By our assumptions, we see that the functions  $\Phi_n, \Phi_F: Q_0 \times \cdots \times Q_n \times (B_{X^*}, w^*) \rightarrow \mathbb{R}$ , defined by  $\Phi_n(\sigma_0, \dots, \sigma_n, x^*) = \sum_{i=0}^n a_i z_{\sigma_i}^{**}(x^*)$  and  $\Phi_F(\sigma_0, \dots, \sigma_n, x^*) = \sum_{i \in F} a_i z_{\sigma_i}^{**}(x^*)$  respectively, are both Borel. Notice that

$$\begin{aligned} (\sigma_0, \dots, \sigma_n) \in D &\Leftrightarrow \exists p \in \mathbb{Q} \left( \left\| \sum_{i \in F} a_i z_{\sigma_i}^{**} \right\| \leq p \text{ and } \frac{p}{1+\varepsilon} < \left\| \sum_{i=0}^n a_i z_{\sigma_i}^{**} \right\| \right) \\ &\Leftrightarrow \exists p \in \mathbb{Q} \left[ (\forall x^* \in B_{X^*} \text{ we have } \Phi_F(\sigma_0, \dots, \sigma_n, x^*) \leq p) \right. \\ &\quad \left. \text{and } (\exists x^* \in B_{X^*} \text{ with } \frac{p}{1+\varepsilon} < \Phi_n(\sigma_0, \dots, \sigma_n, x^*)) \right] \end{aligned}$$

Hence,  $D$  belongs to the  $\sigma$ -algebra generated by the analytic sets. Finally, we recall that the  $\sigma$ -algebra generated by the analytic sets is included in the  $\sigma$ -algebra of all sets with Baire property. The claim is proved.  $\square$

**Claim 2.** For every tuple  $P_0, \dots, P_n$  of perfect subsets of  $Q_0, \dots, Q_n$  there exists  $(\sigma_0, \dots, \sigma_n) \in D(F, \varepsilon, (a_i)_{i=0}^n) \cap (P_0 \times \cdots \times P_n)$ .

*Proof of the claim.* For every  $i \in \{0, \dots, n\}$  we fix  $\tau_i \in P_i$ . Let  $x_0^* \in B_{X^*}$  such that

$$\left\| \sum_{i \in F} a_i z_{\tau_i}^{**} \right\| < (1+\varepsilon) \sum_{i \in F} a_i z_{\tau_i}^{**}(x_0^*).$$

The family  $\{z_{\sigma}^{**} : \sigma \in Q\}$  accumulates to 0 in the weak\* topology. Therefore,  $z_{\sigma}^{**}(x_0^*) = 0$  for all but countable many  $\sigma \in Q$ . For every  $i \in \{0, \dots, n\} \setminus F$  we may select  $\sigma'_i \in P_i$  with  $z_{\sigma'_i}^{**}(x_0^*) = 0$ . Finally, for every  $i \in \{0, \dots, n\}$  we define  $\sigma_i := \tau_i$  if  $i \in F$  and  $\sigma_i := \sigma'_i$  otherwise. Then  $(\sigma_0, \dots, \sigma_n) \in P_0 \times \cdots \times P_n$  and, moreover,

$$\begin{aligned} \left\| \sum_{i \in F} a_i z_{\sigma_i}^{**} \right\| &= \left\| \sum_{i \in F} a_i z_{\tau_i}^{**} \right\| < (1+\varepsilon) \sum_{i \in F} a_i z_{\tau_i}^{**}(x_0^*) \\ &= (1+\varepsilon) \sum_{i=0}^n a_i z_{\sigma_i}^{**}(x_0^*) \leq (1+\varepsilon) \left\| \sum_{i=0}^n a_i z_{\sigma_i}^{**} \right\|. \end{aligned}$$

Thus,  $(\sigma_0, \dots, \sigma_n) \in D(F, \varepsilon, (a_i)_{i=0}^n) \cap (P_0 \times \cdots \times P_n)$  and the claim is proved.  $\square$

By Claim 1, for every  $F, \varepsilon$  and  $(a_i)_{i=0}^n$  the set  $D(F, \varepsilon, (a_i)_{i=0}^n)$  has the Baire property in  $Q_0 \times \cdots \times Q_n$ . We claim that the set  $D(F, \varepsilon, (a_i)_{i=0}^n)$  must be co-meager in  $Q_0 \times \cdots \times Q_n$ . Indeed, if not, then by Theorem 9 there would exist perfect subsets  $P_0, \dots, P_n$  of  $Q_0, \dots, Q_n$  such that  $D(F, \varepsilon, (a_i)_{i=0}^n) \cap (P_0 \times \cdots \times P_n) = \emptyset$  which clearly contradicts Claim 2. It follows that  $D(F, \varepsilon, (a_i)_{i=0}^n)$  is co-meager. By (1), so is the set  $U(Q_0, \dots, Q_n)$ . By Theorem 9 once more, there exist perfect subsets  $R_0, \dots, R_n$  of  $Q_0, \dots, Q_n$  such that  $R_0 \times \cdots \times R_n \subseteq U(Q_0, \dots, Q_n)$  and the proof is completed.  $\square$

We are ready to proceed to the proof of Theorem 4.

*Proof of Theorem 4.* By recursion on the length of finite sequences in  $2^{<\mathbb{N}}$ , we shall construct a family  $(R_t)_{t \in 2^{<\mathbb{N}}}$  of perfect subsets of  $Q$  such that the following are satisfied.

- (C1) For every  $t \in 2^{<\mathbb{N}}$  we have  $\text{diam}(R_t) \leq \frac{1}{2^{|t|}}$ .
- (C2) For every  $t \in 2^{<\mathbb{N}}$  we have  $R_{t \smallfrown 0}, R_{t \smallfrown 1} \subseteq R_t$  and  $R_{t \smallfrown 0} \cap R_{t \smallfrown 1} = \emptyset$ .
- (C3) For every  $n \geq 1$ , every  $t \in 2^n$  and every  $\sigma_t \in R_t$  the family  $\{z_{\sigma_t}^{**} : t \in 2^n\}$  is 1-unconditional.

Assuming that the construction has been carried out, we set

$$R := \bigcup_{\sigma \in 2^{\mathbb{N}}} \bigcap_{n \in \mathbb{N}} R_{\sigma|n}.$$

Clearly,  $R$  is a perfect subset of  $Q$ . Moreover, using condition (C2) above, it is easy to see that the family  $\{z_{\sigma}^{**} : \sigma \in R\}$  is 1-unconditional.

We proceed to the construction. We set  $R_{\emptyset} := Q$ . Assume that for some  $n \geq 1$  the family  $(R_t)_{t \in 2^{n-1}}$  has been constructed. For every  $t \in 2^{n-1}$  and every  $i \in \{0, 1\}$  we select  $Q_{t \smallfrown i}$  perfect subset of  $R_t$  with  $\text{diam}(Q_{t \smallfrown i}) \leq \frac{1}{2^n}$  and such that  $Q_{t \smallfrown 0} \cap Q_{t \smallfrown 1} = \emptyset$ . Let  $t_0 \prec \dots \prec t_{2^n-1}$  denote the  $\prec$ -increasing enumeration of  $2^n$ . We apply Lemma 11 to the family of perfect sets  $Q_{t_0}, \dots, Q_{t_{2^n-1}}$  and for every  $t \in 2^n$  we obtain a perfect subset  $R_t$  of  $Q_t$  such that for every  $(\sigma_t)_{t \in 2^n} \in \prod_{t \in 2^n} R_t$  the family  $\{z_{\sigma_t}^{**} : t \in 2^n\}$  is 1-unconditional. Clearly, the family  $(R_t)_{t \in 2^n}$  satisfies (C1)–(C3) above. This completes the construction and the proof is completed.  $\square$

**Remark 2.** We notice that the existence of a subset of  $X^{**}$  of the size of the continuum which is weak\* discrete and having 0 as the unique weak\* accumulation point can be obtained by the results of Todorćević in [To1], after observing that  $(B_{X^{**}}, w^*)$  is a separable Rosenthal compact containing 0 as a non- $G_\delta$  point. His remarkable proof uses, among others, forcing arguments and absoluteness. This result has been strengthened and extended to a wider class of Rosenthal compacta in [ADK], with a proof avoiding metamathematics.

## 5. THE MAIN RESULTS

In this section we present the proof of Theorem 1 stated in the introduction. We also state and prove some of its consequences. As we have mentioned, the proof is based on the following fundamental construction due to Stegall [St]. A variation of Stegall's construction has been presented by Godefroy and Talagrand [GT] in the more general context of representable Banach spaces (see, also, [GL]). We refer the reader to [AGR] for a full account of related results.

**Theorem 12.** *Let  $X$  be a separable Banach space with non-separable dual. Then for every  $\varepsilon > 0$  there exist a family  $\Delta_\varepsilon = \{x_t : t \in 2^{<\mathbb{N}}\}$  in  $(1 + \varepsilon)B_X$  and a subset  $D_\varepsilon = \{x_\sigma^* : \sigma \in 2^{\mathbb{N}}\}$  in the sphere of  $X^*$  which is weak\* homeomorphic to the Cantor set  $2^{\mathbb{N}}$  via the map  $\sigma \mapsto x_\sigma^*$  and such that for every  $\sigma \in 2^{\mathbb{N}}$  and every  $t \in 2^{<\mathbb{N}}$  we have*

$$|x_\sigma^*(x_t) - \delta_{\sigma t}| < \frac{1}{2^{|t|}}$$

where  $\delta_{\sigma t} = 1$  if  $t \sqsubset \sigma$  and  $\delta_{\sigma t} = 0$  otherwise.

Although the above statement is not explicitly isolated in [St], it is the precise content of the proof.

We notice the following property of the sets  $\Delta_\varepsilon$  and  $D_\varepsilon$  obtained by Theorem 12. For every  $\sigma \in 2^{\mathbb{N}}$  let  $x_\sigma^{**}$  be any weak\* accumulation point of the family  $\{x_{\sigma|n} : n \in \mathbb{N}\}$ . Then the family  $\{(x_\sigma^*, x_\sigma^{**}) : \sigma \in 2^{\mathbb{N}}\} \subseteq X^* \times X^{**}$  forms a bi-orthogonal system, and so, the set  $\{x_\sigma^{**} : \sigma \in 2^{\mathbb{N}}\}$  is weak\* discrete.

We are ready to proceed to the proof of Theorem 1.

*Proof of Theorem 1.* We apply Theorem 12 for  $\varepsilon = 1$  and we obtain a family  $\Delta_1 = \{x_t : t \in 2^{<\mathbb{N}}\}$  in  $2B_X$  and a family  $D_1 = \{x_\sigma^* : \sigma \in 2^{\mathbb{N}}\}$  in the sphere of  $X^*$  as described in Theorem 12. Next, we apply Theorem 10 for the family  $\Delta = \{x_t/2 : t \in 2^{<\mathbb{N}}\}$  and we obtain a regular dyadic subtree  $T$  of  $2^{<\mathbb{N}}$  and a family  $\{y_\sigma^0, y_\sigma^+, y_\sigma^- : \sigma \in P\} \subseteq B_{X^{**}}$  where  $P = [\hat{T}]$ . Notice that the set  $\{(y_\sigma^0, 2x_\sigma^*) : \sigma \in P\}$  forms a bi-orthogonal system. We fix a perfect subset  $Q$  of  $P$  with the following property. For every  $\tau \in Q$  there exists a sequence  $(\tau_n)$  in  $P$  with  $\tau_n \prec \tau$  for all  $n \in \mathbb{N}$  and such that  $\tau_n \rightarrow \tau$ . This condition guarantees that the function  $y_\tau^+$  is not trivially equal to  $y_\tau^0$ . For every  $\tau \in Q$  we set  $z_\tau^{**} = y_\tau^0 - y_\tau^+$  and  $z_\tau^* = 2x_\tau^*$ .

**Claim.** *The following hold.*

- (1) *For every  $\tau \in Q$  we have  $z_\tau^{**} \neq 0$ .*
- (2) *The family  $\{(z_\tau^*, z_\tau^{**}) : \tau \in Q\}$  forms a bounded bi-orthogonal system in  $X^* \times X^{**}$ .*
- (3) *The family  $\{z_\tau^{**} : \tau \in Q\}$  is weak\* discrete having 0 as the unique weak\* accumulation point.*
- (4) *The function  $\Phi : Q \times (B_{X^*}, w^*) \rightarrow \mathbb{R}$  defined by  $\Phi(\tau, x^*) = z_\tau^{**}(x^*)$  is Borel.*

Granting the claim, we complete the proof as follows. By (3) and (4) above, we see that Theorem 4 can be applied to the family  $\mathcal{D} = \{z_\tau^{**} : \tau \in Q\}$ . Hence, there exists a further perfect subset  $R$  of  $Q$  such that the family  $\{z_\tau^{**} : \tau \in R\}$  is 1-unconditional. By (2) above and identifying  $R$  with  $2^{\mathbb{N}}$ , we conclude that the family  $\{(z_\tau^*, z_\tau^{**}) : \tau \in R\}$  is as desired.

So it only remains to prove the claim. First we argue for (1). Fix  $\tau \in Q$  and pick a sequence  $(\tau_n)$  in  $P$  with  $\tau_n \rightarrow \tau$  and such that  $\tau_n \prec \tau$  for every  $n \in \mathbb{N}$ . By property (2) of Theorem 10, we see that  $y_{\tau_n}^0(x^*) \rightarrow y_\tau^+(x^*)$  for all  $x^* \in B_{X^*}$ . By the bi-orthogonality of the family  $\{(y_\sigma^0, 2x_\sigma^*) : \sigma \in P\}$ , we see that

$$0 = y_{\tau_n}^0(x_\tau^*) \rightarrow y_\tau^+(x_\tau^*)$$

and so  $z_\tau^{**}(z_\tau^*) = 2y_\tau^0(x_\tau^*) = 1$ . Hence,  $z_\tau^{**} \neq 0$ . With identical arguments, we see that for every  $\tau, \tau' \in Q$  with  $\tau \neq \tau'$  we have  $z_\tau^{**}(z_{\tau'}^*) = 0$ . Thus, the family  $\{(z_\tau^*, z_\tau^{**}) : \tau \in Q\}$  forms a bi-orthogonal system in  $X^* \times X^{**}$ , that is, (2) is satisfied. To see (3), it is enough to show that for every sequence  $(\tau_n)$  in  $Q$  with  $t_n \neq t_m$  if  $n \neq m$ , the sequence  $(z_{\tau_n}^{**})$  has a subsequence weak\* convergent to 0. So, let  $(\tau_n)$  be one. By passing to a subsequence, we may assume that there exists  $\tau \in Q$  such

that  $\tau_n \rightarrow \tau$  and either  $\tau_n \prec \tau$  for all  $n \in \mathbb{N}$  or vice versa. We will treat the first case (the argument is symmetric). By property (2) of Theorem 10, we see that both  $(y_{\tau_n}^0)$  and  $(y_{\tau_n}^+)$  are weak\* convergent to  $y_\tau^+$ . Therefore,

$$z_{\tau_n}^{**} = y_{\tau_n}^0 - y_{\tau_n}^+ \xrightarrow{w^*} y_\tau^+ - y_\tau^+ = 0.$$

This shows that  $\{z_\tau^{**} : \tau \in Q\}$  is weak\* discrete having 0 as the unique weak\* accumulation point. Finally, the Borelness of the map  $\Phi$  is an immediate consequence of the Borelness of the maps 0 and + obtained by Theorem 10. This completes the proof of the claim, and so, the entire proof is completed.  $\square$

**5.1. Consequences.** Below we state and prove some consequences of Theorem 1. We start with the following theorem.

**Theorem 13.** *Let  $X$  be a separable Banach space with non-separable dual. Then  $X^{**}$  contains an unconditional family of size  $|X^{**}|$ .*

*Proof.* If  $\ell_1(\mathbb{N})$  embeds into  $X$ , then  $\ell_1(2^{\mathfrak{c}})$  embeds into  $X^{**}$ . Hence,  $X^{**}$  contains an unconditional family of size  $2^{\mathfrak{c}} = |X^{**}|$ . If  $\ell_1(\mathbb{N})$  does not embed into  $X$ , then the cardinality of  $X^{**}$  is equal to the continuum (see [OR]). By Theorem 1, the result follows.  $\square$

The following trichotomy provides the first positive answer to the “reflexivity or unconditionality problem”.

**Theorem 14.** *Let  $X$  be a separable Banach space. Then one of the following holds.*

- (a) *The space  $X$  is saturated with reflexive subspaces.*
- (b) *There exists an unconditional family in  $X^{**}$  of size  $|X^{**}|$ .*
- (c) *There exists an unconditional family in  $X^{***}$  of size  $|X^{***}|$ .*

*Proof.* Let  $X$  be a separable Banach space. If  $X^{**}$  is separable, then by a result stated in [M] and proved in [JR] (see also [EW, Theorem 4.1] for a somewhat more general result), we see that the space  $X$  is reflexive saturated, that is, part (a) holds. So assume that  $X^{**}$  is non-separable. If  $X^*$  is non-separable, then by Theorem 13 we see that (b) is satisfied. Finally, if  $X^*$  is separable, then invoking again Theorem 13 we conclude that (c) holds. The proof is completed.  $\square$

We close this section with the following result which provides a positive answer for the class of dual spaces to Banach’s classical “separable quotient problem”.

**Theorem 15.** *Let  $X$  be a Banach space which is isomorphic to a dual Banach space. Then one of the following holds.*

- (i) *The space  $X$  has the Radon–Nikodym property.*
- (ii) *The space  $X$  has a separable quotient with an unconditional basis.*

*Thus, every dual Banach space has a separable quotient.*



For the proof of Theorem 15 we need the following well-known result ([HJ]). We include the proof for completeness.

**Proposition 16.** *Let  $X$  be a Banach space. If  $X^*$  contains an unconditional basic sequence, then  $X$  has a separable quotient with an unconditional basis.*

*Proof.* Let  $(x_n^*)$  be an unconditional basic sequence in  $X^*$  and set  $R := \overline{\text{span}}\{x_n^* : n \in \mathbb{N}\}$ . By a classical result of James (see [LT]), either  $R$  is reflexive, or  $\ell_1$  embeds into  $R$ , or  $c_0$  embeds into  $R$ . If  $R$  is reflexive, then the weak and weak\* topologies on  $R$  coincide. Hence  $R$  is weak\* linearly homeomorphic to a subspace of  $X^*$ , which yields that  $X$  maps onto  $R^*$ . Also observe that if  $\ell_1$  embeds into  $X$ , then  $L^1[0, 1]$  embeds into  $X^*$  (see [Pe]). Therefore,  $\ell_2$  embeds into  $X^*$  which implies that  $\ell_2$  is a quotient of  $X$ .

From now on we assume that  $\ell_1$  does not embed into  $X$ . By [BP], we conclude that  $c_0$  does not embed into  $X^*$ . Hence,  $R$  does not contain  $c_0$ . What remains is to treat the case where  $\ell_1$  embeds into  $R$ . Since  $\ell_1$  does not embed into  $X$ , by [HJ], we conclude that there exists a weak\* null sequence  $(z_n^*)$  in  $R$  equivalent to the usual basis of  $\ell_1$ . Denote by  $T: \ell_1 \rightarrow \overline{\text{span}}\{z_n^* : n \in \mathbb{N}\} \hookrightarrow X^*$  the natural isomorphism and let  $T^*: X^{**} \rightarrow \ell_\infty$  be the dual onto operator. Observe that  $T^*|_X$  maps  $X$  to  $c_0$ , and so, it is weak\*-weak continuous. It follows that  $T^*$  maps  $X$  onto  $c_0$ . This completes the proof.  $\square$

We are ready to proceed to the proof of Theorem 15.

*Proof of Theorem 15.* Let  $Y$  be a Banach space such that  $X$  is isomorphic to  $Y^*$ . Assume that (i) does not hold. It follows that there exists a separable subspace  $Z$  of  $Y$  such that  $Z^*$  is non-separable (see [St]). By Theorem 13, we see that  $Z^{**}$  contains an unconditional basic sequence. Hence, so does  $X^*$ . By Proposition 16, we conclude that  $X$  has a separable quotient with an unconditional basis and the result follows.  $\square$

Let us mention that Todorćević has shown that there exists a model of set theory where the continuum hypothesis fails and in which every Banach space of density character  $\aleph_1$  has a separable quotient ([To3]).

## 6. TREE BASES IN BANACH SPACES

We start with the following theorem.

**Theorem 17.** *Let  $X$  be a separable Banach space not containing  $\ell_1$  and such that  $X^*$  is non-separable. Then there exists a seminormalized family  $(e_t)_{t \in 2^{<\mathbb{N}}}$  such that the following are satisfied.*

- (1) *For every  $\sigma \in 2^{\mathbb{N}}$  the sequence  $(e_{\sigma|_n})$  is weak\* convergent to an element  $z_\sigma^{**} \in X^{**}$ .*
- (2) *For every antichain  $A$  of  $2^{<\mathbb{N}}$  the sequence  $(e_t)_{t \in A}$  is weakly-null.*

- (3) The family  $\{z_\sigma^{**} : \sigma \in 2^{\mathbb{N}}\}$  is weak\* discrete and has 0 as the unique weak\* accumulation point.

Theorem 17 follows by the general structural result obtained in [ADK] and concerning the behavior of non- $G_\delta$  points in a large class of Rosenthal compacta. The proof, however, given in [ADK] uses deep results from the theory of Rosenthal compacta and it is rather involved. The one we present below is based on Stegall's construction as well as on the analysis behind the proof of Theorem 1.

*Proof of Theorem 17.* First we argue as in the proof of Theorem 1. Specifically, applying Theorem 12 for  $\varepsilon = 1$  we obtain  $\Delta_1 = (x_t)_{t \in 2^{<\mathbb{N}}}$  and  $D_1 = \{x_\sigma^* : \sigma \in 2^{\mathbb{N}}\}$ . Next, we apply Theorem 10 for the family  $\Delta_1$  and we obtain a regular dyadic subtree  $T$  of  $2^{<\mathbb{N}}$  and a family  $\{y_\sigma^0, y_\sigma^+, y_\sigma^- : \sigma \in P\}$ , where  $P = [T]$ , as described in Theorem 10. Without loss of generality and by re-enumerating if necessary (which can be done since the tree  $T$  is regular dyadic), we may assume that  $T = 2^{<\mathbb{N}}$  and so  $P = 2^{\mathbb{N}}$ .

We fix a regular dyadic subtree  $R = (r_t)_{t \in 2^{<\mathbb{N}}}$  of  $2^{<\mathbb{N}}$  with the following property.

- (P) For every  $t \in R$  we have  $t \hat{\cap} 0 \notin \hat{R}$  while  $t \hat{\cap} 1 \in \hat{R}$ .

A possible choice can be as follows. For every  $t = (\varepsilon_0, \dots, \varepsilon_k) \in 2^{<\mathbb{N}}$  set  $r_t := (1, \varepsilon_0, 1, \varepsilon_1, \dots, 1, \varepsilon_k)$  if  $t \neq \emptyset$  and  $r_\emptyset = \emptyset$ . It is easy to see that  $R = (r_t)_{t \in 2^{<\mathbb{N}}}$  satisfies (P) above. We denote by  $Q$  the body of  $\hat{R}$ . For every  $\sigma \in 2^{\mathbb{N}}$  let  $\tau_\sigma = \bigcup_n r_{\sigma|n} \in Q$ . The map  $2^{\mathbb{N}} \ni \sigma \mapsto \tau_\sigma \in Q$  is a homeomorphism. We isolate the following properties of  $R$  and  $Q$ .

- (a) If  $t \in 2^{<\mathbb{N}}$  and  $\sigma \in 2^{\mathbb{N}}$  with  $r_t \prec \tau_\sigma$  (respectively,  $\tau_\sigma \prec r_t$ ), then  $r_t \hat{\cap} 0 \prec \tau_\sigma$  (respectively,  $\tau_\sigma \prec r_t \hat{\cap} 0$ ).
- (b) If  $(t_n)$  is a sequence in  $2^{<\mathbb{N}}$  and  $\sigma \in 2^{\mathbb{N}}$  are such that  $r_{t_n} \rightarrow \tau_\sigma$ , then  $r_{t_n} \hat{\cap} 0 \rightarrow \tau_\sigma$ .
- (c) For every  $\sigma \in 2^{\mathbb{N}}$ , the sequence  $(r_{\sigma|n} \hat{\cap} 0)$  is an increasing antichain converging to  $\tau_\sigma$ .

For every  $t \in 2^{<\mathbb{N}}$  we define

$$e_t := x_{r_t} - x_{r_t \hat{\cap} 0}.$$

We claim that the family  $(e_t)_{t \in 2^{<\mathbb{N}}}$  is the desired one. Using (c) above and properties (1) and (4) of Theorem 10, we see that for every  $\sigma \in 2^{\mathbb{N}}$  the sequence  $(e_{\sigma|n})$  is weak\* convergent to the element

$$z_\sigma^{**} := y_{\tau_\sigma}^0 - y_{\tau_\sigma}^+ \in X^{**}.$$

With identical arguments as in the proof of Theorem 1, we see that the family  $\{z_\sigma^{**} : \sigma \in 2^{\mathbb{N}}\}$  is weak\* discrete having 0 as the unique weak\* accumulation point. Hence (1) and (3) in the statement of the theorem are satisfied. Let us see that (2) is also satisfied. Notice that it is enough to prove that for every infinite antichain  $A$  of  $2^{<\mathbb{N}}$  there exists  $B \subseteq A$  infinite such that the sequence  $(e_t)_{t \in B}$  is weakly-null. So let  $A$  be one. There exist  $\sigma \in 2^{\mathbb{N}}$  and an infinite subset  $B$  of  $A$  such that  $B \rightarrow \sigma$

and either  $t \prec \sigma$  for all  $t \in B$  or vice versa. Assume that the first case occurs (the argument is symmetric). Observe that  $r_t \prec \tau_\sigma$  for every  $t \in B$ . By (a) and (b) above and property (4) in Theorem 10, we obtain that

$$w^* - \lim_{t \in B} e_t = w^* - \lim_{t \in B} (x_{t_t} - x_{r_t \wedge 0}) = y_{\tau_\sigma}^+ - y_{\tau_\sigma}^+ = 0.$$

Therefore, the sequence  $(e_t)_{t \in B}$  is weakly-null and the proof is completed.  $\square$

Actually, we can considerably strengthen the properties of the sequence  $(e_t)_{t \in 2^{< \mathbb{N}}}$  obtained by Theorem 17, as follows.

**Theorem 18.** *Let  $X$  be a separable Banach space not containing  $\ell_1$  and with non-separable dual. Then there exist a family  $(w_t)_{t \in 2^{< \mathbb{N}}}$  in  $X$  and a family  $\{w_\sigma^{**} : \sigma \in 2^{\mathbb{N}}\}$  in  $X^{**}$  satisfying (1), (2) and (3) of Theorem 17 as well as the following properties.*

- (i) *The family  $(w_t)_{t \in 2^{< \mathbb{N}}}$  is basic when it is enumerated appropriately.*
- (ii) *The family  $\{w_\sigma^{**} : \sigma \in 2^{\mathbb{N}}\}$  is 1-unconditional.*
- (iii) *For every  $n \geq 1$  if  $t_1 \prec \dots \prec t_{2^n}$  is the  $\prec$ -increasing enumeration of  $2^n$ , then for every  $\{\sigma_1, \dots, \sigma_{2^n}\} \subseteq 2^{\mathbb{N}}$  with  $t_i \sqsubset \sigma_i$  the families  $\{w_{t_i}\}_{i=1}^{2^n}$  and  $\{w_{\sigma_i}^{**}\}_{i=1}^{2^n}$  are  $(1 + \frac{1}{n})$ -equivalent.*
- (iv) *For every  $n \geq 1$  the family  $\{w_t : t \in 2^n\}$  is  $(1 + \frac{1}{n})$ -unconditional.*

The proof of Theorem 18 is based on Theorem 17, as well as, on the following lemmas. In the first one we use Theorem 9 in a similar way as in the proof of Lemma 11.

**Lemma 19.** *Let  $X$  be a separable Banach space. Also let  $Q$  be a perfect subset of  $2^{\mathbb{N}}$  and let  $\{z_\sigma^{**} : \sigma \in Q\}$  be a bounded family in  $X^{**}$ . Assume that the map  $\Phi: Q \times (B_{X^*}, w^*) \rightarrow \mathbb{R}$  defined by  $\Phi(\sigma, x^*) = z_\sigma^{**}(x^*)$  is Borel. Let  $n \in \mathbb{N}$  and let  $Q_0, \dots, Q_n$  be pairwise disjoint perfect subsets of  $Q$ . Then, for every  $\varepsilon > 0$  there exist perfect subsets  $P_0, \dots, P_n$  of  $Q_0, \dots, Q_n$  such that*

$$\left| \left\| \sum_{i=0}^n \lambda_i z_{\sigma_i}^{**} \right\| - \left\| \sum_{i=0}^n \lambda_i z_{\tau_i}^{**} \right\| \right| < \varepsilon$$

for every  $(\sigma_i)_{i=0}^n$  and  $(\tau_i)_{i=0}^n$  in  $P_0 \times \dots \times P_n$  and every  $(\lambda_i)_{i=0}^n$  in  $[-1, 1]^{n+1}$ .

*Proof.* Let  $\delta > 0$  be sufficiently small which will be determined later, and let  $\Lambda \subseteq [-1, 1]$  and  $N \subseteq [0, (n+1)M]$  be finite  $\delta$ -nets where  $M > 0$  is such that  $\|z_\sigma^{**}\| \leq M$  for all  $\sigma \in Q$ . For every  $(a_i)_{i=0}^n$  in  $\Lambda^{n+1}$  and every  $a \in N$  set

$$D(a_0, \dots, a_n, a) := \left\{ (\sigma_0, \dots, \sigma_n) \in Q_0 \times \dots \times Q_n : a - \delta < \left\| \sum_{i=0}^n a_i z_{\sigma_i}^{**} \right\| < a + \delta \right\}.$$

Arguing as in the proof of Claim 1 in Lemma 11, it is easy to verify that the set  $D(a_0, \dots, a_n, a)$  belongs to the  $\sigma$ -algebra generated by the analytic sets, and so, it

has the Baire property in  $Q_0 \times \cdots \times Q_n$ . It is easy to see that for every  $(a_i)_{i=0}^n$  in  $\Lambda^{n+1}$  we have

$$Q_0 \times \cdots \times Q_n = \bigcup_{a \in N} D(a_0, \dots, a_n, a).$$

Applying successively Theorem 9 for every  $(a_0, \dots, a_n) \in \Lambda^{n+1}$ , we obtain perfect subsets  $P_0, \dots, P_n$  of  $Q_0, \dots, Q_n$  such that the following property is satisfied. For every  $(a_0, \dots, a_n) \in \Lambda^{n+1}$  there exists unique  $a \in N$  such that  $P_0 \times \cdots \times P_n \subseteq D(a_0, \dots, a_n, a)$ . We claim that the perfect set  $P_0, \dots, P_n$  satisfy the conclusion of the lemma for a sufficiently small  $\delta$ . Indeed, for every  $(\sigma_i)_{i=0}^n$  and  $(\tau_i)_{i=0}^n$  in  $P_0 \times \cdots \times P_n$  and every  $(a_i)_{i=0}^n$  in  $\Lambda^{n+1}$  we have

$$-2\delta \leq \left\| \sum_{i=0}^n a_i z_{\sigma_i}^{**} \right\| - \left\| \sum_{i=0}^n a_i z_{\tau_i}^{**} \right\| \leq 2\delta.$$

Using this, it is easy to check that for every  $(\lambda_i)_{i=0}^n$  in  $[-1, 1]^{n+1}$  we have

$$\left| \left\| \sum_{i=0}^n \lambda_i z_{\sigma_i}^{**} \right\| - \left\| \sum_{i=0}^n \lambda_i z_{\tau_i}^{**} \right\| \right| \leq 2(n+1)\delta M + 2\delta.$$

Choosing  $\delta > 0$  so that  $2(n+1)\delta M + 2\delta < \varepsilon$ , the lemma is proved.  $\square$

**Lemma 20.** *Let  $X$  be a separable Banach space not containing  $\ell_1$ . Let  $n \in \mathbb{N}$  and  $(z_i^{**})_{i=0}^n$  in  $X^{**}$ . For every  $i \in \{0, \dots, n\}$  let  $(e_k^i)$  be a sequence in  $X$  which is weak\* convergent to  $z_i^{**}$ . Then, for every  $\varepsilon > 0$  there exist  $w_0, \dots, w_n$  finite convex combinations of  $(e_k^0), \dots, (e_k^n)$  respectively, such that*

$$\left| \left\| \sum_{i=0}^n \lambda_i w_i \right\| - \left\| \sum_{i=0}^n \lambda_i z_i^{**} \right\| \right| < \varepsilon$$

for every  $(\lambda_i)_{i=0}^n$  in  $[-1, 1]^{n+1}$ .

*Proof.* We will need the following claim.

**Claim.** *Let  $d \in \mathbb{N}$  and  $y_0^{**}, \dots, y_d^{**}$  in  $X^{**}$ . For every  $j \in \{0, \dots, d\}$  let  $(y_k^j)$  be a sequence in  $X$  which is weak\* convergent to  $y_j^{**}$ . Then, for every  $\theta > 0$  there exist  $k_0 \in \mathbb{N}$  and  $\mu_0, \dots, \mu_{k_0}$  in  $[0, 1]$  with  $\sum_{k=0}^{k_0} \mu_k = 1$  such that*

$$(2) \quad \left| \left\| \sum_{k=0}^{k_0} \mu_k y_k^j \right\| - \|y_j^{**}\| \right| < \theta$$

for every  $j \in \{0, \dots, d\}$ .

Granting the claim, we proceed as follows. Let  $\delta > 0$  be sufficiently small, which we will determine later, and let  $\Lambda$  be a finite  $\delta$ -net in  $[-1, 1]$ . Let  $(a_i^0)_{i=0}^n, \dots, (a_i^d)_{i=0}^n$  be an enumeration of the set  $\Lambda^{n+1}$ . For every  $j \in \{0, \dots, d\}$  and every  $k \in \mathbb{N}$  we set  $y_j^{**} = \sum_{i=0}^n a_i^j z_i^{**}$  and  $y_k^j = \sum_{i=0}^n a_i^j e_k^i$ . Notice that the sequence  $(y_k^j)$  is weak\* convergent to  $y_j^{**}$  for every  $j \in \{0, \dots, d\}$ . We apply the above claim for  $\theta = \frac{\varepsilon}{2}$  and

we obtain  $k_0 \in \mathbb{N}$  and  $\mu_0, \dots, \mu_{k_0}$  in  $[0, 1]$  satisfying inequality (2) above. For every  $i \in \{0, \dots, n\}$  we set

$$w_i = \sum_{k=0}^{k_0} \mu_k e_k^i.$$

Notice that for every  $j \in \{0, \dots, d\}$  we have

$$\sum_{k=0}^{k_0} \mu_k y_k^j = \sum_{k=0}^{k_0} \mu_k \left( \sum_{i=0}^n a_i^j e_k^i \right) = \sum_{i=0}^n a_i^j \left( \sum_{k=0}^{k_0} \mu_k e_k^i \right) = \sum_{i=0}^n a_i^j w_i.$$

Hence, inequality (2) is reformulated as follows. For every  $j \in \{0, \dots, d\}$  we have

$$\left| \left\| \sum_{i=0}^n a_i w_i \right\| - \left\| \sum_{i=0}^n a_i z_i^{**} \right\| \right| < \frac{\varepsilon}{2}.$$

Let  $M > 0$  be such that  $\|e_k^i\| \leq M$  and  $\|z_i^{**}\| \leq M$  for every  $i \in \{0, \dots, n\}$  and every  $k \in \mathbb{N}$ . It follows that for every  $(\lambda_i)_{i=0}^n$  in  $[-1, 1]^{n+1}$  we have

$$\left| \left\| \sum_{i=0}^n \lambda_i w_i \right\| - \left\| \sum_{i=0}^n \lambda_i z_i^{**} \right\| \right| \leq 2(n+1)\delta M + \frac{\varepsilon}{2}.$$

Hence, by choosing  $\delta$  sufficiently small, the result follows.

It remains to prove the claim. For every  $j \in \{0, \dots, d\}$  we select  $x_j^* \in X^*$  with  $\|x_j^*\| = 1$  such that  $\|y_j^{**}\| - \frac{\theta}{4} < y_j^{**}(x_j^*)$ . By [OR], for every  $j \in \{0, \dots, d\}$  we may select a sequence  $(x_k^j)$  in  $X$  satisfying the following.

- (a) The sequence  $(x_k^j)$  is weak\* convergent to  $y_j^{**}$ .
- (b) For every  $k \in \mathbb{N}$  we have  $\|x_k^j\| \leq \|y_j^{**}\|$ .
- (c) For every  $k \in \mathbb{N}$  we have  $|x_k^j(x_k^j) - y_j^{**}(x_k^j)| < \frac{\theta}{4}$ .

Notice that for every convex combination  $w$  of  $(x_k^j)$  we have

$$(3) \quad \|y_j^{**}\| - \frac{\theta}{2} \leq \|w\| \leq \|y_j^{**}\|.$$

For every  $j \in \{0, \dots, d\}$  and every  $k \in \mathbb{N}$  we set  $d_k^j := y_k^j - x_k^j$ . Observe that the sequence  $(d_k^j)$  is weakly-null. Applying successively Mazur's theorem (for every  $j$ ), we obtain  $k_0 \in \mathbb{N}$  and  $\mu_0, \dots, \mu_{k_0}$  in  $[0, 1]$  with  $\sum_{k=0}^{k_0} \mu_k = 1$  such that for every  $j \in \{0, \dots, d\}$  we have

$$(4) \quad \left\| \sum_{k=0}^{k_0} \mu_k d_k^j \right\| < \frac{\theta}{4}.$$

Since

$$\left| \left\| \sum_{k=0}^{k_0} \mu_k y_k^j \right\| - \left\| \sum_{k=0}^{k_0} \mu_k x_k^j \right\| \right| \leq \left\| \sum_{k=0}^{k_0} \mu_k d_k^j \right\|,$$

by inequalities (3) and (4) above, the proof of the claim follows and the lemma is proved.  $\square$

We recall that a subset  $I$  of  $2^{<\mathbb{N}}$  is said to be a (finite) segment if there exist  $s, t \in 2^{<\mathbb{N}}$  with  $s \sqsubseteq t$  and such that  $I = \{w : s \sqsubseteq w \sqsubseteq t\}$ . If  $I = \{w : s \sqsubseteq w \sqsubseteq t\}$  is a segment, then we set  $\min(I) := s$  and  $\max(I) := t$ . By  $\phi: 2^{<\mathbb{N}} \rightarrow \mathbb{N}$  we denote the unique bijection satisfying  $\phi(s) < \phi(t)$  if either  $|s| < |t|$ , or  $|s| = |t|$  and  $s \prec t$  for all  $s, t \in 2^{<\mathbb{N}}$ . For every  $t \in 2^{<\mathbb{N}}$  by  $V_t$  we denote the clopen subset  $\{\sigma : t \sqsubset \sigma\}$  of  $2^{\mathbb{N}}$ . We are ready to proceed to the proof of Theorem 18.

*Proof of Theorem 18.* First, we start with the families  $(e_t)_{t \in 2^{<\mathbb{N}}}$  and  $\{z_\sigma^{**} : \sigma \in 2^{\mathbb{N}}\}$  obtained by Theorem 17. Using Theorem 4 and by passing to regular dyadic subtree if necessary, we may assume that the family  $\{z_\sigma^{**} : \sigma \in 2^{\mathbb{N}}\}$  is 1-unconditional. We observe the following. For every  $t \in 2^{<\mathbb{N}}$  there exists an infinite antichain  $(s_n)$  of  $2^{<\mathbb{N}}$  such that  $t \sqsubset s_n$  for every  $n \in \mathbb{N}$ . By property (2) of Theorem 17, we see that  $(e_{s_n})$  is weakly-null. Hence, considering the space  $X$  as a subspace of  $C[0, 1]$ , using a standard sliding hump argument and by passing to a dyadic (but not necessarily regular) subtree of  $2^{<\mathbb{N}}$ , we may assume the following.

- (i) If  $(t_n)$  is the enumeration of  $2^{<\mathbb{N}}$  according to  $\phi$ , then the sequence  $(e_{t_n})$  is Schauder basic.

Let  $(\varepsilon_n)$  be a decreasing sequence of positive reals converging sufficiently fast to zero. By recursion on the length of finite sequences in  $2^{<\mathbb{N}}$ , we shall construct

- (C1) a Cantor scheme  $(P_t)_{t \in 2^{<\mathbb{N}}}$  of perfect subsets of  $2^{\mathbb{N}}$ ,
- (C2) a family  $(I_t)_{t \in 2^{<\mathbb{N}}}$  of segments of  $2^{<\mathbb{N}}$ , and
- (C3) a family  $(w_t)_{t \in 2^{<\mathbb{N}}}$  of convex combinations of  $(e_t)_{t \in 2^{<\mathbb{N}}}$ .

The construction is done so that for every  $t \in 2^{<\mathbb{N}}$  the following are satisfied.

- (P1)  $w_t$  is a convex combination of  $\{e_s : s \in I_t\}$ .
- (P2)  $P_t \subseteq V_{\max(I_t)}$ .
- (P3) For every  $\epsilon \in \{0, 1\}$  we have  $\max(I_t \hat{\ } \epsilon) \sqsubseteq \min(I_t \smallfrown \epsilon)$ .
- (P4) For every  $s, t \in 2^{<\mathbb{N}}$  we have  $|s| < |t|$  if and only if  $|\max(I_s)| < |\min(I_t)|$ .
- (P5) For every  $n \in \mathbb{N}$  and every  $(\sigma_t)_{t \in 2^{2^n}}$  and  $(\tau_t)_{t \in 2^{2^n}}$  in  $\prod_{t \in 2^{2^n}} P_t$  we have
  - (a)  $(z_{\sigma_t}^{**})_{t \in 2^{2^n}}$  is  $(1 + \varepsilon_n)$ -equivalent to  $(z_{\tau_t}^{**})_{t \in 2^{2^n}}$ , and
  - (b)  $(z_{\sigma_t}^{**})_{t \in 2^{2^n}}$  is  $(1 + \varepsilon_n)$ -equivalent  $(w_t)_{t \in 2^{2^n}}$ .

Using Lemma 19 and Lemma 20, one can easily realize that such a construction can be carried out.

For every  $\sigma \in 2^{\mathbb{N}}$  let  $\tau_\sigma$  be the unique element of  $2^{\mathbb{N}}$  determined by the infinite chain  $\bigcup_n I_{\sigma|n}$ . Clearly the sequence  $(w_{\sigma|n})$  is weak\* convergent to  $w_\sigma^{**} := z_{\tau_\sigma}^{**}$ . It is easy to check using properties (P3), (P4) and (P5) above that the families  $(w_t)_{t \in 2^{<\mathbb{N}}}$  and  $\{w_\sigma^{**} : \sigma \in 2^{\mathbb{N}}\}$  are as desired. The proof is completed.  $\square$

## REFERENCES

- [AAK] D. Apatsidis, S. A. Argyros and V. Kanellopoulos, *On the subspaces of  $JF$  and  $JT$  with non-separable dual*, J. Funct. Anal. 254 (2008), 632–674.

- [AAT] S. A. Argyros, A. D. Arvanitakis and A. G. Toliaş, *Saturated extensions, the attractors method and hereditarily James tree spaces*, in “Methods in Banach spaces”, LMS Lecture Notes, Vol. 337, 1–90.
- [ADK] S. A. Argyros, P. Dodos and V. Kanellopoulos, *A classification of separable Rosenthal compacta and its applications*, *Dissertationes Math.* 449 (2008), 1–52.
- [AFK] S. A. Argyros, V. Felouzis and V. Kanellopoulos, *A proof of Halpern-L auchli Partition Theorem*, *Europ. J. Comb.* 23 (2002), 1–10.
- [AGR] S. A. Argyros, G. Godefroy and H. P. Rosenthal, *Descriptive Set Theory and Banach spaces*, *Handbook of the Geometry of Banach Spaces*, Vol. 2, edited by W. B. Johnson and J. Lindenstrauss, Elsevier, 2003.
- [AT] S. A. Argyros and S. Todor evi c, *Ramsey Methods in Analysis*, *Advanced Courses in Mathematics*, CRM Barcelona, Birkh user, Verlag, Basel, 2005.
- [ATo] S. A. Argyros and A. Toliaş, *Methods in the Theory of Hereditarily Indecomposable Banach spaces*, *Memoirs Amer. Math. Soc.* 806, 2004.
- [BP] C. Bessaga and A. Pełczyński, *On bases and unconditional convergence of series in Banach spaces*, *Studia Math.* 17 (1958), 151–164.
- [B] A. Blass, *A partition for perfect sets*, *Proc. Amer. Math. Soc.* 82 (1981), 271–277.
- [C] T. J. Carlson, *Some unifying principles in Ramsey theory*, *Discrete Math.* 68 (1988), 117–169.
- [EW] G. A. Edgar and R. F. Wheeler, *Topological properties of Banach spaces*, *Pacific J. Math.* 115 (1984), 317–350.
- [E] E. Ellentuck, *A new proof that analytic sets are Ramsey*, *J. Symb. Logic* 39 (1974), 163–165.
- [F] J. Farahat, *Espaces  $bc$  Banach contenant  $\ell_1$  d’apres H. P. Rosenthal*, *Seminaire Maurey-Schwartz*, Ecole Polytechnique, 1973–74.
- [GL] G. Godefroy and A. Louveau, *Axioms of determinacy and biorthogonal systems*, *Israel J. Math.* 67 (1989), 109–116.
- [GT] G. Godefroy and M. Talagrand, *Espaces de Banach repr esentable*, *Israel J. Math.* 41 (1982), 321–330.
- [G1] W. T. Gowers, *An infinite Ramsey theorem and some Banach-space dichotomies*, *Ann. Math.* 156 (2002), 797–833.
- [G2] W. T. Gowers, *Ramsey methods in Banach spaces*, *Handbook of the Geometry of Banach Spaces*, Vol. 2, edited by W. B. Johnson and J. Lindenstrauss, Elsevier, 2003.
- [GM] W. T. Gowers and B. Maurey, *The unconditional basic sequence problem*, *J. Amer. Math. Soc.* 6 (1993), 851–874.
- [HJ] J. N. Hagler and W. B. Johnson, *On Banach spaces whose dual balls are not weak\* sequentially compact*, *Israel J. Math.* 28 (1977), 325–330.
- [HL] J. D. Halpern and H. L auchli, *A partition theorem*, *Trans. Amer. Math. Soc.* 124 (1966), 360–367.
- [J] R. C. James, *A separable somewhat reflexive Banach space with non-separable dual*, *Bull. Amer. Math. Soc.* 80 (1974), 738–743.
- [JR] W. B. Johnson and H. P. Rosenthal, *On weak\* basic sequences and their applications to the study of Banach spaces*, *Studia Math.* 43 (1972), 77–92.
- [Ka] V. Kanellopoulos, *Ramsey families of subtrees of the dyadic tree*, *Trans. Amer. Math. Soc.* 357 (2005), 3865–3886.
- [Ke] A. S. Kechris, *Classical Descriptive Set Theory*, *Grad. Texts in Math.*, Vol. 156, Springer-Verlag, 1995.
- [LT] J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces I and II*, Springer, 1996.
- [LSV] A. Louveau, S. Shelah and B. Velickovi c, *Borel partitions of infinite subtrees of a perfect tree*, *Annals Pure Appl. Logic* 63 (1993), 271–281.

- [Mau] B. Maurey, *Types and  $\ell_1$  subspaces*, Texas functional analysis seminar 1982–1983 (Austin, Tex.), 123–137, Longhorn Notes, Univ. Texas Press, Austin, TX, 1983.
- [Mil] K. Milliken, *A partition theorem for the infinite subtrees of a tree*, Trans. Amer. Math. Soc. 263 (1981), 137–148.
- [Mi] A. W. Miller, *Infinite combinatorics and definability*, Annals Pure Appl. Logic 41 (1989), 179–203.
- [M] V. D. Milman, *Geometric theory of Banach spaces. I. Theory of basic and minimal systems*, (Russian) Uspehi Mat. Nauk 25 (1970), 113–174.
- [Od] E. Odell, *Applications of Ramsey theorems to Banach space theory*, Notes in Banach spaces, 379–404, Univ. Texas Press, Austin, Texas, 1980.
- [OR] E. Odell and H. P. Rosenthal, *A double-dual characterization of separable Banach spaces not containing  $\ell_1$* , Israel J. Math. 20 (1975), 375–384.
- [Pa] J. Pawlikowski, *Parametrized Ellentuck Theorem*, Topol. Appl. 37 (1990), 65–73.
- [Pe] A. Pelczyński, *On Banach spaces containing  $L_1(\mu)$* , Studia Math. 30 (1968), 231–246.
- [Ro] H. P. Rosenthal, *A characterization of Banach spaces containing  $\ell_1$* , Proc. Nat. Acad. Sci. USA 71 (1974), 2411–2413.
- [St] Ch. Stegall, *The Radon-Nikodym property in conjugate Banach spaces*, Trans. Amer. Math. Soc. 206 (1975), 213–223.
- [Ste] J. Stern, *A Ramsey theorem for trees with an application to Banach spaces*, Israel J. Math. 29 (1978), 179–188.
- [To1] S. Todorčević, *Compact subsets of the first Baire class*, J. Amer. Math. Soc. 12 (1999), 1179–1212.
- [To2] S. Todorčević, *Introduction to Ramsey Spaces*, Annals Math. Studies, No. 174, Princeton Univ. Press, 2010.
- [To3] S. Todorčević, *Biorthogonal systems and quotient spaces via Baire category methods*, Math. Annalen 335 (2006), 687–715.

NATIONAL TECHNICAL UNIVERSITY OF ATHENS, FACULTY OF APPLIED SCIENCES, DEPARTMENT OF MATHEMATICS, ZOGRAFOU CAMPUS, 157 80, ATHENS, GREECE.

*E-mail address:* sargyros@math.ntua.gr, pdodos@math.ntua.gr, bkanel@math.ntua.gr