

# ON ANTICHAINS OF SPREADING MODELS OF BANACH SPACES

PANDELIS DODOS

ABSTRACT. We show that for every separable Banach space  $X$  either  $\text{SP}_w(X)$  (the set of all spreading models of  $X$  generated by weakly-null sequences in  $X$ , modulo equivalence) is countable, or  $\text{SP}_w(X)$  contains an antichain of the size of the continuum. This answers a question of Dilworth, Odell and Sari.

## 1. INTRODUCTION

Let  $X$  be a separable Banach space and let  $\text{SP}_w(X)$  denote the set of all spreading models of  $X$  generated by weakly-null sequences in  $X$ , modulo equivalence. By  $\leq$  we denote the usual relation on  $\text{SP}_w(X)$  of domination. The study of the structure  $(\text{SP}_w(X), \leq)$  was initiated by Androulakis, Odell, Schlumprecht and Tomczak-Jaegermann [AOST]. They showed, for instance, that  $(\text{SP}_w(X), \leq)$  is a semi-lattice, that is, any two elements of  $\text{SP}_w(X)$  admit a least upper bound. The question of determining which countable lattices can be realized as  $(\text{SP}_w(X), \leq)$ , for some separable Banach space  $X$ , was answered by Dilworth, Odell and Sari [DOS].

This note is motivated by the following problem posed by the authors of [DOS] (see [DOS, Problem 1.13]).

**Problem 1.** *If  $\text{SP}_w(X)$  is uncountable must there exist  $\{(x_n^\xi) : \xi < \omega_1\}$  in  $\text{SP}_w(X)$  which is either strictly increasing with respect to  $\xi$ , or strictly decreasing, or consists of mutually incomparable elements?*

To state our first result, let us say that a seminormalized basic sequence  $(x_n)$  in a Banach space  $X$  is *C-Schreier spreading* for some  $C \geq 1$  (or simply *Schreier spreading*, if  $C$  is understood) if for every  $k \in \mathbb{N}$ , every  $k \leq n_0 < \dots < n_k$  and every  $k \leq m_0 < \dots < m_k$  we have that  $(x_{n_i})_{i=0}^k$  is  $C$ -equivalent to  $(x_{m_i})_{i=0}^k$ . Observe that if  $(x_n)$  is Schreier spreading, then there exists a unique spreading model (up to equivalence) generated by subsequences of  $(x_n)$ . Also let  $2^{<\mathbb{N}}$  denote the Cantor tree and let  $\varphi: 2^{<\mathbb{N}} \rightarrow \mathbb{N}$  be the unique bijection satisfying  $\varphi(s) < \varphi(t)$  if either  $|s| < |t|$ , or  $|s| = |t| = n$  and  $s <_{\text{lex}} t$ . (Here,  $<_{\text{lex}}$  stands for the usual lexicographical order on  $2^n$ .)

We show the following theorem.

---

2000 *Mathematics Subject Classification*: 46B20, 03E15.

**Theorem 1.** *Let  $X$  be a separable Banach space such that  $\text{SP}_w(X)$  is uncountable. Then there exist a family  $(x_t)_{t \in 2^{<\mathbb{N}}}$  in  $X$  and  $C \geq 1$  such that the following hold.*

- (1) *If  $(t_n)$  is the enumeration of  $2^{<\mathbb{N}}$  according to  $\varphi$ , then the sequence  $(x_{t_n})$  is a seminormalized basic sequence.*
- (2) *For every  $\sigma \in 2^{\mathbb{N}}$  the sequence  $(x_{\sigma|n})$  is weakly-null and  $C$ -Schreier spreading.*
- (3) *For every  $\sigma, \tau \in 2^{\mathbb{N}}$  with  $\sigma \neq \tau$  if  $(y_n^\sigma)$  and  $(y_n^\tau)$  are spreading models generated by subsequences of  $(x_{\sigma|n})$  and  $(x_{\tau|n})$  respectively, then  $(y_n^\sigma)$  and  $(y_n^\tau)$  are incomparable with respect to domination.*

Theorem 1 yields the following corollary.

**Corollary 2.** *Let  $X$  be a separable Banach space such that  $\text{SP}_w(X)$  is uncountable. Then  $\text{SP}_w(X)$  contains an antichain of the size of the continuum.*

We notice that, independently, Ferenczi and Rosendal have proved Corollary 2 under the additional assumption that  $X$  has separable dual ([FR]).

In [AOST] (see also [DOS]), it was shown that  $\text{SP}_w(X)$  can contain a strictly decreasing infinite sequence, yet no strictly increasing infinite sequence can be found in  $\text{SP}_w(X)$ . This is not, however, the case of the uncountable.

**Theorem 3.** *Let  $X$  be a separable Banach space.*

- (a) *If  $\text{SP}_w(X)$  contains a strictly decreasing sequence of length  $\omega_1$ , then  $\text{SP}_w(X)$  contains a strictly increasing sequence of length  $\omega_1$ .*

*On the other hand,*

- (b) *if  $\text{SP}_w(X)$  does not contain a strictly increasing infinite sequence, then there exists a countable ordinal  $\xi_X$  such that  $\text{SP}_w(X)$  does not contain strictly decreasing sequences of order type greater than  $\xi_X$ .*

It was shown in [DOS, Theorem 3.7] that for every countable ordinal  $\xi$  there exists a separable Banach space  $X_\xi$  such that  $(\text{SP}_w(X_\xi), \leq)$  does not contain a strictly increasing infinite sequence, yet  $\text{SP}_w(X_\xi)$  contains a strictly decreasing sequence of order type  $\xi$ . Thus, the ordinal  $\xi_X$  obtained by part (b) of Theorem 3 is not uniformly bounded within the class of separable Banach spaces for which  $\text{SP}_w(X)$  does not contain a strictly increasing infinite sequence.

In the proofs of Theorem 1 and part (a) of Theorem 3 we use the structural result obtained by Sari in [Sa]. The central argument, however, in the proof of Theorem 1 is essentially based on the work of Leo Harrington and Saharon Shelah on Borel orders. Deep as it is, the theory developed by Harrington and Shelah is highly sophisticated. In particular, all known proofs of their results use either effective descriptive set theory or forcing. However, for the proof of Theorem 1 we need only some instances of the theory and merely for  $F_\sigma$  orders. Thus, we have included “elementary” proofs of all the results that we need, making the paper

essentially self-contained and accessible to anyone with basic knowledge of classical descriptive set theory. None of these proofs should be considered as a contribution to the field of Borel orders.

The paper is organized as follows. In Section 2 we state and prove all results on Borel orders that are needed for the proof of Theorem 1. In Section 3 we show that for every separable Banach space  $X$  the structure  $(\text{SP}_w(X), \leq)$  can be realized as an  $F_\sigma$  order. In Section 4 we give the proof of Theorem 1 while the proof of Theorem 3 is given in Section 5.

**Notation.** By  $\mathbb{N} = \{0, 1, 2, \dots\}$  we denote the natural numbers, and by  $[\mathbb{N}]^\infty$  we denote the set of all infinite subsets of  $\mathbb{N}$  (which is clearly a Polish subspace of  $2^\mathbb{N}$ ). By  $2^{<\mathbb{N}}$  we denote the set of all finite sequences of 0's and 1's (the empty sequence is included). We view  $2^{<\mathbb{N}}$  as a tree equipped with the (strict) partial order  $\sqsubset$  of extension. For every  $t \in 2^{<\mathbb{N}}$  by  $|t|$  we denote the length of  $t$ , that is, the cardinality of the set  $\{s \in 2^{<\mathbb{N}} : s \sqsubset t\}$ . For every  $n \in \mathbb{N}$  set  $2^n := \{t \in 2^{<\mathbb{N}} : |t| = n\}$ . If  $s, t \in 2^{<\mathbb{N}}$ , then by  $s^\frown t$  we denote their concatenation. For every  $\sigma \in 2^\mathbb{N}$  and every  $n \geq 1$  let  $\sigma|n = (\sigma(0), \dots, \sigma(n-1))$ , while  $\sigma|0 = \emptyset$ .

If  $(x_n)$  and  $(y_n)$  are basic sequences in a Banach space  $X$  and  $C \geq 1$ , then we say that  $(x_n)$  is  $C$ -dominated by  $(y_n)$  (or simply *dominated*, if  $C$  is understood) if for every  $k \in \mathbb{N}$  and every  $a_0, \dots, a_k \in \mathbb{R}$  we have

$$\left\| \sum_{n=0}^k a_n x_n \right\| \leq C \left\| \sum_{n=0}^k a_n y_n \right\|.$$

We write  $(x_n) \leq (y_n)$  to denote the fact that  $(x_n)$  is dominated by  $(y_n)$ . All the other pieces of notation we use are standard (see, e.g., [Ke, LT, AOST]).

## 2. QUASI-ORDERS AND BOREL ORDERS

A *quasi-order* is a set  $X$  with a binary relation  $\leq$  on  $X$  which is reflexive and transitive. Given  $x, y \in X$  we set

- (a)  $x \equiv y \iff (x \leq y) \text{ and } (y \leq x)$ ,
- (b)  $x < y \iff (x \leq y) \text{ and } (y \not\leq x)$ ,
- (c)  $x \perp y \iff (x \not\leq y) \text{ and } (y \not\leq x)$ .

If  $x, y \in X$  are as in case (c) above, then we say that  $x$  and  $y$  are *incomparable*. An *antichain* is a subset of  $X$  consisting of pairwise incomparable elements. An  $\omega_1$ -chain in  $X$  is a sequence  $(x_\xi)_{\xi < \omega_1}$  in  $X$  such that either  $x_\xi < x_\zeta$  for every  $\xi < \zeta < \omega_1$ , or  $x_\xi < x_\zeta$  for every  $\zeta < \xi < \omega_1$ .

A *Borel order* is a quasi-order  $(X, \leq)$  where  $X$  is Polish and  $\leq$  is Borel in  $X^2$ . A Borel order is called *thin* if  $X$  does not contain a perfect set of pairwise incomparable elements. We will need the following lemma concerning the structure of  $F_\sigma$  thin orders.

**Lemma 4.** *Let  $X$  be a Polish space and let  $\leq$  be an  $F_\sigma$  thin order on  $X$ . Then  $(X, \leq)$  does not contain  $\omega_1$ -chains.*

Lemma 4 is a very special case of a deep result due to Harrington and Shelah (see [HS, HMS]) asserting that *any* Borel thin order does not contain  $\omega_1$ -chains. We notice that, prior to [HS], Friedman had shown ([F]) that any Borel linear order does not contain  $\omega_1$ -chains.

*Proof of Lemma 4.* Let  $(F_n)$  be an increasing sequence of closed subsets of  $X^2$  with  $\leq = \bigcup_n F_n$ . By symmetry, it is enough to show that if  $(X, \leq)$  contains a strictly increasing sequence  $(x_\xi)_{\xi < \omega_1}$ , then there exists a perfect subset  $P$  of  $X$  such that  $x \perp y$  for every  $x, y \in P$  with  $x \neq y$ . Set  $\Gamma := \{x_\xi : \xi < \omega_1\}$ . Refining if necessary, we may assume that for every  $\xi < \omega_1$  the point  $x_\xi$  is a condensation point of  $\Gamma$ . Let  $\rho$  be a compatible complete metric for  $X$ . By recursion on the length of sequences in  $2^{<\mathbb{N}}$ , we shall construct a family  $(U_t)_{t \in 2^{<\mathbb{N}}}$  of open subsets of  $X$  such that the following are satisfied.

- (a) For every  $t \in 2^{<\mathbb{N}}$  we have  $\overline{U_{t \smallfrown 0}}, \overline{U_{t \smallfrown 1}} \subseteq U_t$  and  $\overline{U_{t \smallfrown 0}} \cap \overline{U_{t \smallfrown 1}} = \emptyset$ .
- (b) For every  $t \in 2^{<\mathbb{N}}$  with  $|t| \geq 1$  we have  $\rho - \text{diam}(U_t) \leq \frac{1}{|t|}$ .
- (c) For every  $n \geq 1$  and every  $t, s \in 2^n$  with  $t \neq s$  we have  $(U_t \times U_s) \cap F_n = \emptyset$  and  $(U_s \times U_t) \cap F_n = \emptyset$ .
- (d) For every  $t \in 2^{<\mathbb{N}}$  we have  $U_t \cap \Gamma \neq \emptyset$ .

Assuming that the construction has been carried out, we set

$$P := \bigcup_{\sigma \in 2^{\mathbb{N}}} \bigcap_{n \in \mathbb{N}} U_{\sigma \restriction n}.$$

By (a) and (b) above, we see that  $P$  is a perfect subset of  $X$ . Moreover, using (c), it is easy to check that  $P$  is in addition an antichain.

We proceed to the construction. For  $n = 0$  set  $U_\emptyset := X$ . Let  $\xi < \zeta < \omega_1$ . Then  $x_\xi < x_\zeta$ , and so,  $x_\zeta \not\perp x_\xi$ . In particular, we have that  $(x_\zeta, x_\xi) \notin F_1$ . Hence, there exist  $V^0, W^0$  open subsets of  $X$  such that  $x_\zeta \in V^0$ ,  $x_\xi \in W^0$  and  $(V^0 \times W^0) \cap F_1 = \emptyset$ . Notice that both  $V^0 \cap \Gamma$  and  $W^0 \cap \Gamma$  are uncountable. So, we may select  $\eta < \theta < \omega_1$  such that  $x_\eta \in V^0$  and  $x_\theta \in W^0$ . Since  $x_\theta \not\perp x_\eta$ , we find  $V^1, W^1$  open subsets of  $V^0$  and  $W^0$  respectively such that  $x_\theta \in W^1$ ,  $x_\eta \in V^1$  and  $(W^1 \times V^1) \cap F_1 = \emptyset$ . Notice that conditions (c) and (d) above are satisfied for  $V^1$  and  $W^1$  except, possibly, (a) and (b). Thus, refining, we find  $U_{(0)}$  and  $U_{(1)}$  open subsets of  $V^1$  and  $W^1$  respectively such that conditions (a)–(d) are satisfied. For the general step we proceed similarly. The proof is completed.  $\square$

For more information on the structure of Borel thin orders we refer to the work of Louveau [L], and Louveau and Saint Raymond [LStR]. For applications of the theory of Borel orders to Banach space theory we refer to the work of Rosendal [Ros].

We will also need the following special case of the theorem of Silver [Si] on the number of equivalence classes of co-analytic equivalence relations. The proof given

below is an adaptation of Louveau's approach on Silver's theorem (via the, so called, Gandy–Harrington topology—see [MK]) in an easier setting.

**Lemma 5.** *Let  $X$  be a Polish space and let  $\sim$  be an  $F_\sigma$  equivalence relation on  $X$ . Then, either the equivalence classes of  $\sim$  are countable, or there exists a Cantor set  $P \subseteq X$  consisting of pairwise inequivalent elements.*

*Proof.* Let  $\mathcal{B} = (U_n)$  be a countable basis of  $X$ . For every closed subset  $F$  of  $X$  set

$$D(F) := F \setminus \bigcup \{U_n \in \mathcal{B} : \exists x \in F \text{ with } U_n \cap F \subseteq [x]\}$$

where  $[x] = \{y \in X : x \sim y\}$ . That is,  $D(F)$  results by removing from  $F$  all basic relatively open subsets of  $F$  which are contained in a single equivalence class. Clearly  $D(F)$  is closed and  $D(F) \subseteq F$ . By transfinite recursion, we define a decreasing sequence  $(X_\xi)_{\xi < \omega_1}$  of closed subsets of  $X$  as follows. We set  $X_0 = X$ ,  $X_{\xi+1} = D(X_\xi)$  and  $X_\lambda = \bigcap_{\xi < \lambda} X_\xi$  if  $\lambda$  is limit. There exists  $\xi_0 < \omega_1$  such that  $X_{\xi_0} = X_{\xi_0+1}$ . We consider the following cases.

CASE 1:  $X_{\xi_0} = \emptyset$ . Notice that for every  $\xi < \xi_0$  the set  $X_\xi \setminus X_{\xi+1}$  is contained in at most countable many equivalence classes. Since  $X_{\xi_0} = \emptyset$ , we see that

$$X = \bigcup_{\xi < \xi_0} X_\xi \setminus X_{\xi+1}.$$

Hence, this case implies that the equivalence classes of  $\sim$  are countable.

CASE 2:  $X_{\xi_0} \neq \emptyset$ . We set  $Y = X_{\xi_0}$  and  $\sim' = \sim \cap Y^2$ . Clearly  $\sim'$  is  $F_\sigma$  in  $Y^2$ . We claim that  $\sim'$  is meager in  $Y^2$ . Indeed, by the Kuratowski–Ulam theorem (see, e.g., [Ke, Theorem 8.41]), it is enough to show that for every  $x \in Y$  the set  $[x]' = \{y \in Y : x \sim' y\} = \{y \in Y : x \sim y\}$  is meager. Notice that  $[x]'$  is  $F_\sigma$  in  $Y$ . Therefore, if  $[x]'$  was not meager, then there would exist  $U_n \in \mathcal{B}$  such that  $U_n \cap Y \subseteq [x]'$ . This implies that  $D(X_{\xi_0}) \subsetneq X_{\xi_0}$ , a contradiction. Thus,  $\sim'$  is meager in  $Y^2$ , as claimed. It follows, by a classical result of Mycielski (see, e.g., [Ke, Theorem 19.1]), that there exists a Cantor set  $P \subseteq Y$  such that  $x \not\sim' y$  for every  $x, y \in P$  with  $x \neq y$ . This clearly implies that  $x \not\sim y$  for every  $x, y \in P$  with  $x \neq y$ . The proof is completed.  $\square$

### 3. CODING $(\text{SP}_w(X), \leq)$ AS AN $F_\sigma$ ORDER

Let  $X$  be a separable Banach space. Our aim is to show that the quasi-order  $(\text{SP}_w(X), \leq)$  can be realized as an  $F_\sigma$  order. This is done in a rather standard and natural way.

Let  $U$  be the universal space of Pelczynski for unconditional basic sequences (see [P]). That is, the space  $U$  has an unconditional Schauder basis  $(u_n)$  and for any other unconditional basic sequence  $(y_n)$  in some Banach space  $Y$  there exists  $L = \{l_0 < l_1 < \dots\} \in [\mathbb{N}]^\infty$  such that  $(y_n)$  is equivalent to  $(u_{l_n})$ . In what follows,

for every  $L = \{l_0 < l_1 < \dots\} \in [\mathbb{N}]^\infty$  by  $(u_n)_{n \in L}$  we denote the subsequence  $(u_{l_n})$  of  $(u_n)$  determined by  $L$ . Define  $\leq$  in  $[\mathbb{N}]^\infty \times [\mathbb{N}]^\infty$  by the rule

$$L \leq M \Leftrightarrow (u_n)_{n \in L} \text{ is dominated by } (u_n)_{n \in M}.$$

Clearly  $\leq$  is a quasi-order. Let  $\sim$  be the associated equivalence relation (that is,  $L \sim M$  if and only if  $L \leq M$  and  $M \leq L$ ) and let  $<$  be the strict part of  $\leq$  (that is,  $L < M$  if and only if  $L \leq M$  and  $M \not\leq L$ ). Notice that  $L \sim M$  if and only if the sequences  $(u_n)_{n \in L}$  and  $(u_n)_{n \in M}$  are equivalent as basic sequences. We have the following easy fact whose proof is sketched for completeness.

**Fact 6.** *Both  $\leq$  and  $\sim$  are  $F_\sigma$ .*

*Proof.* It is enough to show that  $\leq$  is  $F_\sigma$ . For every  $K \in \mathbb{N}$  with  $K \geq 1$  let  $\leq_K$  be the relation on  $[\mathbb{N}]^\infty \times [\mathbb{N}]^\infty$  defined by setting

$$L \leq_K M \Leftrightarrow (u_n)_{n \in L} \text{ is } K\text{-dominated by } (u_n)_{n \in M}.$$

It is easy to see that  $\leq_K$  is closed in  $[\mathbb{N}]^\infty \times [\mathbb{N}]^\infty$ . Since  $\leq$  is the union of  $\leq_K$  over all  $K \geq 1$ , the result follows.  $\square$

Our coding of  $(\text{SP}_w(X), \leq)$  as an  $F_\sigma$  order will be done via the following lemma.

**Lemma 7.** *Let  $X$  be a separable Banach space. Then there exists  $A_X \subseteq [\mathbb{N}]^\infty$  analytic such that the following are satisfied.*

- (1) *For every  $(y_n) \in \text{SP}_w(X)$  there exists  $L \in A_X$  such that  $(y_n)$  is equivalent to  $(u_n)_{n \in L}$ .*
- (2) *For every  $L \in A_X$  there exists  $(y_n) \in \text{SP}_w(X)$  such that  $(u_n)_{n \in L}$  is equivalent to  $(y_n)$ .*

*Proof.* Recall that a sequence  $(x_n)$  in  $X$  is said to be *Cesaro summable* if

$$\lim_{n} \frac{x_0 + \dots + x_{n-1}}{n} = 0.$$

Let  $\text{SPC}$  be the subset of  $X^\mathbb{N}$  defined by

$$(x_n) \in \text{SPC} \Leftrightarrow (x_n) \text{ is seminormalized, basic, Cesaro summable} \\ \text{and } C\text{-Schreier spreading for some } C \geq 1.$$

It is easy to check that  $\text{SPC}$  is a Borel subset of  $X^\mathbb{N}$  (actually, it is  $F_{\sigma\delta}$ ). Consider the subset  $A$  of  $[\mathbb{N}]^\infty$  defined by

$$L \in A \Leftrightarrow \text{if } L = \{l_0 < l_1 < \dots\}, \text{ then } \exists (x_n) \in X^\mathbb{N} \exists \theta \geq 1 \text{ with } \left[ (x_n) \in \text{SPC} \right. \\ \left. \text{and } (\forall k \forall \ell \leq n_0 < \dots < n_k \text{ we have } (x_{n_i})_{i=0}^k \overset{\theta}{\sim} (u_{l_i})_{i=0}^k \right].$$

As  $\text{SPC}$  is Borel in  $X^\mathbb{N}$ , it is easy to see that the set  $A$  is analytic. Denote by  $(e_n)$  the standard basis of  $\ell_1$ . Let us isolate the following property of the set  $A$ .

- (P) If  $L \in A$ , then the sequence  $(u_n)_{n \in L}$  is not equivalent to  $(e_n)$ . This follows from the fact that every sequence  $(x_n)$  belonging to SPC is a Cesaro summable Schauder basic sequence.

The proof of the lemma will be finished once we show the following claim.

**Claim.** *Let  $(y_n) \in \text{SP}_w(X)$  which is not equivalent to  $(e_n)$ . Then there exists  $L \in A$  such that  $(y_n)$  is equivalent to  $(u_n)_{n \in L}$ . Conversely, for every  $L \in A$  there exists  $(y_n) \in \text{SP}_w(X)$  which is not equivalent to  $(e_n)$  and such that  $(u_n)_{n \in L}$  is equivalent to  $(y_n)$ .*

*Proof of the claim.* Let  $(y_n) \in \text{SP}_w(X)$  not equivalent to  $(e_n)$  and let  $(x_n)$  be a seminormalized weakly-null sequence in  $X$  that generates it. By passing to a subsequence, we may assume that  $(x_n)$  is a seminormalized,  $C$ -Schreier spreading (for some  $C \geq 1$ ) basic sequence. Since  $(y_n)$  is not equivalent to  $(e_n)$ , by a result of Rosenthal, we see that  $(x_n)$  has a subsequence  $(x_{n_k})$  which is additionally Cesaro summable (see [AT, Theorem II.9.8]). Therefore,  $(x_{n_k}) \in \text{SPC}$ . Since  $(x_{n_k})$  still generates  $(y_n)$  as spreading model, we conclude that there exists  $L \in A$  such that  $(u_n)_{n \in L}$  is equivalent to  $(y_n)$ .

Conversely, let  $L \in A$ . We select  $(x_n) \in \text{SPC}$  witnessing that  $L \in A$ . By property (P) above, we see that  $(u_n)_{n \in L}$  is not equivalent to  $(e_n)$ . We claim that  $(x_n)$  is weakly-null. Assume not. Then there exist  $M = \{m_0 < m_1 < \dots\} \in [\mathbb{N}]^\infty$ ,  $x^* \in X^*$  and  $\varepsilon > 0$  such that  $x^*(x_{m_n}) > \varepsilon$  for every  $n \in \mathbb{N}$  (also notice that  $m_n \geq n$ ). Let  $K \geq 1$  be the basis constant of  $(x_n)$ . Also let  $C \geq 1$  be such that  $(x_n)$  is  $C$ -Schreier spreading. Observe that for every  $n \in \mathbb{N}$  we have

$$\begin{aligned} \left\| \frac{x_0 + \dots + x_{2n-1}}{2n} \right\| &\geq \frac{1}{2(K+1)} \left\| \frac{x_n + \dots + x_{2n-1}}{n} \right\| \\ &\geq \frac{1}{2C(K+1)} \left\| \frac{x_{m_n} + \dots + x_{m_{2n-1}}}{n} \right\| \geq \frac{\varepsilon}{2C(K+1)} \end{aligned}$$

which implies that  $(x_n)$  is not Cesaro summable, a contradiction. Thus,  $(x_n)$  is weakly-null. Let  $(y_n)$  be a spreading model generated by a subsequence of  $(x_n)$ . Then  $(y_n) \in \text{SP}_w(X)$ . Invoking the definition of the set  $A$  again, we see that  $(y_n)$  is equivalent to  $(u_n)_{n \in L}$ . This yields additionally that  $(y_n)$  is not equivalent to  $(e_n)$ . The proof of the claim is completed.  $\square$

If  $(e_n) \notin \text{SP}_w(X)$ , then we set  $A_X := A$ . If  $(e_n) \in \text{SP}_w(X)$ , then we set

$$A_X := A \cup \{L \in [\mathbb{N}]^\infty : (u_n)_{n \in L} \sim (e_n)\}.$$

Clearly  $A_X$  is analytic and, by the above claim,  $A_X$  is as desired.  $\square$

#### 4. PROOF OF THEOREM 1

Let  $X$  be a separable Banach space such that  $\text{SP}_w(X)$  is uncountable. Let  $A_X$  be the analytic subset of  $[\mathbb{N}]^\infty$  obtained by Lemma 7. We fix a continuous map

$\Phi: \mathbb{N}^{\mathbb{N}} \rightarrow [\mathbb{N}]^{\infty}$  with  $\Phi(\mathbb{N}^{\mathbb{N}}) = A_X$ . We define  $\preceq$  on  $\mathbb{N}^{\mathbb{N}}$  by the rule

$$\alpha \preceq \beta \Leftrightarrow \Phi(\alpha) \leq \Phi(\beta).$$

By Fact 6 and the continuity of  $\Phi$ , we see that  $\preceq$  is an  $F_{\sigma}$  quasi-order on the Baire space  $\mathbb{N}^{\mathbb{N}}$ . We have the following lemma.

**Lemma 8.** *Let  $X$  be a separable Banach space such that  $\text{SP}_w(X)$  is uncountable, and consider the  $F_{\sigma}$  quasi-order  $(\mathbb{N}^{\mathbb{N}}, \preceq)$ . Then, either*

- (a)  $(\mathbb{N}^{\mathbb{N}}, \preceq)$  is not thin, or
- (b)  $(\mathbb{N}^{\mathbb{N}}, \preceq)$  contains a strictly increasing sequence of length  $\omega_1$ .

*Proof.* Let  $\cong$  be the equivalence relation associated with  $\preceq$  (that is,  $\alpha \cong \beta$  if  $\alpha \preceq \beta$  and  $\beta \preceq \alpha$ ). Notice that

$$\alpha \cong \beta \Leftrightarrow \Phi(\alpha) \sim \Phi(\beta)$$

for every  $\alpha, \beta \in \mathbb{N}^{\mathbb{N}}$ . Also observe that  $\cong$  is an  $F_{\sigma}$  equivalence relation. Since  $\text{SP}_w(X)$  is uncountable, we see that  $\cong$  has uncountable many equivalence classes. Thus, by Lemma 5, there exists a Cantor set  $P \subseteq \mathbb{N}^{\mathbb{N}}$  such that  $\alpha \not\cong \beta$  for every  $\alpha, \beta \in P$  with  $\alpha \neq \beta$ . Fix a homeomorphism  $h: 2^{\mathbb{N}} \rightarrow P$ . Let  $<_{\text{lex}}$  be the (strict) lexicographical ordering on  $2^{\mathbb{N}}$ . For every  $Q \subseteq 2^{\mathbb{N}}$  denote by  $[Q]^2$  the set of unordered pairs of elements of  $Q$ . Consider the following subsets  $\mathcal{I}$  and  $\mathcal{D}$  of  $[2^{\mathbb{N}}]^2$  defined by

$$\{\sigma, \tau\} \in \mathcal{I} \Leftrightarrow \text{if } \sigma <_{\text{lex}} \tau \text{ then } h(\sigma) \preceq h(\tau),$$

$$\{\sigma, \tau\} \in \mathcal{D} \Leftrightarrow \text{if } \sigma <_{\text{lex}} \tau \text{ then } h(\tau) \preceq h(\sigma).$$

It is easy to check that both  $\mathcal{I}$  and  $\mathcal{D}$  are Borel in  $[2^{\mathbb{N}}]^2$ , in the sense that the sets

$$\mathcal{I}^* = \{(\sigma, \tau) \in 2^{\mathbb{N}} \times 2^{\mathbb{N}} : \{\sigma, \tau\} \in \mathcal{I}\} \text{ and } \mathcal{D}^* = \{(\sigma, \tau) \in 2^{\mathbb{N}} \times 2^{\mathbb{N}} : \{\sigma, \tau\} \in \mathcal{D}\}$$

are both Borel subsets of  $2^{\mathbb{N}} \times 2^{\mathbb{N}}$ . By result of Galvin (see [Ke, Theorem 19.7]), there exists a perfect set  $Q \subseteq 2^{\mathbb{N}}$  such that one of the following cases holds true.

CASE 1:  $[Q]^2 \subseteq \mathcal{I}$ . We fix a sequence  $(\sigma_n)$  in  $Q$  which is increasing with respect to  $<_{\text{lex}}$ . Then  $h(\sigma_n) \preceq h(\sigma_m)$  for every  $n < m$ . Since  $h(Q) \subseteq P$  and  $P$  consists of inequivalent elements with respect to  $\cong$ , we see that the sequence  $(h(\sigma_n))$  is strictly increasing. This yields that  $(\text{SP}_w(X), \leq)$  contains a strictly increasing sequence. By a result of Sari [Sa], we conclude that  $\text{SP}_w(X)$  must contain a strictly increasing sequence of length  $\omega_1$ . This clearly implies that  $(\mathbb{N}^{\mathbb{N}}, \preceq)$  contains a strictly increasing sequence of length  $\omega_1$ , that is, part (b) of the lemma is valid.

CASE 2:  $[Q]^2 \subseteq \mathcal{D}$ . Let  $(\tau_n)$  be a sequence in  $Q$  which is decreasing with respect to  $<_{\text{lex}}$ . Arguing as in Case 1 above, we see that the sequence  $(h(\tau_n))$  is strictly increasing. So, this case also implies part (b) of the lemma.

CASE 3:  $[Q]^2 \cap (\mathcal{I} \cup \mathcal{D}) = \emptyset$ . We set  $R := h(Q)$ . Clearly  $R$  is a perfect subset of  $\mathbb{N}^{\mathbb{N}}$ . It is easy to check that if  $\alpha, \beta \in R$  with  $\alpha \neq \beta$ , then  $\alpha$  and  $\beta$  are incomparable with



respect to  $\preceq$ . Hence,  $R$  is a perfect antichain of  $(\mathbb{N}^{\mathbb{N}}, \preceq)$ , that is,  $(\mathbb{N}^{\mathbb{N}}, \preceq)$  is not thin. Thus, this case implies part (a) of the lemma. The proof is completed.  $\square$

**Lemma 9.** *Let  $X$  be a separable Banach space such that  $\text{SP}_w(X)$  is uncountable. Then there exists a Cantor set  $P \subseteq A_X$  consisting of pairwise incomparable elements with respect to domination.*

*Proof.* Assume, towards a contradiction, that such a Cantor set  $P$  does not exist. This easily implies that  $(\mathbb{N}^{\mathbb{N}}, \preceq)$  is a thin quasi-order. By Lemma 8, we see that  $(\mathbb{N}^{\mathbb{N}}, \preceq)$  is an  $F_\sigma$  thin order that contains an  $\omega_1$ -chain. But this possibility is ruled out by Lemma 4. Having arrived to the desired contradiction, the lemma is proved.  $\square$

**Remark 1.** We notice that Lemmas 7 and 9 immediately yield that if  $X$  is a separable Banach space such that  $\text{SP}_w(X)$  is uncountable, then  $\text{SP}_w(X)$  must contain an antichain of the size of the continuum.

We are ready to proceed to the proof of Theorem 1.

*Proof of Theorem 1.* Let  $P \subseteq A_X$  be the Cantor set obtained by Lemma 9. By passing to a perfect subset of  $P$  if necessary, we may assume that

- (A) for every  $L \in P$  the sequence  $(u_n)_{n \in L}$  is not equivalent to the standard basis of  $\ell_1$ .

We will construct the family  $(x_t)_{t \in 2^{<\mathbb{N}}}$  by “pulling back” inside  $X$  the spreading models coded by  $P$ . To this end let  $(d_m)$  be a countable dense subset of  $X$ . Let SPC be the Borel subset of  $X^{\mathbb{N}}$  defined in the proof of Lemma 7. Consider the following subset  $G$  of  $P \times [\mathbb{N}]^\infty$  defined by the rule

$$(L, M) \in G \Leftrightarrow \text{if } L = \{l_0 < l_1 < \dots\} \text{ and } M = \{m_0 < m_1 < \dots\}, \text{ then} \\ \left[ L \in P \text{ and } (d_{m_n}) \in \text{SPC and } (\exists \theta \geq 1 \right. \\ \left. \forall k \forall n_0 < \dots < n_k \text{ we have } (d_{m_{n_i}})_{i=0}^k \overset{\theta}{\sim} (u_{l_i})_{i=0}^k \right].$$

We gather, below, some properties of the set  $G$ .

- (P1) The set  $G$  is Borel.
- (P2) For every  $(L, M) \in G$  and every  $N$  infinite subset of  $M$  if  $(y_n)$  is a spreading model generated by a subsequence of  $(d_m)_{m \in N}$ , then  $(y_n)$  is equivalent to  $(u_n)_{n \in L}$ .
- (P3) For every  $L \in P$  there exists  $M \in [\mathbb{N}]^\infty$  such that  $(L, M) \in G$ .
- (P4) For every  $(L, M) \in G$  the sequence  $(d_m)_{m \in M}$  is weakly-null.

Properties (P1) and (P2) are rather straightforward consequences of the definition of the set  $G$ . Property (P3) follows by assumption (A) above, the fact that  $P$  is a subset of  $A_X$  and a standard perturbation argument. Property (P4) has already been verified in the proof of Lemma 7.

Since  $G$  is a Borel subset of  $P \times [\mathbb{N}]^\infty$ , by (P3) above and the Yankov-von Neumann uniformization theorem (see, e.g., [Ke, Theorem 18.1]), there exists a map  $f: P \rightarrow [\mathbb{N}]^\infty$  which is measurable with respect to the  $\sigma$ -algebra generated by the analytic sets and such that  $(L, f(L)) \in G$  for every  $L \in P$ . Notice that the map  $f$  must be one-to-one. Invoking the classical fact that analytic sets have the Baire property, by [Ke, Theorem 8.38] and by passing to a perfect subset of  $P$ , we may assume that  $f$  is actually continuous. Moreover, by passing to a further perfect subset of  $P$  if necessary, we may also assume that there exist  $j_0, k_0 \in \mathbb{N}$  such that for every  $L \in P$ , the sequence  $(d_m)_{m \in f(L)}$  is  $j_0$ -Schreier spreading and satisfies  $\frac{1}{k_0} \leq \|d_m\| \leq k_0$  for every  $m \in f(L)$ .

The function  $f$  is one-to-one and continuous. Hence, identifying every element of  $[\mathbb{N}]^\infty$  with its characteristic function (that is, an element of  $2^\mathbb{N}$ ), we see that the set  $f(P)$  is a perfect subset of  $2^\mathbb{N}$ . Recall that by  $\varphi: 2^{<\mathbb{N}} \rightarrow \mathbb{N}$  we denote the canonical bijection described in the introduction. By recursion on the length of finite sequences in  $2^{<\mathbb{N}}$ , we may select a family  $(m_s)_{s \in 2^{<\mathbb{N}}}$  in  $\mathbb{N}$  with the following properties.

- (P5) For every  $s_1, s_2 \in 2^{<\mathbb{N}}$  we have  $\varphi(s_1) < \varphi(s_2)$  if and only if  $m_{s_1} < m_{s_2}$ .
- (P6) For every  $\sigma \in 2^\mathbb{N}$ , setting  $M_\sigma := \{m_{\sigma|n} : n \in \mathbb{N}\} \in [\mathbb{N}]^\infty$ , there exist a unique  $L_\sigma \in P$  such that  $M_\sigma \subseteq f(L_\sigma)$ .

We set  $x_s := d_{m_s}$  for every  $s \in 2^{<\mathbb{N}}$ . Observe that  $\frac{1}{k_0} \leq \|x_s\| \leq k_0$  for every  $s \in 2^{<\mathbb{N}}$ . Also notice that for every  $\sigma \in 2^\mathbb{N}$  the sequence  $(x_{\sigma|n})$  is  $j_0$ -Schreier spreading.

Now let  $s \in 2^{<\mathbb{N}}$  with  $|s| = k$  and  $\sigma \in 2^\mathbb{N}$  with  $\sigma|k = s$ . By properties (P4) and (P6), we see that the sequence  $(x_{\sigma|n})_{n > k}$  is weakly-null. Using this observation and the classical procedure of Mazur for constructing basic sequences (see [LT]), we may select a family  $(s_t)_{t \in 2^{<\mathbb{N}}}$  in  $2^{<\mathbb{N}}$  such that, setting  $x_t := x_{s_t}$  for every  $t \in 2^{<\mathbb{N}}$ , the following are satisfied.

- (P7) For every  $t_1, t_2 \in 2^{<\mathbb{N}}$  we have  $s_{t_1} \sqsubset s_{t_2}$  if and only if  $t_1 \sqsubset t_2$ . Moreover,  $|s_{t_1}| < |s_{t_2}|$  if and only if  $\varphi(s_1) < \varphi(s_2)$ .
- (P8) If  $(t_n)$  is the enumeration of  $2^{<\mathbb{N}}$  according to the bijection  $\varphi$ , then the sequence  $(x_{t_n})$  is basic.

It is easy to verify that the family  $(x_t)_{t \in 2^{<\mathbb{N}}}$  has all properties stated in Theorem 1. The proof is completed.  $\square$

**Remark 2.** We would like to comment on the richness of the structure  $(\text{SP}_w(X), \leq)$  when  $\text{SP}_w(X)$  is uncountable. Let  $X$  be a separable Banach space and assume that there exist  $C \geq 1$  and a family  $\{(y_n^\xi) : \xi < \omega_1\}$  of mutually inequivalent spreading models generated by weakly-null sequences in  $X$  such that for every  $\xi < \zeta < \omega_1$  either the sequence  $(y_n^\xi)$  is  $C$ -dominated by  $(y_n^\zeta)$  or vice versa. By Lemma 7, there exist  $K \geq 1$  and an uncountable subset  $U$  of  $A_X$  such that the following hold. For every  $L, M \in U$  either  $(u_n)_{n \in L}$  is  $K$ -dominated by  $(u_n)_{n \in M}$  or vice versa, and moreover, for every  $L \in U$  there exists a unique ordinal  $\xi_L < \omega_1$  such that  $(u_n)_{n \in L}$

is equivalent to  $(y_n^{\xi_L})$ . Let  $\bar{U}$  be the closure of  $U$  in  $[\mathbb{N}]^\infty$  and set  $F := \bar{U} \cap A_X$ . Then  $F$  is an uncountable analytic set. Consider the following symmetric relation  $\approx_K$  in  $[\mathbb{N}]^\infty \times [\mathbb{N}]^\infty$  defined by

$$L \approx_K M \Leftrightarrow \text{either } (u_n)_{n \in L} \text{ is } K\text{-dominated by } (u_n)_{n \in M} \text{ or vice versa.}$$

It is easy to see that  $\approx_K$  is closed in  $[\mathbb{N}]^\infty \times [\mathbb{N}]^\infty$ . By the choice of  $U$ , we have  $L \approx_K M$  for every  $L, M \in U$ . Since  $\approx_K$  is closed, we see that  $L \approx_K M$  for every  $L, M \in \bar{U}$ . In particular,  $L \approx_K M$  for every  $L, M \in F$ . Notice that  $U \subseteq F$ , and so, the relation  $\sim$  of equivalence restricted on  $F$  has uncountable many equivalence classes. By Lemma 5, there exists a perfect subset  $P$  of  $F$  such that for every  $L, M \in P$  the sequences  $(u_n)_{n \in L}$  and  $(u_n)_{n \in M}$  are not equivalent<sup>2</sup>. Thus, we have shown the following proposition.

**Proposition 10.** *Let  $X$  be a separable Banach space and assume that there exist  $C \geq 1$  and a family  $\{(y_n^\xi) : \xi < \omega_1\}$  of mutually inequivalent spreading models generated by weakly-null sequences in  $X$  such that for every  $\xi < \zeta < \omega_1$  either the sequence  $(y_n^\xi)$  is  $C$ -dominated by  $(y_n^\zeta)$  or vice versa. Then  $(\text{SP}_w(X), \leq)$  contains a linearly ordered subset of the size of the continuum.*

Related to Proposition 10, the following question is open to us. Let  $X$  be a separable Banach space and assume that  $\text{SP}_w(X)$  is uncountable. Does  $(\text{SP}_w(X), \leq)$  contain a linearly ordered subset of the size of the continuum, or at least uncountable?

## 5. PROOF OF THEOREM 3

(a) First we need to recall some standard facts (see [Ke, page 351]). Let  $S$  be a set and let  $\prec$  be a strict, well-founded (binary) relation on  $S$ . This is equivalent to asserting that there is no infinite decreasing chain  $\cdots \prec s_1 \prec s_0$ . By recursion on  $\prec$ , we define the *rank* function  $\rho_\prec : S \rightarrow \text{Ord}$  of  $\prec$  by the rule

$$\rho_\prec(s) = \sup\{\rho_\prec(x) + 1 : x \prec s\}.$$

In particular,  $\rho_\prec(s) = 0$  if and only if  $s$  is minimal. The *rank*  $\rho(\prec)$  of  $\prec$  is defined by  $\rho(\prec) = \sup\{\rho_\prec(s) + 1 : s \in S\}$ .

We are ready to proceed to the proof. So, let  $X$  be a separable Banach space such that  $\text{SP}_w(X)$  contains a strictly decreasing sequence of length  $\omega_1$ . Let  $A_X$  be the analytic subset of  $[\mathbb{N}]^\infty$  obtained by Lemma 7. Consider the following relation  $\prec$  on  $[\mathbb{N}]^\infty$  defined by

$$L \prec M \Leftrightarrow (L \in A_X \text{ and } (M \in A_X \text{ and } (M < L)).$$

That is,  $\prec$  is the relation  $>$  (the reverse of  $<$ ) restricted on  $A_X \times A_X$ . Clearly  $\prec$  is analytic (as a subset of  $[\mathbb{N}]^\infty \times [\mathbb{N}]^\infty$ ). Let  $\{(y_n^\xi) : \xi < \omega_1\}$  be a strictly decreasing

<sup>2</sup>This does not follow directly by Lemma 5 since  $F$  is not Polish. One has to observe that  $F$  is the continuous surjective image of  $\mathbb{N}^{\mathbb{N}}$  and use an argument as in the beginning of Section 4.

sequence in  $\text{SP}_w(X)$ . By Lemma 7, for every  $\xi < \omega_1$  we may select  $L_\xi \in A_X$  such that  $(u_n)_{n \in L_\xi}$  is equivalent to  $(y_n^\xi)$ . It follows that  $L_\xi < L_\zeta$  if and only if  $\zeta < \xi$ .

Assume, towards a contradiction, that  $\text{SP}_w(X)$  does not contain a strictly increasing sequence of length  $\omega_1$ . Then, by the result of Sari [Sa] already quoted in the proof of Theorem 1,  $\text{SP}_w(X)$  does not contain a strictly increasing sequence of length  $\omega$ . It follows that  $\prec$  is a well-founded relation on  $[\mathbb{N}]^\infty$  which is in addition analytic. By the Kunen–Martin theorem (see [Ke, Theorem 31.5]), we see that  $\rho(\prec)$  is a countable ordinal, say  $\xi_0$ . For every  $\eta < \xi_0$  set

$$A_X^\eta := \{L \in A_X : \rho_\prec(L) = \eta\}.$$

Since  $\rho_\prec(L) < \xi_0$  for every  $L \in A_X$ , we see that  $A_X = \bigcup_{\eta < \xi_0} A_X^\eta$ . Moreover, for every  $L, M \in A_X^\eta$  we have that either  $L \sim M$  or  $L \perp M$ . That is, we have partitioned the quotient  $A_X / \sim$  into countable many antichains. As the family  $\{L_\xi : \xi < \omega_1\}$  is uncountable, we see that there exist  $\xi, \zeta < \omega_1$  with  $\xi \neq \zeta$  and  $\eta < \xi_0$  such that  $L_\xi, L_\zeta \in A_X^\eta$ . But this is clearly impossible. Having arrived to the desired contradiction the proof of part (a) is completed.

(b) Again we need to discuss some standard facts. Let  $R$  be a binary relation on  $\mathbb{N}$ , that is,  $R \subseteq \mathbb{N} \times \mathbb{N}$ . By identifying  $R$  with its characteristic function, we view every binary relation on  $\mathbb{N}$  as an element of  $2^{\mathbb{N} \times \mathbb{N}}$ . Let LO be the subset of  $2^{\mathbb{N} \times \mathbb{N}}$  consisting of all (strict) linear orderings on  $\mathbb{N}$ . It is easy to see that LO is a closed subset of  $2^{\mathbb{N} \times \mathbb{N}}$  (see also [Ke, page 212]).

For every  $\alpha \in \text{LO}$  and every  $n, m \in \mathbb{N}$  we write

$$n <_\alpha m \Leftrightarrow \alpha(n, m) = 1.$$

Let WO be the subset of LO consisting of all well-orderings on  $\mathbb{N}$ . For every  $\alpha \in \text{WO}$  let  $|\alpha|$  denote the unique ordinal which is isomorphic to  $(\mathbb{N}, <_\alpha)$ . We will need the following boundedness principle for WO (see [Ke, page 240]): *if  $B$  is an analytic subset of WO, then  $\sup\{|\alpha| : \alpha \in B\} < \omega_1$ .*

We proceed to the proof of part (b). Let  $X$  be a separable Banach space. Let  $A_X$  be the analytic subset of  $[\mathbb{N}]^\infty$  obtained by Lemma 7. Consider the following subset  $O_X$  of LO defined by the rule

$$\begin{aligned} \alpha \in O_X \quad \Leftrightarrow \quad & \exists (L_n) \in ([\mathbb{N}]^\infty)^\mathbb{N} \text{ with } \left[ (\forall n \ L_n \in A_X) \text{ and} \right. \\ & \left. [\forall n, m \ (n <_\alpha m \Leftrightarrow L_n > L_m)] \right]. \end{aligned}$$

Since  $A_X$  is analytic, it easy to check that  $O_X$  is an analytic subset of LO.

**Claim.** *The set  $\text{SP}_w(X)$  does not contain a strictly increasing sequence if and only if  $O_X \subseteq \text{WO}$ .*

*Proof of the claim.* First assume that there exists  $\alpha \in O_X$  with  $\alpha \notin \text{WO}$ . By definition, there exists a sequence  $(L_n)$  in  $A_X$  such that for every  $n, m \in \mathbb{N}$  we have

$$n <_\alpha m \Leftrightarrow L_n > L_m.$$

Since  $\alpha \notin \text{WO}$ , there exists a sequence  $(n_i)$  in  $\mathbb{N}$  such that  $n_{i+1} <_\alpha n_i$  for every  $i \in \mathbb{N}$ . It follows that  $(L_{n_i})$  is a strictly increasing sequence, which clearly implies that  $\text{SP}_w(X)$  contains a strictly increasing sequence.

Conversely, assume that  $\text{SP}_w(X)$  contains a strictly increasing sequence. Hence, we may find a sequence  $(L_n)$  in  $A_X$  such that  $L_n < L_m$  if and only if  $n < m$ . Let  $\alpha \in \text{LO}$  be defined by

$$n <_\alpha m \Leftrightarrow n > m \ (\Leftrightarrow L_n > L_m).$$

Then  $\alpha \in \text{O}_X$  and  $\alpha \notin \text{WO}$ . The claim is proved.  $\square$

Now let  $X$  be a separable Banach space that does not contain a strictly increasing sequence. By the previous claim, we see that the set  $\text{O}_X$  is an analytic subset of  $\text{WO}$ . Hence, by boundedness, we see that

$$\sup\{|\alpha| : \alpha \in \text{O}_X\} = \xi_X < \omega_1.$$

We claim that  $\xi_X$  is the desired ordinal. Indeed, let  $\xi$  be a countable ordinal and let  $\{(y_n^\zeta) : \zeta < \xi\}$  be a strictly decreasing sequence in  $\text{SP}_w(X)$ . By Lemma 7, there exists  $(L_\zeta)_{\zeta < \xi}$  in  $A_X$  which is strictly decreasing. Fix a bijection  $e : \mathbb{N} \rightarrow \{\zeta : \zeta < \xi\}$  and define  $\alpha \in \text{WO}$  by setting

$$n <_\alpha m \Leftrightarrow e(n) < e(m) \ (\Leftrightarrow L_{e(n)} > L_{e(m)}).$$

It follows that  $\alpha \in \text{O}_X$ , and so,  $\xi = |\alpha| \leq \xi_X$ . The proof is completed.

**Remark 3.** Denote by  $\text{SB}$  the standard Borel space of all separable Banach spaces as it is discussed in [AD, B, Ke]. Consider the subset  $\text{NCI}$  of  $\text{SB}$  defined by

$$X \in \text{NCI} \Leftrightarrow \text{SP}_w(X) \text{ does not contain a strictly increasing infinite sequence.}$$

It can be shown, using some results from [DOS], that the set  $\text{NCI}$  is co-analytic non-Borel in  $\text{SB}$ . Moreover, there exists a co-analytic rank  $\phi : \text{NCI} \rightarrow \omega_1$  on  $\text{NCI}$  such that for every  $X \in \text{NCI}$  we have

$$\sup\{|\alpha| : \alpha \in \text{O}_X\} \leq \phi(X)$$

where  $\text{O}_X$  is as in the proof of part (b) of Theorem 3 (for the definition of co-analytic ranks we refer to [Ke], and for applications of rank theory to Banach space theory we refer to [AD]).

**Acknowledgments.** I would like to thank Spiros Argyros for many discussions on the subject as well as for his comments on the paper.

## REFERENCES

- [AOST] G. Androulakis, E. Odell, Th. Schlumprecht and N. Tomczak-Jaegermann, *On the structure of the spreading models of a Banach space*, Canadian J. Math. 57 (2005), 673–707.
- [AD] S. A. Argyros and P. Dodos, *Genericity and amalgamation of classes of Banach spaces*, Adv. Math. 209 (2007), 666–748.

- [AT] S. A. Argyros and S. Todorcević, *Ramsey Methods in Analysis*, Advanced Courses in Mathematics, CRM Barcelona, Birkhäuser, Verlag, Basel, 2005.
- [B] B. Bossard, *A coding of separable Banach spaces. Analytic and coanalytic families of Banach spaces*, Fund. Math. 172 (2002), 117–152.
- [DOS] S. J. Dilworth, E. Odell and B. Sari, *Lattice structures and spreading models*, Israel J. Math. 161 (2007), 387–411.
- [FR] V. Ferenczi and C. Rosendal, *Complexity and homogeneity in Banach spaces*, in “Banach Spaces and their Applications in Analysis”, editors B. Randrianantoanina and N. Randrianantoanina, Walter de Gruyter, Berlin, 2007.
- [F] H. Friedman, *Borel structures in mathematics*, manuscript, Ohio State University, 1979.
- [HMS] L. Harrington, D. Marker and S. Shelah, *Borel orderings*, Trans. Amer. Math. Soc. 310 (1988), 293–302.
- [HS] L. Harrington and S. Shelah, *Counting the equivalence classes of co- $\kappa$ -Suslin equivalence relations*, Logic Colloquium '80, North-Holland, Amsterdam, 1982, 147–152.
- [Ke] A. S. Kechris, *Classical Descriptive Set Theory*, Grad. Texts in Math., 156, Springer-Verlag, 1995.
- [LT] J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces I and II*, Springer, 1996.
- [L] A. Louveau, *Two results on Borel orders*, J. Symb. Logic 54 (1989), 865–874.
- [LStR] A. Louveau and J. Saint Raymond, *On the quasi-ordering of Borel linear orders under embeddability*, J. Symb. Logic 55 (1990), 537–560.
- [MK] D. A. Martin and A.S. Kechris, *Infinite games and effective descriptive set theory*, in “Analytic Sets”, editors C.A. Rogers et al., Academic Press, 1980.
- [P] A. Pelczynski, *Universal bases*, Studia Math. 19 (1960), 247–268.
- [Ros] C. Rosendal, *Incomparable, non-isomorphic and minimal Banach spaces*, Fund. Math. 183 (2004), 253–274.
- [Sa] B. Sari, *On Banach spaces with few spreading models*, Proc. Amer. Math. Soc. 134 (2005), 1339–1345.
- [Si] J. H. Silver, *Counting the number of equivalence classes of Borel and coanalytic equivalence relations*, Ann. Math. Logic 18 (1980), 1–28.

NATIONAL TECHNICAL UNIVERSITY OF ATHENS, FACULTY OF APPLIED SCIENCES, DEPARTMENT OF MATHEMATICS, ZOGRAFOU CAMPUS, 157 80, ATHENS, GREECE.

*E-mail address:* `pdodos@math.ntua.gr`