

# FPT Algorithms for Plane Completion Problems\*

Dimitris Chatzidimitriou<sup>†</sup>    Archontia C. Giannopoulou<sup>‡</sup>    Spyridon Maniatis<sup>†</sup>  
Clément Requilé<sup>§¶</sup>    Dimitrios M. Thilikos<sup>†||</sup>    Dimitris Zoros<sup>†</sup>

July 26, 2016

## Abstract

The PLANE SUBGRAPH (resp. TOPOLOGICAL MINOR) COMPLETION problem asks, given a (possibly disconnected) plane (multi)graph  $\Gamma$  and a connected plane (multi)graph  $\Delta$ , whether it is possible to add edges in  $\Gamma$  without violating the planarity of its embedding so that it contains some subgraph (resp. topological minor) that is topologically isomorphic to  $\Delta$ . We give FPT algorithms that solve both problems in  $f(|E(\Delta)|) \cdot |E(\Gamma)|^2$  steps. Moreover, for the PLANE SUBGRAPH COMPLETION problem we show that  $f(k) = 2^{\mathcal{O}(k \log k)}$ .

## 1 Introduction

*Completion problems* on graphs are defined as follows: Consider a graph class  $\mathcal{P}$  and ask whether we may add edges to a given graph  $G$  in order to obtain a graph  $G^+$ , where  $G^+ \in \mathcal{P}$ . Numerous results have appeared for the case where the objective is to minimize the number of edges added in  $G$  [16, 12, 14, 9, 3].

In this paper, we consider the PLANE SUBGRAPH (resp. TOPOLOGICAL MINOR) COMPLETION (PSC) (resp. PTMC) problem which, given a (possibly disconnected) plane graph  $\Gamma$ , called the *host graph*, and a connected plane graph  $\Delta$ , called the *pattern graph*, asks whether it is possible to add edges in  $\Gamma$  such that the resulting graph remains plane and contains some subgraph (resp. topological minor) that is topologically isomorphic to  $\Delta$ . Both  $\Gamma$  and  $\Delta$  are allowed to have multiple edges but not loops. When the input graph  $\Gamma$  is planar triangulated, both PSC and PTMC are NP-complete. Indeed, let  $G$  be any planar triangulated graph. Note here, that as any planar triangulated graph is 3-connected,  $G$  is 3-connected and from Whitney’s Theorem [15] admits a unique embedding on the plane (up to equivalence), say  $\Gamma$ . Let also  $\Delta$  be the cycle on  $n = |V(G)|$  vertices. Then  $\Delta$  also has unique embedding on the plane

---

\*The first, the third, the fifth, and the last authors were supported by the E.U. (European Social Fund - ESF) and Greek national funds through the Operational Program “Education and Lifelong Learning” of the National Strategic Reference Framework (NSRF) - Research Funding Program: “Thales. Investing in knowledge society through the European Social Fund”. The second author’s research has been supported by the European Research Council (ERC) under the European Unions Horizon 2020 research and innovation programme (ERC consolidator grant DISTRUCT, agreement No 648527) and by the Warsaw Center of Mathematics and Computer Science. The fourth author was supported by the FP7-PEOPLE-2013-CIG project CountGraph (ref. 630749), and the Berlin Mathematical School. Emails: [hatzisdimitris@gmail.com](mailto:hatzisdimitris@gmail.com), [archontia.giannopoulou@gmail.com](mailto:archontia.giannopoulou@gmail.com), [spyridon.maniatis@gmail.com](mailto:spyridon.maniatis@gmail.com), [clement.requile@gmail.com](mailto:clement.requile@gmail.com), [sedthilk@thilikos.info](mailto:sedthilk@thilikos.info), [dzoros@math.uoa.gr](mailto:dzoros@math.uoa.gr)

<sup>†</sup>Department of Mathematics, National and Kapodistrian University of Athens, Athens, Greece.

<sup>‡</sup>Institute of Software Technology and Theoretical Computer Science, Technische Universität Berlin, Germany

<sup>§</sup>Freie Universität Berlin, Institut für Mathematik und Informatik, Berlin, Germany.

<sup>¶</sup>Department of Mathematics, Universitat Politècnica de Catalunya, Barcelona, Spain

<sup>||</sup>AIGCo project team, CNRS, LIRMM, France.

(up to equivalence). Since  $\Gamma$  is triangulated no edge can be added to it while preserving its planarity. Thus, both PSC and PTMC become equivalent to the HAMILTON CYCLE PROBLEM which is NP-complete on planar triangulated graphs [5] (see also [8]). This observation further implies that PSC and PTMC parameterized by the number of added edges  $k$ , and in particular even for  $k = 0$ , are NP-complete. Thus, PSC and PTMC are not FPT when parameterized by the number of added edges unless  $\mathbf{P} = \mathbf{NP}$ . Thus, in order to obtain a tractable algorithm, we need to find an alternative way to parameterize these problems. In particular, we will consider  $|E(\Delta)|$  as our parameter. Our two main results are the following.

**Theorem.** PSC parameterized by the number of edges of the pattern graph  $\Delta$ , say  $k$ , can be solved in  $2^{\mathcal{O}(k \log k)} \cdot m^2$  time, where  $m = |E(\Gamma)|$ .

**Theorem.** PTMC parameterized by the number of edges of the pattern graph  $\Delta$ , say  $k$ , can be solved in  $f(k) \cdot m^2$  time, where  $m = |E(\Gamma)|$  and  $f$  is a computable function.

For the PTMC algorithm our approach is the following. Let  $\Gamma$  and  $\Delta$  be an input of the problem as above. We first apply a series of transformations on our input graph  $\Gamma$  that turn it into a combinatorial structure  $\mathbf{G}$  (while the topological properties of  $\Gamma$  are retained) whose treewidth is bounded by a function of  $|E(\Delta)|$ . Then, we apply a series of transformations on our input graph  $\Delta$  that allow us to encode both the topological and combinatorial information of  $\Delta$  using a combinatorial structure  $\mathbf{D}$ . Finally, we show that  $(\Delta, \Gamma)$  is a yes-instance of our problem if and only if an MSO-expressible relation holds for  $\mathbf{G}$  and  $\mathbf{D}$ , thus translating our problem into a purely combinatorial one. Then by employing Courcelle's Theorem we prove the existence of an algorithm for PTMC. We remark here that a similar approach could also solve the PLANE SUBGRAPH COMPLETION problem. However, with a more careful analysis we are able to derive an algorithm which avoids the heavy parametric dependence (caused by the use of Courcelle's theorem) for the case of plane topological minors.

Our approach towards solving PSC is the following. Let  $\Gamma$  and  $\Delta$  be an input of PSC, where  $|E(\Delta)| = k$  for some positive integer  $k$ . We construct a family  $\mathcal{G}$  consisting of  $\mathcal{O}(n)$  combinatorial structures depending only on  $\Gamma$  whose underlying graphs have treewidth  $\mathcal{O}(k)$ . We also construct a family  $\mathcal{H}$  consisting of  $2^{\mathcal{O}(k \log k)}$  combinatorial structures depending only on  $\Delta$ , again by applying a series of appropriate transformations on them (different than the transformations for PTMC). For the graphs  $\Gamma$  and  $\Delta$  and the families  $\mathcal{G}$  and  $\mathcal{H}$ , it holds that  $(\Delta, \Gamma)$  is a yes-instance if and only if some structure  $\mathbf{D} \in \mathcal{H}$  is contained as a *contraction* in a structure  $\mathbf{G} \in \mathcal{G}$ , denoted  $\mathbf{D} \leq_c \mathbf{G}$ . Therefore, we again translate our problem into one of combinatorial nature. Finally, for a fixed pair of structures  $(\mathbf{D}, \mathbf{G}) \in \mathcal{H} \times \mathcal{G}$  with the above properties, we can decide in  $2^{\mathcal{O}(k \log k)} \cdot m$  time whether  $\mathbf{D} \leq_c \mathbf{G}$ . Therefore, by testing for all pairs  $(\mathbf{D}, \mathbf{G}) \in \mathcal{H} \times \mathcal{G}$  whether  $\mathbf{D} \leq_c \mathbf{G}$ , we decide in  $2^{\mathcal{O}(k \log k)} \cdot m^2$  steps whether  $(\Delta, \Gamma)$  is a yes-instance.

The paper is organized as follows. In Section 2 we give the necessary definitions. In Section 3 we present the algorithm for the PSC problem and in Section 4 we present the algorithm for the PTMC problem. In the concluding Section 5 we discuss about other completion problems that can be solved by modifying our results, such as the PLANE INDUCED SUBGRAPH COMPLETION, the PLANE MINOR COMPLETION, the PLANAR ROOTED TOPOLOGICAL MINOR, and the PLANAR DISJOINT PATHS COMPLETION problems.

## 2 Definitions

For a positive integer  $n$ , we denote  $[n] = \{1, 2, \dots, n\}$ . Given a set  $S$ , a near-partition of  $S$  is a family of sets  $S_1, S_2, \dots, S_k$ , where  $S_i \cap S_j = \emptyset$ , for every  $i \neq j$ , and  $\cup_{i \in [k]} S_i = S$  (note that

by the definition it is possible that  $S_i = \emptyset$  for some  $i \in [k]$ ). Unless stated otherwise, the graphs considered do not have loops but may have multiple edges. Given a graph  $G$ , we will denote by  $V(G)$  the set of its vertices and  $E(G)$  the set of its edges. We denote by  $\mathbf{dist}_G(u, v)$  the distance of two vertices  $u$  and  $v$  in the graph  $G$ . Also, given a graph  $G$ , a vertex  $u \in V(G)$ , and  $V_0 \subseteq V(G)$ , we denote by  $N_G(u)$  the neighborhood of  $u$  in  $G$  and by  $N_G(V_0) := \bigcup_{v \in V_0} N_G(v) \setminus V_0$ . Given a vertex  $v$  with exactly two neighbors  $v_1$  and  $v_2$ , the dissolution of  $v$  is the operation where we delete  $v$  and add an edge  $\{v_1, v_2\}$  (even if one existed already).

Let  $G$  be a graph. A subset  $S$  of its vertices is a *separator* of  $G$  if the graph  $G - S := (V(G) \setminus S, E[V(G) \setminus S])$  is not connected. The size of a separator  $S$  is equal to  $|S|$ . The vertex contained in a separator of size 1 will be called a *cut-vertex*, while the vertices of a separator of size 2 will be called a *cut-pair*. For every integer  $k > 1$ , a graph  $G$  with at least  $k + 1$  vertices is *k-connected* if  $G$  has no separators of size less than  $k$ . For definitions not explicitly stated on the paper as well as more details on general graphs, see [7].

We say that a graph is *plane* when it is embedded without crossings between its edges on the sphere  $\Sigma = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 = 1\}$ . We treat a plane graph as its embedding in  $\Sigma$ , that is, we do not distinguish between a vertex of the graph and the point of the sphere used in the drawing to represent the vertex or between an edge and the open curve representing it. We often use the term “general graph” in order to stress that a graph is treated as a combinatorial structure and not as a topological (i.e., embedded) one. Also, given a plane graph  $\Gamma$  we use the term *general graph* of  $\Gamma$  to refer to  $\Gamma$  as a combinatorial structure. We use capital greek letters for plane graphs and capital latin letters for general graphs.

We denote by  $\subseteq$ ,  $\subseteq_{sp}$ ,  $\subseteq_{in}$ ,  $\leq_m$ , and  $\simeq$  the usual subgraph, spanning subgraph, induced subgraph, minor, and isomorphism relation between two graphs, respectively. Given a graph  $G$  and  $V_0 \subseteq V(G)$ , we denote by  $G[V_0]$  the subgraph of  $G$  induced by  $V_0$ . We call  $V_0$  *connected* if  $G[V_0]$  is connected.

Let  $\Gamma$  be a plane graph and  $u \in V(\Gamma)$ . Then a tuple  $(u_1, \dots, u_k)$  (with possible repetitions) will be called a *cyclic neighborhood* of  $u$ , and will be denoted by  $\mathcal{N}_\Gamma(u)$ , if  $\{u, u_1\}, \dots, \{u, u_k\}$  are exactly the edges incident to  $u$ , as we meet them starting from  $(u, u_1)$  and proceeding clockwise.

Let  $A$  be a subset of  $\mathbb{R}^n$ . We define  $\mathbf{int}(A)$  to be the interior of  $A$ ,  $\mathbf{cl}(A)$  its closure and  $\mathbf{bd}(A) = \mathbf{cl}(A) \setminus \mathbf{int}(A)$  its border. Given a plane graph  $\Gamma$  we denote its *faces* by  $F(\Gamma)$ , i.e.,  $F(\Gamma)$  is the set of the connected components of  $\Sigma \setminus \Gamma$  (in the operation  $\Sigma \setminus \Gamma$  we treat  $\Gamma$  as the set of points of  $\Sigma$  corresponding to its vertices and its edges). Given a graph  $G$  we denote by  $\mathcal{C}(G)$  the set of the connected components of  $G$ . For every  $f \in F(\Gamma)$  we denote by  $B_\Gamma(f)$  the graph induced by the vertices and edges of  $\Gamma$  whose embeddings are subsets of  $\mathbf{bd}(f)$  and we call it the *boundary* of  $f$ . We also denote by  $V(B_\Gamma(f))$  and  $E(B_\Gamma(f))$  the vertices and the edges of  $B_\Gamma(f)$ , respectively.

We define a *closed walk* of a graph  $G$  to be a cyclic ordering  $w = (v_1, \dots, v_l, v_1)$  of vertices of  $V(G)$  such that for any two consecutive vertices, say  $v_i, v_{i+1}$ , there is an edge between them in  $G$ , i.e.,  $\{v_i, v_{i+1}\} \in E(G)$ . Note here that there may exist two distinct indices  $i, j$  such that  $v_i, v_j \in w$  and  $v_i = v_j$  (the walk can revisit a vertex). We will denote by  $\ell_w = l$  the length of the respective closed walk  $w$ . We say that a walk  $w$  of a plane graph  $\Gamma$  is *facial* if there exist  $f_i \in F(\Gamma)$  and  $\Theta_j \in \mathcal{C}(B_\Gamma(f_i))$  such that the vertices of  $w$  are the vertices of  $V(\Theta_j)$  and the cyclic ordering of  $w$  indicates the way these vertices are met when making a closed walk along  $\Theta_j$  while always keeping  $f_i$  on the same side of the walk. Notice that a facial walk is unique (up to cyclic permutation).

Given that  $\Gamma$  is a plane graph and  $\mathbf{w} = \{w_1, \dots, w_p\}$  is a non-empty set of closed walks of  $\Gamma$ , we say that  $\mathbf{w}$  is a *facial mapping* if there exists some face  $f$  of  $\Gamma$  such that  $\mathcal{C}(B_\Gamma(f)) = \{\Theta_1, \Theta_2, \dots, \Theta_p\}$  and  $w_j$  is a facial walk of  $\Theta_j$ ,  $j \in [p]$ . We define the length of the facial mapping  $\mathbf{w}$  to be  $\ell_{\mathbf{w}} = \sum_{i=1}^p \ell_{w_i}$ . Given a plane graph  $\Gamma$  and  $f \in F(\Gamma)$ , we define  $\mathbf{w}(f)$  as the

facial mapping of  $\Gamma$  corresponding to  $f$  and define its length  $\ell_f$  to be the length  $\ell_{\mathbf{w}}(f)$  of its corresponding facial mapping. Observe that for every face  $f \in F(\Gamma)$ , its facial mapping  $\mathbf{w}(f)$  is unique (up to permutations). Let  $C_1, C_2$  be two disjoint closed curves of  $\Sigma$ . Let also  $D_i$  be the open disk of  $\Sigma \setminus C_i$  that does not contain points of  $C_{3-i}$ ,  $i \in [2]$ . The *annulus between  $C_1$  and  $C_2$*  is the set  $\Sigma \setminus (D_1 \cup D_2)$  and we denote it by  $A[C_1, C_2]$ . Notice that  $A[C_1, C_2]$  is a closed set.

Let  $\Gamma$  and  $\Delta$  be two plane graphs. We say that  $\Gamma$  and  $\Delta$  are *topologically isomorphic* if they are isomorphic via a bijection  $g : V(\Gamma) \rightarrow V(\Delta)$  and there exists a function  $h : F(\Gamma) \rightarrow F(\Delta)$ , such that for every  $f \in F(\Gamma)$ ,  $g(\mathbf{w}(f)) = \mathbf{w}(h(f))$  (where  $g(\mathbf{w}(f))$  is the result of applying  $g$  to every vertex of every closed walk in  $\mathbf{w}$ ). We call the function  $\alpha : V(\Gamma) \cup F(\Gamma) \rightarrow V(\Delta) \cup F(\Delta)$  such that  $\alpha = g \cup h$ , a *topological isomorphism* between  $\Gamma$  and  $\Delta$ .

We say that a general graph  $G$  is *uniquely embeddable* if any two plane graphs  $\Gamma$  and  $\Gamma'$  that are embeddings of  $G$  in the sphere, are topologically isomorphic. We say that a plane graph  $\Gamma$  is *uniquely embedded* if its general graph  $G$  is uniquely embeddable, i.e.,  $\Gamma$  is the unique embedding of  $G$ , up to topological isomorphism. Given two plane graphs  $\Gamma_1$  and  $\Gamma_2$  we say that they are the *same graph* if they are topologically isomorphic (and not just isomorphic).

Let  $\Gamma$  and  $\Delta$  be two plane graphs and let  $Z \subseteq V(\Gamma)$ . We say that  $\Delta$  is a  *$Z$ -embedded subgraph* of  $\Gamma$ , and write  $\Delta \leq_{es}^Z \Gamma$ , if  $\Delta$  is topologically isomorphic to some subgraph of  $\Gamma \setminus Z$ . When  $Z = \emptyset$ , we say that  $\Delta$  is an *embedded subgraph* of  $\Gamma$  and write  $\Delta \leq_{es} \Gamma$ .

**Definition 1 ( *$Z$ -embedded topological minors*).** *Let  $\Gamma$  and  $\Delta$  be two plane graphs and let  $Z \subseteq V(\Gamma)$ . We say that  $\Delta$  is a  $Z$ -embedded topological minor of  $\Gamma$ , and write  $\Delta \leq_{etm}^Z \Gamma$  if there exist a function  $\rho_1 : V(\Delta) \rightarrow V(\Gamma)$  and a function  $\rho_2 : E(\Delta) \rightarrow \mathcal{P}(\Gamma)$ , where  $\mathcal{P}(\Gamma)$  denotes the set of all paths of  $\Gamma$ , such that*

1. *For every  $v \in V(\Delta)$ ,  $\rho_1(v) \notin Z$ .*
2. *For every  $e = \{u, v\} \in E(\Delta)$ , the path  $\rho_2(e)$  of  $\Gamma$  has  $\rho_1(u)$  and  $\rho_1(v)$  as its endpoints and if  $e_1 \neq e_2$ , then  $\rho_2(e_1)$  and  $\rho_2(e_2)$  are internally vertex-disjoint.*
3. *If  $\Gamma\langle\rho_2\rangle$  is the graph obtained by the union of all paths in  $\rho_2(E(\Delta))$  after we dissolve all vertices that are not vertices in  $\rho_1(V(\Delta))$ , then there is a topological isomorphism  $\alpha : V(\Delta) \cup F(\Delta) \rightarrow V(\Gamma\langle\rho_2\rangle) \cup F(\Gamma\langle\rho_2\rangle)$  between  $\Delta$  and  $\Gamma\langle\rho_2\rangle$  where  $\alpha|_{V(\Delta)} = \rho_1$ .*

When  $Z = \emptyset$ , we just write  $\Delta \leq_{etm} \Gamma$ .

If in the 3rd condition of the above definition we replace topological isomorphism by isomorphism and consider general graphs, say  $H$  and  $G$ , we define the relation of  $H$  being a  *$Z$ -topological minor* of  $(G, Z)$ .

For definitions regarding plane graphs not explicitly stated on the paper as well as more details on the subject, see [13].

## 2.1 Radial Enhancements

Let  $\Gamma$  be a plane graph. A *subdivided radial enhancement* of  $\Gamma$  is defined as a plane graph that can be constructed as follows: consider  $\Gamma$ , subdivide every edge once, add a vertex  $v_f$  inside each face  $f$  of  $\Gamma$ . Consider a permutation  $(H_1, H_2, \dots, H_s)$  of the connected components of  $B_\Gamma(f)$  and a facial walk of each connected component. Then add edges connecting  $v_f$  with the vertices incident to  $B_\Gamma(f)$  in such a way that the first vertices in the cyclic neighborhood of  $v_f$  are the vertices of  $H_1$  and appear in the order of the fixed facial walk. Then the vertices of  $H_2$  follow, etc. Observe that in the resulting embedding, every face that is incident to an edge of  $E(\Gamma)$  is (planar) triangulated. This triangulation may have multiple edges unless the boundary of each face of  $\Gamma$  is a cycle, as can be seen in the two distinct examples of a subdivided radial enhancement of a disconnected plane graph  $\Gamma$  in Figure 1.




Figure 1: A disconnected plane graph  $\Gamma$  and two members of  $\mathcal{R}_\Gamma$ .

Notice that the vertices of the resulting plane graph can be partitioned into three independent sets: the *original vertices* of  $\Gamma$  denoted by  $V_o(\Gamma)$ , the *subdivision vertices* denoted by  $V_s(\Gamma)$ , which are the ones that were introduced after subdividing the edges, and the *radial vertices* denoted by  $V_r(\Gamma)$ , which are the ones that were added inside each face. Notice also that the edges of the resulting plane graph can be partitioned into two independent sets: the *subdivision edges* denoted by  $E_s(\Gamma)$  and the *radial edges*, denoted by  $E_r(\Gamma)$ , that were introduced after adding the radial vertices.

We denote by  $\mathcal{R}_\Gamma$  the set of all different (in terms of topological isomorphism) subdivided radial enhancements of  $\Gamma$ . Observe that if  $\Gamma$  is connected, then the boundary of each face of  $\Gamma$  is connected and we obtain the following.

**Observation 1.** *For every connected plane graph  $\Gamma$ ,  $R(\Gamma)$  is uniquely defined and thus  $\mathcal{R}_\Gamma$  contains only one member.*

From the subdivided radial enhancement's construction we obtain the following.

**Observation 2.** *For every plane graph  $\Gamma$  and every  $R(\Gamma) \in \mathcal{R}_\Gamma$  it holds that  $|E(R(\Gamma))| = \mathcal{O}(|E(\Gamma)|)$ .*

Given a plane graph  $\Gamma$  and a graph  $R(\Gamma) \in \mathcal{R}_\Gamma$ , for every integer  $i > 1$ , we denote by  $R^i(\Gamma)$  the graph  $R(R^{i-1}(\Gamma))$ , where  $R^1(\Gamma) = R(\Gamma)$ . We define then  $V_o^i(\Gamma) = V(R^{i-1}(\Gamma))$ ,  $V_s^i = V_s(R^{i-1}(\Gamma))$ , and  $V_r^i = V_r(R^{i-1}(\Gamma))$ . For notational consistency, we define  $V_s^1(\Gamma) = V_s(\Gamma)$ ,  $V_r^1(\Gamma) = V_r(\Gamma)$  and  $V_o^1(\Gamma) = V_o(\Gamma)$ . We also define the sets of edges  $E_s^i(\Gamma)$  which are the edges obtained in  $R^i(\Gamma)$  after subdividing the edges  $E_s^{i-1}(\Gamma)$  and  $E_r^i(\Gamma) = E(R^i(\Gamma)) \setminus E_s^i(\Gamma)$ . Let  $\Gamma$  be a plane graph and  $R(\Gamma)$  be a member of  $\mathcal{R}_\Gamma$ . For each vertex  $u$  of  $R(\Gamma)$ , we define the function  $R^{-1} : V(R(\Gamma)) \rightarrow V(\Gamma) \cup E(\Gamma) \cup F(\Gamma)$  as follows:

- if  $u \in V_o(R(\Gamma)) \equiv V(\Gamma)$  then  $R^{-1}(u) = u$ ,
- if  $u \in V_f(R(\Gamma))$  then  $R^{-1}(u) = f_u$ , where  $f_u \in F(\Gamma)$  is the face in which  $u$  was added, and
- if  $u \in V_s(R(\Gamma))$  then  $R^{-1}(u) = e_u$ , where  $e_u \in E(\Gamma)$  is the edge whose subdivision created  $u$ .

We will call  $R^{-1}(u)$  the *preimage* of  $u$ .

**Lemma 1.** *Let  $\Gamma$  be a plane graph (possibly with multiple edges). Then for every  $R(\Gamma) \in \mathcal{R}_\Gamma$ , it holds that  $R^3(\Gamma)$  is 3-connected. Moreover, for every plane graph  $\Gamma$  each member of  $\mathcal{R}_\Gamma$  is connected and if  $\Gamma$  is  $i$ -connected, then  $R(\Gamma)$  is  $(i + 1)$ -connected, for  $i \in [2]$ .*

In order to prove this lemma and only, we also need the following definitions as well as the next observation. Given a plane graph  $\Gamma$ , an *alternating face walk* is defined as a sequence of the form  $(a_1, a_2, \dots, a_k)$ , for  $k > 1$ , such that:

- for every  $i \in [k - 1]$ , either  $a_i \in V(\Gamma) \cup E(\Gamma)$ ,  $a_{i+1} \in F(\Gamma)$ , and  $a_i$  is in the boundary of  $a_{i+1}$ , or  $a_i \in F(\Gamma)$ ,  $a_{i+1} \in V(\Gamma) \cup E(\Gamma)$ , and  $a_{i+1}$  is in the boundary of  $a_i$ ,
- for every  $i, j \in [k]$ ,  $i \neq j \Rightarrow a_i \neq a_j$ .

We will call  $a_1$  and  $a_k$  the *endpoints* of the alternating face walk. The length of an alternating face walk  $(a_1, a_2, \dots, a_k)$  is  $k$ . The *alternating face distance* between two elements  $u, v$  of  $\Gamma$  is the smallest  $k$  for which there exists an alternating face walk of length  $k$  joining them.

The following observation follows directly from the definition of an alternating face walk.

**Observation 3.** *Let  $\Gamma$  be a plane graph,  $R(\Gamma)$  be a subdivided radial enhancement of  $\Gamma$ , and  $u, v \in V(R(\Gamma))$ . Then  $R(\Gamma)$  contains a  $(u, v)$ -path if and only if  $\Gamma$  contains an alternating face walk with endpoints  $R^{-1}(u)$  and  $R^{-1}(v)$ .*

*Proof of Lemma 1.* The assertion that every member of  $R(\Gamma)$  is connected follows immediately from Observation 3 together with the fact that a plane graph  $\Gamma$  always contains an alternating face walk between any pair  $\{x, y\} \subseteq V(\Gamma) \cup E(\Gamma)$ .

Suppose now that  $\Gamma$  is connected but  $R(\Gamma)$  is not 2-connected and let  $s \in V(R(\Gamma))$  be a cut-vertex of  $R(\Gamma)$ . Then  $s$  cannot be a radial vertex otherwise  $\Gamma$  would not be connected. Moreover,  $s$  cannot be a subdivision vertex, since the endpoints of the corresponding edge of  $\Gamma$  are connected to every neighbor of  $s$  that is a radial vertex (which is, at least one). Therefore  $s$  must be an original vertex of  $\Gamma$  and also, consequently, a cut-vertex of  $\Gamma$ . We choose a pair of vertices  $u, v$  in  $V(R(\Gamma)) \setminus \{s\}$  in such a way that it satisfies the following conditions:

- If  $u_\Gamma := R^{-1}(u)$  and  $v_\Gamma := R^{-1}(v)$  are the preimages of  $u$  and  $v$  in  $\Gamma$  then every alternating face walk joining them in  $\Gamma$  contains  $s$ , and
- among all pairs satisfying the above condition they have the shortest alternating face distance.

Let us first assume that the shortest alternating face walk joining them has length  $k > 3$ . Let  $u'_\Gamma$  be the element of the alternating face walk that is the unique neighbor of  $u_\Gamma$  in this walk and assume that  $u'_\Gamma \neq s$ . Then  $u'_\Gamma$  and  $v_\Gamma$  are joined by an alternating face walk  $w$  of length  $k - 1$ . Thus, from the choice of  $u$  and  $v$ , there exists an alternating face walk joining  $u'_\Gamma$  and  $v_\Gamma$  that does not contain  $s$  and hence we can trivially extend  $w'$  to an alternating face walk joining  $u_\Gamma$  and  $v_\Gamma$  that does not contain  $s$ , a contradiction. Thus  $u'_\Gamma = s$ . Let then  $v'_\Gamma$  be the unique neighbor of  $v_\Gamma$  in the alternating face walk. Then the alternating face walk joining  $u_\Gamma$  and  $v'_\Gamma$  contains  $s$  and has length  $k - 1$ . From the choice of  $u, v$  there exists an alternating face walk  $w'$  joining  $u_\Gamma$  and  $v'_\Gamma$  that avoids  $s$ . Then,  $w'$  can trivially be extended to an alternating face walk joining  $u_\Gamma$  and  $v_\Gamma$  that avoids  $s$ . This is, again, a contradiction. Thus, any shortest alternating face walk joining  $u_\Gamma$  and  $v_\Gamma$  has length 3. This implies that the shortest alternating face walk between  $u_\Gamma$  and  $v_\Gamma$  is  $(u_\Gamma, s, v_\Gamma)$ . Moreover,  $u_\Gamma, v_\Gamma \in F(\Gamma)$ . Then by utilising the edges that are incident to  $s$  and the faces  $f$  where  $s \in \mathbf{bd}(f)$  we may find an alternating face walk joining  $u_\Gamma$  and  $v_\Gamma$  that avoids  $s$ , a contradiction. Therefore,  $R(\Gamma)$  is 2-connected.

Suppose now that  $\Gamma$  is 2-connected but  $R(\Gamma)$  is not 3-connected. Let then  $s, t$  be a cut-pair of  $R(\Gamma)$  and  $s_\Gamma, t_\Gamma$  be their respective preimages in  $R(\Gamma)$ . Notice that neither  $s$  nor  $t$  is a radial vertex, as otherwise  $\Gamma$  would contain a separator of size less than 2. Again, as previously, we choose a pair of vertices  $u, v$  in  $V(R(\Gamma)) \setminus \{s_\Gamma, t_\Gamma\}$  in such a way that it satisfies the following conditions:

- If  $u_\Gamma := R^{-1}(u)$  and  $v_\Gamma := R^{-1}(v)$  are the preimages of  $u$  and  $v$  in  $\Gamma$  then every alternating face walk joining them in  $\Gamma$  contains at least one of the  $s_\Gamma$  and  $t_\Gamma$ , and

- among all pairs satisfying the above condition they have the shortest alternating face distance.

We first show that if  $w$  is a shortest alternating face walk joining  $u_\Gamma, v_\Gamma$  then it contains exactly one of the  $s_\Gamma, t_\Gamma$ . Towards, a contradiction let us assume that both  $s_\Gamma$  and  $t_\Gamma$  belong to  $w$ . Then as  $s_\Gamma, t_\Gamma \in V(\Gamma) \cup E(\Gamma)$  the elements do not appear consecutively in  $w$ . Let  $x_\Gamma$  be an element that appears between  $s_\Gamma$  and  $t_\Gamma$  in  $w$ . Then there exists an alternating face walk joining  $u_\Gamma$  and  $x_\Gamma$  and an alternating face walk joining  $x_\Gamma$  and  $v_\Gamma$  such that both of them have length strictly less than the length of  $w$ . From the choice of  $u$  and  $v$ , we obtain that there exist two alternative face walks  $w_u$  and  $w_v$  joining  $x_\Gamma$  with  $u_\Gamma$  and  $v_\Gamma$  respectively and such that none of them contains any of  $s_\Gamma, t_\Gamma$ . By combining them, we may obtain an alternating face walk joining  $u_\Gamma$  and  $v_\Gamma$  and containing neither  $s_\Gamma$  nor  $t_\Gamma$ , a contradiction. Thus  $w$  contains exactly one of the  $s_\Gamma$  and  $t_\Gamma$ , say  $s_\Gamma$ . Let us assume that  $w$  has length  $k > 3$  and let  $u'_\Gamma$  be the element of the alternating face walk that is the unique neighbor of  $u_\Gamma$  in this walk and assume that  $u'_\Gamma \neq s_\Gamma$ . Then  $u'_\Gamma$  and  $v_\Gamma$  are joined by an alternating face walk  $w'$  of length  $k - 1$ . Thus, from the choice of  $u$  and  $v$ ,  $w'$  does not contain  $s_\Gamma$  and hence the alternating face walk joining  $u_\Gamma$  and  $v_\Gamma$  does not contain  $s_\Gamma$  or  $t_\Gamma$ , a contradiction. Thus  $u'_\Gamma = s_\Gamma$ . Let then  $v'_\Gamma$  be the unique neighbor of  $v_\Gamma$  in the alternating face walk  $w$ . Then the alternating face walk joining  $u_\Gamma$  and  $v'_\Gamma$  contains  $s_\Gamma$  and has length  $k - 1$ . From the choice of  $u, v$  there exists an alternating face walk  $w'$  joining  $u_\Gamma$  and  $v'_\Gamma$  that avoids  $s_\Gamma$ . Then,  $w'$  can trivially be extended to an alternating face walk joining  $u_\Gamma$  and  $v_\Gamma$  that avoids  $s_\Gamma$ . This is a contradiction. Thus, any shortest alternating face walk joining  $u_\Gamma$  and  $v_\Gamma$  has length 3. Moreover,  $w = (u_\Gamma, s_\Gamma, t_\Gamma)$ . If  $s_\Gamma \in V(\Gamma)$  then as above we may find an alternating face walk joining  $u_\Gamma$  and  $v_\Gamma$  that avoids  $s_\Gamma$ , a contradiction. Therefore,  $s_\Gamma \in E(\Gamma)$ . However, in that case,  $s_\Gamma$  belongs to the boundary of both faces  $u_\Gamma$  and  $v_\Gamma$  and at least one of the endpoints of  $s_\Gamma$ , say  $z_\Gamma$ , is different from  $t_\Gamma$ . Therefore,  $(u_\Gamma, z_\Gamma, v_\Gamma)$  is an alternating face walk of  $\Gamma$  avoiding both  $s_\Gamma$  and  $t_\Gamma$ , a contradiction. We conclude that  $R(\Gamma)$  is 3-connected.  $\square$

*Remark 1.* If  $\Gamma$  is 2-connected then  $R(\Gamma)$  can also be shown to be 4-connected. However, 3-connectivity is sufficient for our purposes.

## 2.2 Graph Structures

A key-concept in our algorithms is the notion of the vertex and the edge structure which is formally defined as follows. Let  $G$  be a simple planar graph,  $k, l \in \mathbb{N}$ ,  $(S_1, S_2, \dots, S_k)$  be a near-partition of  $V(G)$  and  $E_1, E_2, \dots, E_l$  be a near-partition of  $E(G)$ . A *vertex structure*  $\mathbf{G}$  is a tuple  $(G, S_1, S_2, \dots, S_k)$  and an *edge structure*  $\mathbf{G}'$  is a tuple  $(G, E_1, E_2, \dots, E_l)$ .

Let  $\mathbf{G} = (G, A, X_1, \dots, X_l)$  and  $\mathbf{D} = (D, B, Y_1, \dots, Y_l)$  be vertex structures, where  $l \in \mathbb{N}$ . We say that  $\mathbf{D}$  is a *contraction* of  $\mathbf{G}$ , denoted by  $\mathbf{D} \leq_c \mathbf{G}$ , if and only if there exists a function  $\sigma : V(G) \rightarrow V(D)$  satisfying the following *contraction properties*:

1. if  $u, v \in V(D)$ ,  $u \neq v \Leftrightarrow \sigma^{-1}(u) \cap \sigma^{-1}(v) = \emptyset$ ,
2. for every  $u \in V(D)$ ,  $G[\sigma^{-1}(u)]$  is connected,
3.  $\{u, v\} \in E(D) \Leftrightarrow G[\sigma^{-1}(u) \cup \sigma^{-1}(v)]$  is connected,
4.  $\sigma(A) \subseteq B$ , and
5. for every  $i \in [l]$  and every  $x \in Y_i$  it holds that  $|\sigma^{-1}(x)| = 1$  and  $\sigma^{-1}(x) \in X_i$ .

In particular, a graph  $D$  is a *contraction* of a graph  $G$  if  $(D, V(D)) \leq_c (G, V(G))$  and we write  $D \leq_c G$ . Notice that  $\leq_c$  defined for graphs is the usual contraction relation where only conditions 1, 2, and 3 apply. Observe that for any two vertex structures  $\mathbf{G}$  and  $\mathbf{D}$ , where  $G$  and  $D$  respectively are their associated planar graphs,  $\mathbf{D} \leq_c \mathbf{G}$  implies that  $D \leq_c G$ .

We will also need the following proposition, which follows from the results in [1].

**Proposition 1.** *There exists an algorithm that receives as input a vertex structure  $\mathbf{G}$ , whose graph has  $m$  edges and treewidth at most  $h$ , and a vertex structure  $\mathbf{D}$ , whose graph is connected and has  $k$  edges, and outputs whether  $\mathbf{D} \leq_c \mathbf{G}$  in  $2^{\mathcal{O}(k+h+k \log h)} \cdot m$  steps.*

Let  $\mathbf{G} = (G, S_1, \dots, S_l)$  be a vertex structure on a planar graph  $G$ , where  $l \in \mathbb{N}$ . Given a possibly empty  $Q \subseteq V(G)$ , notice that the tuple  $(Q, S_1 \setminus Q, \dots, S_l \setminus Q)$  also forms a near-partition of  $V(G)$ . Then, we can define the following operator on vertex structures:

$$\mathbf{d}(\mathbf{G}, Q) := (G, Q, S_1 \setminus Q, \dots, S_l \setminus Q).$$

Obviously,  $\mathbf{d}(\mathbf{G}, Q)$  is also a vertex structure on  $G$ .

Let  $\Gamma$  be a plane graph and consider an  $R(\Gamma) \in \mathcal{R}_\Gamma$ . By Lemma 1 and Observation 1, the graph  $R^3(\Gamma)$  is uniquely defined according to  $R(\Gamma)$ . The following operators on  $(\Gamma, R(\Gamma))$  uniquely define a vertex and an edge structure:

$$\begin{aligned} \mathbf{p}(\Gamma, R(\Gamma)) &:= (R^3(\Gamma), V(\Gamma), V_s^1(\Gamma), V_r^1(\Gamma), V_s^2(\Gamma), V_r^2(\Gamma), V_s^3(\Gamma), V_r^3(\Gamma)) \\ \mathbf{e}(\Gamma, R(\Gamma)) &:= (R^3(\Gamma), E_s^3(\Gamma), E_r^3(\Gamma)). \end{aligned}$$

The underlying graph of the above structure is the general graph of  $R^3(\Gamma)$  and the vertex sets that form the partition of  $V(R^3(\Gamma))$  are the original vertices  $V(\Gamma)$ , followed by the sets of the subdivision and the radial vertices of each of the three subdivided radial enhancements. Moreover, the edges are separated to those that have been obtained in  $R^3(\Gamma)$  only by subdividing original edges of the graph and those that were obtained after adding radial vertices and edges and subdividing those edges.

Throughout the rest of the paper we will only use structures defined by those three operators. The main purpose is to associate three subdivided radial enhancements to a given plane graph so that (i) the resulting graph is 3-connected and therefore uniquely embeddable, so we can disregard the embedding and treat it as a combinatorial object, and (ii) the vertices and edges of the original graph and each subdivided radial enhancement are distinguishable. In addition, both in PSC and the PTMC problems we try to match the faces of the pattern graph to faces, or parts of faces, of the host graph, the radial enhancements and the corresponding structures seem to be the appropriate tool to use, since we actually need to match just the radial vertices that are added inside each face.

Given a graph  $G$  and a non-negative integer  $k$ , we define the *ball* around a vertex  $v$  of  $G$  as the subgraph  $B_G^k(v)$  of  $G$  induced by the set of vertices at distance at most  $k$  from  $v$ . Consider now the subgraph  $\tilde{G}$  of  $G$  induced by the set of vertices that lay outside a given ball  $B_G^k(v)$ , i.e.,  $\tilde{G} = G \setminus B_G^k(v)$ , and consider the set  $\mathcal{C}(\tilde{G})$  of all its connected components. Then by contracting all the edges of every  $C \in \mathcal{C}(\tilde{G})$  to a single vertex in  $G$ , denoted  $v_C$ , we obtain the *k-contracted graph* around  $v$ , that will be denoted by  $G_v^{(k)}$ . We can now make this contracted graph into a structure as follows. Given a vertex structure  $\mathbf{G} = (G, \emptyset, S_1, \dots, S_l)$  and a non-negative integer  $k$ , we define the *k-contracted vertex structure* around a vertex  $v$  of the graph  $G$  as  $\mathbf{G}_v^{(k)} := (G_v^{(k)}, \{v_C \mid C \in \mathcal{C}(\tilde{G})\}, S'_1, \dots, S'_l)$ , where  $S'_i = S_i \cap B_G^k(v)$  for every  $i \in [l]$ .



### 3 An FPT algorithm for the PSC problem

Given a plane graph  $\Gamma$  we define *the set of non-edges of  $\Gamma$* :  $\overline{E(\Gamma)} = \binom{V(\Gamma)}{2} \setminus E(\Gamma)$ . A set of non-edges  $S \subseteq \overline{E(\Gamma)}$  will be called *insertable* if there is a way to add the edges to  $\Gamma$  such that no two edges of  $E(\Gamma) \cup S$  intersect (apart from any common endpoints). Finally, we define the following relation between two plane graphs  $\Gamma$  and  $\Delta$ . We say that  $\Delta \preceq \Gamma$  if there exists a set  $S \subseteq \overline{E(\Gamma)}$  of insertable edges of  $\Gamma$  such that  $\Delta \leq_{es} \Gamma'$ , where  $\Gamma'$  is obtained from  $\Gamma$  after adding  $S$ . Then PSC asks, given two plane graphs  $\Gamma$  and  $\Delta$ , whether  $\Delta \preceq \Gamma$ .

The main idea of our algorithm is to create two families of vertex structures, one from the host graph  $\Gamma$  and the other from the pattern graph  $\Delta$ , such that  $\Delta \preceq \Gamma$  if and only if there are two structures  $\mathbf{D}$  and  $\mathbf{G}$  from each of the above families such that  $\mathbf{D} \leq_c \mathbf{G}$ . Then, we bound the size of these families and use the algorithm from Proposition 1 to check all pairs of their members for the required property. From now on, in this section, whenever we refer to a structure we will assume that it is a vertex structure.

We define the first family of structures based on the host graph. Given a plane graph  $\Gamma$ , a subdivided radial enhancement of it,  $R(\Gamma)$ , and a positive integer  $k$ , we define the following family of structures:

$$\mathcal{G}_{\Gamma, R(\Gamma), k} := \{\mathbf{d}(\mathbf{p}(\Gamma, R(\Gamma)), \emptyset)_v^{(k)} \mid v \in V(\Gamma)\}.$$

Obviously,  $|\mathcal{G}_{\Gamma, R(\Gamma), k}| = |V(\Gamma)|$ , regardless of the choice of  $R(\Gamma)$  and  $k$ . In the following lemma we bound the treewidth of the underlying graphs of all members of this family.

**Lemma 2.** *Let  $\Gamma$  be a plane graph,  $R(\Gamma)$  a subdivided radial enhancement of  $\Gamma$ ,  $k \in \mathbb{N}$ ,  $v \in V(\Gamma)$ , and  $\mathbf{G}_v := \mathbf{d}(\mathbf{p}(\Gamma, R(\Gamma)), \emptyset)_v^{(k)} \in \mathcal{G}_{\Gamma, R(\Gamma), k}$ . Then the underlying graph  $G_v$  of the structure  $\mathbf{G}_v$  has treewidth at most  $3(k+1)$  and size  $\mathcal{O}(|E(\Gamma)|)$ .*

*Proof.* Observe that the diameter of  $G_v$  is, by construction, at most  $k+1$  and that  $G_v$  remains planar. From [4, Theorem 4] we obtain that  $G_v$  has treewidth at most  $3(k+1)$ . For the size, notice first that  $G_v \equiv R^3(\Gamma)_v^{(k)}$ . From Observation 2,  $|E(R^3(\Gamma))| = \mathcal{O}(|E(\Gamma)|)$  and since  $R^3(\Gamma)_v^{(k)} \leq_m R^3(\Gamma)$ , it follows that  $|E(G_v)| = \mathcal{O}(|E(\Gamma)|)$ .  $\square$

In order to define the second family of structures based on the pattern graph we need the following two definitions.

A *facial extension* of a connected plane graph  $\Delta$  is a connected plane graph  $\Delta^+$  satisfying the following properties:

1.  $\Delta \subseteq \Delta^+$ ,
2.  $V(\Delta^+) \setminus V(\Delta)$  is an independent set in  $\Delta^+$ , and
3. for every distinct  $x, y \in V(\Delta^+) \setminus V(\Delta)$ ,  $N_{\Delta^+}(x) \not\subseteq N_{\Delta^+}(y)$ .

We will denote by  $\mathcal{F}_\Delta$  the family of all facial extensions of the graph  $\Delta$ .

Given a connected plane graph  $\Delta$  and a subset  $L \subseteq E(\Delta)$  of its edges, we denote by  $\text{span}(\Delta, L)$  the set of all spanning subgraphs of  $\Delta$  that contain all the edges in  $E(\Delta) \setminus L$ . Note that such subgraphs could also contain some edges in  $L$ . A *pattern-guess* of a connected plane graph  $\Delta$  is an element  $\Delta^*$  of  $\text{span}(\Delta^+, E(\Delta))$ , for  $\Delta^+ \in \mathcal{F}_\Delta$ . That is, a spanning subgraph of a facial extension  $\Delta^+$  of  $\Delta$  containing at least all the edges in  $E(\Delta^+) \setminus E(\Delta)$ . The family of all possible pattern-guesses  $\Delta^*$  of  $\Delta$  will be denoted by  $\mathcal{PG}_\Delta$ .

Now, given a connected plane graph  $\Delta$  we define the following family of structures:

$$\mathcal{H}_\Delta := \{\mathbf{d}(\mathbf{p}(\Delta^*, R(\Delta^*)), V(\Delta^+) \setminus V(\Delta)) \mid \Delta^* \in \mathcal{PG}_\Delta, R(\Delta^*) \in \mathcal{R}_{\Delta^*}\}.$$

In the following lemma we bound the size of this second family and also the size and diameter of the underlying graphs of all members of the family.

**Lemma 3.** *If  $\Delta$  is a connected plane graph then  $|\mathcal{H}_\Delta| = 2^{\mathcal{O}(|E(\Delta)| \cdot \log |E(\Delta)|)}$  and, for any structure  $\mathbf{D} \in \mathcal{H}_\Delta$ , the underlying graph  $D$  of  $\mathbf{D}$  has size and diameter bounded by  $\mathcal{O}(|E(\Delta)|)$ .*

*Proof.* First, note that  $D \equiv R^3(\Delta^*)$ , for some  $\Delta^* \in \mathcal{PG}_\Delta$ . Then, recall that since  $\Delta^*$  is a spanning subgraph of a facial extension  $\Delta^+$  of  $\Delta$ , the edges of  $\Delta^*$  consist of the vertices of  $\Delta^+$  and some of the edges of  $\Delta^+$ . By the construction of  $\Delta^+$ , in each face of  $\Delta$  the number of added vertices is a linear function of the size of the boundary of the face. Therefore, since  $\Delta^+$  is plane,  $|E(\Delta^+)|$  is a linear function of  $|E(\Delta)|$ . Thus,  $|E(\Delta^*)| = \mathcal{O}(|E(\Delta)|)$  and from Observation 2 we get  $|E(R^3(\Delta^*))| = \mathcal{O}(|E(\Delta^*)|) = \mathcal{O}(|E(\Delta)|)$ . Obviously, since  $R^3(\Delta^*)$  is connected,  $\text{diam}(R^3(\Delta^*)) \leq |E(R^3(\Delta^*))| = \mathcal{O}(|E(\Delta)|)$ .

Second, note that  $|\mathcal{H}_\Delta| \leq |\mathcal{PG}_\Delta| \cdot \max\{|\mathcal{R}_{\Delta^*}| \mid \Delta^* \in \mathcal{PG}_\Delta\}$ . If  $\Delta^*$  is disconnected, then  $\mathcal{R}_{\Delta^*}$  may contain at most  $\mathcal{O}(|E(\Delta)|!) = 2^{\mathcal{O}(|E(\Delta)| \cdot \log |E(\Delta)|)}$  members. It is easy to see why this holds. In the construction of the subdivided radial enhancement, inside each face there will be created regions linear to the number of edges that are adjacent to that face. Then, the different connected components that lie in that face can be placed inside any of these regions and more edges will be added, equal to size of the facial walks around the connected components (i.e, up to twice the size of the boundary of the connected components). All these can be bounded by a factorial function of the size of the boundary of the face and since the number of faces of the graph is linear to the number of edges, the above bound is reached.

A pattern-guess  $\Delta^*$  of  $\mathcal{PG}_\Delta$  is constructed by first choosing a facial extension  $\Delta^+ \in \mathcal{F}_\Delta$ . Now, observe that in order to enumerate all the possible facial extensions of  $\Delta$ , we can restrict ourselves to the enumeration of the facial extensions of a single face  $F$  of size  $q$  as, due to planarity, the sum of the degrees of its faces is a linear function of the number of its edges. Each facial extension of the face  $F$  can be constructed by adding at most  $q$  vertices inside  $F$  in layers, such that in each layer the neighborhoods of the vertices are non-crossing partitions of the boundary of  $F$ , where a *non-crossing partition*  $P$  of a set  $S$  is a partition with the extra property that if  $u_1, u_2, u_3, u_4$  are vertices (not necessarily successive but strictly in that order) of the facial mapping of  $F$ , then  $\forall S_1, S_2 \in P$  if  $\{u_1, u_3\} \subseteq S_1$  and  $u_2 \in S_2$  then  $u_4 \notin S_2$ . Since the non-crossing partitions of a set of size  $q$  can be bounded by the  $q$ -th Catalan number, i.e., by  $\mathcal{O}(4^q) = 2^{\mathcal{O}(q)}$ , the total number of the facial extensions of  $\Delta$  can be bounded by  $2^{\mathcal{O}(|V(\Delta)| \cdot \log |V(\Delta)|)} = 2^{\mathcal{O}(|E(\Delta)| \cdot \log |E(\Delta)|)}$ , since  $\Delta$  is connected. Therefore  $|\mathcal{F}_\Delta| = 2^{\mathcal{O}(|E(\Delta)| \cdot \log |E(\Delta)|)}$ . After choosing  $\Delta^+$ , we then select an element of  $\text{span}(\Delta^+, E(\Delta))$ , i.e., we choose and remove a subset of  $E(\Delta)$  from  $\Delta^+$ . This way we can obtain any member of  $\mathcal{PG}_\Delta$ . Hence, we conclude that

$$|\mathcal{PG}_\Delta| = 2^{|E(\Delta)|} \cdot 2^{\mathcal{O}(|E(\Delta)| \cdot \log |E(\Delta)|)} = 2^{\mathcal{O}(|E(\Delta)| \cdot \log |E(\Delta)|)}$$

and thus  $|\mathcal{H}_\Delta| = 2^{\mathcal{O}(|E(\Delta)| \cdot \log |E(\Delta)|)}$ . □

The next two lemmata will lead us to Theorem 1 which ensures the correctness of our algorithm.

**Lemma 4.** *Let  $\Gamma$  be a plane graph and  $\Delta$  be a connected plane graph. If for every  $R(\Gamma) \in \mathcal{R}_\Gamma$  there exists a positive integer  $c$  and two structures  $\mathbf{G} \in \mathcal{G}_{\Gamma, R(\Gamma), c}$  and  $\mathbf{D} \in \mathcal{H}_\Delta$ , such that  $\mathbf{D} \leq_c \mathbf{G}$ , then  $\Delta \preceq \Gamma$ .*

*Proof.* Suppose now that for every  $R(\Gamma) \in \mathcal{R}_\Gamma$  there exist two structures  $\mathbf{G} \in \mathcal{G}_{\Gamma, R(\Gamma), c}$  and  $\mathbf{D} \in \mathcal{H}_\Delta$ , such that  $\mathbf{D} \leq_c \mathbf{G}$ . This is the same as saying that for every  $R(\Gamma) \in \mathcal{R}_\Gamma$  there exist a  $\Delta^+ \in \mathcal{F}_\Delta$ , a  $\Delta^* \in \text{span}(\Delta^+, E(\Delta))$ , and an  $R(\Delta^*) \in \mathcal{R}_{\Delta^*}$  such that  $\mathbf{d}(\mathbf{p}(\Delta^*, R(\Delta^*)), B)$

$\leq_c \mathbf{d}(\mathbf{p}(\Gamma, R(\Gamma)), \emptyset)$ . Let then  $\Gamma'$  be the plane graph that results from  $R^3(\Gamma)$  if we contract all connected components of  $R^3(\Gamma)[\sigma^{-1}(B)]$ . It follows immediately that  $\Gamma' \simeq_{tp} R^3(\Delta^*)$ . Let

$$\alpha : V(\Gamma') \cup F(\Gamma') \rightarrow V(R^3(\Delta^*)) \cup F(R^3(\Delta^*))$$

be a topological isomorphism between  $\Gamma'$  and  $R^3(\Delta^*)$ . Then, for each edge  $\{u, v\} \in E(\Delta) \setminus E(\Delta^*)$  there is a face  $f \in F(\Gamma)$  such that both  $\alpha^{-1}(u)$  and  $\alpha^{-1}(v)$  belong to a member of the facial mapping of  $f$ . Hence, the set  $S = \{\{\alpha^{-1}(u), \alpha^{-1}(v)\} \mid \{u, v\} \in E(\Delta) \setminus E(\Delta^*)\}$  is insertable in  $\Gamma$ . Hence,  $\Delta \preceq \Gamma$ .  $\square$

**Lemma 5.** *Let  $\Gamma$  be a plane graph and  $\Delta$  be a connected plane graph. If  $\Delta \preceq \Gamma$ , then for every  $R(\Gamma) \in \mathcal{R}_\Gamma$  there exist two structures  $\mathbf{G} \in \mathcal{G}_{\Gamma, R(\Gamma), c}$  and  $\mathbf{D} \in \mathcal{H}_\Delta$ , such that  $\mathbf{D} \leq_c \mathbf{G}$ , where  $c$  is a constant such that  $\max_{\Delta^*} \{\mathbf{diam}(R^3(\Delta^*))\} \leq c$ .*

To prove Lemma 5 we need the following lemma, which asserts that in order to find in the host graph a subgraph isomorphic to the pattern graph containing a specific vertex, then all the other vertices participating in the pattern can be found at a distance from the starting vertex bounded by the diameter of the pattern graph. This fact is not true in the original graphs per se, but it holds for their subdivided radial enhancements.

**Lemma 6.** *Let  $\mathbf{G} = (G, \emptyset, S_1, \dots, S_l)$  and  $\mathbf{D} = (D, B, Z_1, \dots, Z_l)$  be two structures, where  $B$  is an independent set and  $l \in \mathbb{N}$ . Then  $\mathbf{D} \leq_c \mathbf{G}$  if and only if there exists some  $v \in V(G)$  such that  $\mathbf{D} \leq_c \mathbf{G}_v^{(k)}$ , where  $k := \mathbf{diam}(D)$ .*

*Proof.* Suppose first that  $\mathbf{D} \leq_c \mathbf{G}$ , i.e., there exists a function  $\sigma : V(G) \rightarrow V(D)$  satisfying the contraction properties. Then  $D \leq_c G$ , and thus there exists a minor  $D'$  of  $G$  such that  $D \simeq D'$ . Moreover, notice that all vertices of  $V(D) \setminus B$  are mapped to single vertices of  $G$ . Then, there exists a vertex  $v \in V(D') \subseteq V(G)$  such that  $D \setminus B \simeq D' \setminus (\sigma^{-1}(B)) \subseteq B_G^k(v)$ . Consider now the structure  $\mathbf{G}_v^{(k)} := (G_v, \{u_C \mid C \in \mathcal{C}(\tilde{G})\}, S'_1, \dots, S'_l)$ , where  $G_v$  is the  $k$ -contracted graph around the vertex  $v$ ,  $\tilde{G} = G \setminus B_G^k(v)$  and  $S'_i = S_i \cap B_G^k(v)$ , for every  $i \in [l]$ . Observe that  $G_v = B_G^k(v) \uplus \{u_C \mid C \in \mathcal{C}(\tilde{G})\}$ . Let then the function  $\rho : V(G_v) \rightarrow V(D)$  be defined as follows.

- $\rho(u) = \sigma(u)$ , for every  $u \in V(B_G^k(v))$ .
- $\rho(u_C) = v$ , where  $v \in B$  and there exists  $u \in C$  such that  $\sigma(u) = v$ .

Note then that since  $D' \setminus \sigma^{-1}(B) \subseteq B_G^k(v)$  it holds that  $\rho^{-1}(D) = \sigma^{-1}(D)$ . This implies that  $\rho$  satisfies the first three contraction properties. Moreover, since  $\rho(u_C) \in B$ , for every  $C \in \mathcal{C}(\tilde{G})$ , then  $\rho(\{u_C \mid C \in \mathcal{C}(\tilde{G})\}) \subseteq B$  and thus,  $\rho$  satisfies the fourth contraction property as well. Finally, as  $S'_i \subseteq S_i \cap B_G^k(v)$ , then  $\rho$  also satisfies the fifth contraction property. Therefore for the chosen  $v \in V(G)$  we obtain that  $\mathbf{D} \leq_c \mathbf{G}_v^{(k)}$ .

Suppose now that there exists a vertex  $v \in V(G)$  such that  $\mathbf{D} \leq_c \mathbf{G}_v^{(k)}$ . Then there exists a function  $\sigma : V(G_v) \rightarrow V(D)$  satisfying the contraction properties. Notice that  $\mathbf{G}_v^{(k)} \leq_c \mathbf{G}$  (where every vertex of  $B_G^k(v)$  is mapped to itself and every vertex  $u_C$  is mapped to the connected component  $C$ ). Thus, there exists a function  $\rho : V(G) \rightarrow V(G_v)$  satisfying the contraction properties. It is easy to confirm that  $\sigma \circ \rho : V(G) \rightarrow V(D)$  satisfies the contraction properties and therefore  $\mathbf{D} \leq_c \mathbf{G}$ .  $\square$

We are now ready to prove the section's main result.

*Proof of Lemma 5.* First of all, we know that such a constant  $c$  exists from Lemma 3 and that in fact  $c = \mathcal{O}(|E(\Delta)|)$ . Since  $\Delta \preceq \Gamma$ , there exists an insertable set of non-edges  $S \subseteq \overline{E(\Gamma)}$  and two plane graphs  $\Gamma' = (V(\Gamma), E(\Gamma) \cup S)$  and  $\Gamma_0$ , such that  $\Gamma_0 \subseteq \Gamma'$  and  $\Delta \simeq_{tp} \Gamma_0$ . Without loss of generality we may assume that all edges of  $S$  are also edges of  $\Gamma_0$ . Let then  $\alpha : V(\Gamma_0) \cup F(\Gamma_0) \rightarrow V(\Delta) \cup F(\Delta)$  be a topological isomorphism between  $\Gamma_0$  and  $\Delta$ . For every edge  $e = \{u, v\}$  of  $S$  let  $e_\alpha = \{\alpha(u), \alpha(v)\}$ . We define the sets  $S_\alpha = \{e_\alpha \mid e \in S\}$ ,  $S_\alpha^\Delta = S_\alpha \cap E(\Delta)$ , and  $S_\alpha^\Gamma = S_\alpha \setminus S_\alpha^\Delta$ .

We first construct a graph  $\Delta^+ \in \mathcal{F}_\Delta$ . For this, we add a set of vertices and edges embedded inside some of the faces of  $\Delta$  in such a way that edges intersect only at their common endpoints. In particular, for each face  $f \in F(\Delta)$  with facial mapping  $\mathbf{w}(f)$  do the following:

- For each edge  $e = \{u, v\}$  that lies inside the region enclosed by  $\alpha^{-1}(\mathbf{w}(f))$  in  $\Gamma$  and whose endpoints belong to  $\Gamma'$ , add the edge  $\{\alpha(u), \alpha(v)\}$  in the interior of  $f$  in  $\Delta$  in such a way that (i) edges intersect only at their common endpoints and (ii) after we extend the mapping  $\alpha$  so that it takes into account those edges of  $\Gamma$  that were added in  $\Delta$ , the following must hold: for any connected component that was inside  $f$  and, after the addition of the edges, is in a face  $f'$ , the preimages of the vertices of that connected component in  $\Gamma_0$  are inside the region enclosed by  $\alpha^{-1}(\mathbf{w}(f'))$ .
- Consider the faces  $f_1, f_2, \dots, f_j$  that form the partition of  $f$  after the addition of the new edges. For every such face  $f_i$  let  $p_i$  be the region enclosed by  $\alpha^{-1}(\mathbf{w}(f_i))$  in  $\Gamma'$ . Notice that since  $\Delta^+$  is connected, the boundary of  $\alpha^{-1}(\mathbf{w}(f_i))$  is connected. For every  $i \in [j]$  let  $\mathcal{C}_{p_i}$  be the set of all connected components that lie entirely in the region enclosed by  $\alpha^{-1}(\mathbf{w}(f_i))$  in  $\Gamma'$ . Let  $\mathcal{C}_{p_i}^\emptyset$  denote the set of all connected components in  $\mathcal{C}_{p_i}$  that do not have any neighbors in  $B_{\Gamma'}(f_i)$ . For every  $C_w \in \mathcal{C}_{p_i}$ , let  $S_w$  be its neighborhood in  $B_\Gamma(f_i)$ . Consider the Hasse diagram defined by the sets  $S_w$  and without loss of generality, let  $S_1, S_2, \dots, S_q$  be its maximal elements. Let then  $O_t = \{C_l \in \mathcal{C}_{p_i} \setminus \mathcal{C}_{p_i}^\emptyset \mid S_l \subseteq S_t\}$ ,  $t \in [q]$ . For every  $t \in [q]$ , add a vertex  $u_t$  in  $f_i$  and make it adjacent to the vertices in  $\alpha(S_t)$  (notice that since the boundary is again connected there is a unique way to construct the cyclic neighborhood of  $u_t$  up to cyclic permutations). We call  $O_t$  the *origin* of  $u_t$ .

The resulting graph  $\Delta^+$  is, by definition, a member of  $\mathcal{F}_\Delta$ . See an example of such a construction in Figure 2.




Figure 2: The construction of  $\Delta^+$ . The new vertices are the vertices of  $B = V(\Delta^*) \setminus V(\Delta)$ .

To construct  $\Delta^*$  from  $\Delta^+$ , for every edge  $\{u, v\} \in S$ , we remove the edge  $\{\alpha(u), \alpha(v)\}$  from  $\Delta^+$ . Since  $\{\alpha(u), \alpha(v)\} \in E(\Delta)$ , it follows that

$$\Delta^* \in \text{span}(\Delta^+, E(\Delta)).$$

We now define a function  $g_0 : E(\Delta^*) \cup F(\Delta^*) \mapsto E(\Gamma) \cup F(\Gamma)$ . Let  $f \in F(\Delta^*)$  with facial mapping  $\mathbf{w}(f)$ . Observe that there is at least one face  $f' \in F(\Gamma)$  with facial mapping  $\mathbf{w}(f')$ , such that for every facial walk  $w = (u_1, \dots, u_k) \in \mathbf{w}(f)$  there is a facial walk  $w' \in \mathbf{w}(f')$  of length at least  $k$  and a subsequence  $(v_1, \dots, v_k)$  of  $w'$  (up to cyclic permutations) with the following properties:  $v_i = \alpha(u_i)$  if  $v_i \in \Delta$  and  $v_i \in V(C)$ , for some  $C$  in the origin of  $u_i$ , if  $u_i \in V(\Delta^*) \setminus V(\Delta)$ .

Notice that due to planarity the regions defined by those walks (unless the walks are trivial) are mutually nested. Of all such faces (if there are multiple), let  $f'$  be the one whose region contains all other regions. Then,  $g_0(f) = f'$ . We will call the connected component whose vertices belong to that walk the *outermost connected component*.

Recall that, by construction, the new vertices of  $V(\Delta^*) \setminus V(\Delta)$  form an independent set. Thus, for each edge  $e = \{u, v\} \in E(\Delta^*)$  at most one of its endpoints belongs in  $V(\Delta^*) \setminus V(\Delta)$ . If both endpoints  $u, v$  of  $e$  belong to  $V(\Delta)$ , then we define  $g_0(e) = \{\alpha^{-1}(u), \alpha^{-1}(v)\} \in E(\Gamma)$ . Otherwise exactly one of  $u$  and  $v$ , say  $v$ , belongs to  $V(\Delta^*) \setminus V(\Delta)$ . In this case, we define  $g_0(e) = \{\alpha^{-1}(u), v'\} \in E(\Gamma)$ , where  $v'$  is a neighbor of  $\alpha^{-1}(u)$  in the outermost connected component in the origin of  $v$ .

Let now  $R(\Gamma)$  be an arbitrary subdivided radial enhancement of  $\Gamma$ . In order to construct a subdivided radial enhancement  $R(\Delta^*)$  of  $\Delta$  recall that we first subdivide all edges of  $R(\Delta^*)$  and then add a radial vertex  $u_f$  inside each face  $f \in F(\Delta^*)$ . For every  $f$  let  $r_{g_0}(f)$  be the radial vertex of  $R(\Gamma)$  that was added in  $g_0(f)$ . Consider the cyclic neighborhood of  $r_{g_0}(f)$  in  $R(\Gamma)$ . Notice that it can be broken down in  $s_1, s_2, \dots, s_l$  segments where  $s_i$  is a facial walk  $w_i$  of  $\mathbf{w}(g_0(f))$ . Let  $w'_i$  be the subsequence of the walk that corresponds to a walk  $z_i$  in  $\mathbf{w}(f)$ . Add edges between the  $u_f$  and the vertices of the boundary of  $u_f$  in such a way that the cyclic neighborhood of  $u_f$  is  $(z_1, z_2, \dots, z_l)$ . Notice that for every subdivision vertex  $x$  of  $R(\Delta^*)$  that appears between  $u_i$  and  $u_{i+1}$  in the facial walk of  $w$ , there is a subdivision vertex  $v_x$  appearing between  $v_i$  and  $v_{i+1}$  in the walk  $w$  of  $\mathbf{w}(f)$ . We add an edge  $\{u_f, v_x\}$  so that  $v_x$  appears between  $u_i$  and  $u_{i+1}$  in the cyclic neighborhood of  $u_f$  (this can be done in a unique way). We extend the mapping  $g_0$  restricted to  $E(\Delta^*)$  to the mapping  $g_1$  by mapping every edge  $\{u_f, u_i\}$  to the edge  $\{r_{g_0}(f), v_i\}$ . We also map the edges  $\{u_f, x\}$  to the edges  $\{r_{g_0}(f), v_x\}$ . Notice that  $g_1$  can be extended to  $F(R(\Delta^*))$  similarly to  $g_0$ . In the same fashion we extend  $g_1$  to the function  $g_2$  on the graphs  $R^2(\Gamma)$  and  $R^2(\Delta^*)$  and then to  $g_3$  on the graphs  $R^3(\Gamma)$  and  $R^3(\Delta^*)$ . Recall that

$$\mathbf{d}(\mathbf{p}(\Gamma, R(\Gamma)), \emptyset) = (R^3(\Gamma), \emptyset, V(\Gamma), V_s^1(\Gamma), V_r^1(\Gamma), \dots, V_s^3(\Gamma), V_r^3(\Gamma)),$$

and that

$$\begin{aligned} \mathbf{d}(\mathbf{p}(\Delta^*, R(\Delta^*)), B) &= \\ &= (R^3(\Delta^*), V(\Delta^*) \setminus V(\Delta), V(\Delta), V_s^1(\Delta^*), V_r^1(\Delta^*), \dots, V_s^3(\Delta^*), V_r^3(\Delta^*)). \end{aligned}$$

Let now  $\sigma : V(R^3(\Gamma)) \rightarrow V(R^3(\Delta^*))$  such that:

$$\sigma(v) = \begin{cases} u & \text{if } v \in V(\Gamma), u \in V(\Delta), \text{ and } \alpha^{-1}(u) = v \in V(\Gamma) \\ z & \text{if } v \in V_s^i(\Gamma) \text{ and there exists } u \in V_s^i(\Delta^*) \text{ with } g^i(e) = e', i \in [3], \\ & \text{where } z \text{ (resp. } v) \text{ is the subdivision vertex of the edge } e \text{ (resp. } e') \\ w & \text{if } v \in V_r^i(\Gamma) \text{ and there exists } u \in V_r^i(\Delta^*) \text{ with } g^i(f) = f', i \in [3], \\ & \text{where } w \text{ (resp. } v) \text{ is the radial vertex added in face } f \text{ (resp. } f') \\ x & \text{where } x \in B \text{ such that the distance between } v \text{ and the vertices in} \\ & O_x \text{ in } R^3(\Gamma) \text{ is minimized} \end{cases}$$

It is quite straightforward to verify that  $\sigma$  satisfies the five required contraction properties and thus  $\mathbf{d}(\mathbf{p}(\Delta^*, R(\Delta^*)), B) \leq_c \mathbf{d}(\mathbf{p}(\Gamma, R(\Gamma)), \emptyset)$ . Therefore, since these two structures satisfy the conditions of Lemma 6, we conclude that there exists some  $v \in V(\Gamma)$  such that  $\mathbf{d}(\mathbf{p}(\Delta^*, R(\Delta^*)), B) \leq_c \mathbf{d}(\mathbf{p}(\Gamma, R(\Gamma)), \emptyset)_v^{\mathbf{diam}(R^3(\Delta^*))}$ . Notice now that  $\mathbf{d}(\mathbf{p}(\Delta^*, R(\Delta^*)), B) \in \mathcal{H}_\Delta$  and that  $\mathbf{d}(\mathbf{p}(\Gamma, R(\Gamma)), \emptyset)_v^{\mathbf{diam}(R^3(\Delta^*))}$  is a minor of  $\mathbf{d}(\mathbf{p}(\Gamma, R(\Gamma)), \emptyset)_v^c \in \mathcal{G}_{\Gamma, R(\Gamma), c}$  and we have proven our claim.  $\square$

The next theorem is a direct consequence of Lemma 4 and Lemma 5.

**Theorem 1.** *Let  $\Gamma$  be a plane graph and  $\Delta$  be a connected plane graph. It holds that  $\Delta \preceq \Gamma$  if and only if for every  $R(\Gamma) \in \mathcal{R}_\Gamma$  there exist two structures  $\mathbf{G} \in \mathcal{G}_{\Gamma, R(\Gamma), c}$  and  $\mathbf{D} \in \mathcal{H}_\Delta$ , such that  $\mathbf{D} \leq_c \mathbf{G}$ , where  $c$  is a constant such that  $\max_{\Delta^*} \{\mathbf{diam}(R^3(\Delta^*))\} \leq c$ .*

**Theorem 2.** *There exists an algorithm that, given as input an  $n$ -edge plane graph  $\Gamma$  and a connected  $k$ -edge plane graph  $\Delta$ , decides whether  $\Delta \preceq \Gamma$  in  $2^{\mathcal{O}(k \log k)} \cdot n^2$  steps.*

*Proof.* Let  $\Gamma$  and  $\Delta$  be two plane graphs, where  $\Delta$  is connected. From Theorem 1, we have  $\Delta \preceq \Gamma$  if and only if for every  $R(\Gamma) \in \mathcal{R}_\Gamma$  there exist two structures  $\mathbf{G} \in \mathcal{G}_{\Gamma, R(\Gamma), c}$  and  $\mathbf{D} \in \mathcal{H}_\Delta$ , such that  $\mathbf{D} \leq_c \mathbf{G}$ , where  $c$  is a constant such that  $\max\{\mathbf{diam}(R^3(\Delta^*)) \mid \Delta^* \in \mathcal{PG}_\Delta\} \leq c = \mathcal{O}(|E(\Delta)|)$ . Thus, in the first step the algorithm fixes an arbitrary  $R(\Gamma) \in \mathcal{R}_\Gamma$  and constructs the families  $\mathcal{G}_{\Gamma, R(\Gamma), c}$  and  $\mathcal{H}_\Delta$ . By definition,  $|\mathcal{G}_{\Gamma, R(\Gamma), c}| = n$  and from Lemma 3,  $|\mathcal{H}_\Delta| = 2^{\mathcal{O}(k \cdot \log k)}$ . Therefore the algorithm needs to check all possible pairs of structures which are  $2^{\mathcal{O}(k \cdot \log k)} \cdot n$ . To do that we can use the algorithm from Proposition 1, since from Lemma 2, the underlying graphs of all structures in  $\mathcal{G}_{\Gamma, R(\Gamma), c}$  have treewidth  $\mathcal{O}(k)$ . The algorithm from Proposition 1 checks each pair in  $2^{\mathcal{O}(k \cdot \log k)} \cdot n$  steps, therefore the whole algorithm outputs an answer in at most  $2^{\mathcal{O}(k \cdot \log k)} \cdot n^2$  steps. For a more detailed description of the algorithm, see also Algorithm 1.  $\square$

**Input** : An encoding an  $n$ -edged plane graph  $\Gamma$  and a  $k$ -edged plane graph  $\Delta$ .  
**Question:** Is it true that  $\Delta \preceq \Gamma$ ?

```

1 arbitrarily construct an  $R(\Gamma) \in \mathcal{R}_\Gamma$ ;
2 foreach facial extension  $\Delta^+$  of  $\Delta$  do
3   | foreach pattern-guess  $\Delta^*$  based on  $\Delta^+$  do
4   |   | construct a member of  $\mathcal{PG}_\Delta$ ;
5   |   | //  $|\mathcal{PG}_\Delta| = 2^{\mathcal{O}(k \log k)}$  by Lemma 3
6   |   | end
7   | end
8   | construct  $\mathcal{R}_{\Delta^*}$  based on  $\mathcal{PG}_\Delta$ ;
9   | construct the family  $\mathcal{H}_\Delta$  based on  $\mathcal{R}_{\Delta^*}$  and  $\mathcal{PG}_\Delta$ ;
10  | //  $|\mathcal{H}_\Delta| = 2^{\mathcal{O}(k \log k)}$  by Lemma 3
11  | set  $c \geq \max\{\text{diam}(R^3(\Delta^*)) \mid \Delta^* \in \mathcal{PG}_\Delta\}$ ;
12  | //  $c = \mathcal{O}(k)$  by Lemma 3
13  | foreach  $u \in V(\Gamma)$  do construct a member of the family  $\mathcal{G}_{\Gamma, R(\Gamma), c}$ ;
14  | foreach  $\mathbf{G} \in \mathcal{G}_{\Gamma, R(\Gamma), c}$  do //  $n$  steps
15  |   | foreach  $\mathbf{D} \in \mathcal{H}_\Delta$  do //  $\mathcal{O}(k \log k)$  steps
16  |   |   | if  $\mathbf{D} \leq_c \mathbf{G}$  then return True;
17  |   |   | // in  $2^{\mathcal{O}(k \log c)} \cdot n$  steps using the algorithm of Proposition 1 since
18  |   |   |   | the underlying graph of  $\mathbf{G}$  has treewidth  $\leq 3(c+1)$  by Lemma 2
19  |   |   | end
20  |   | end
21  | end
22 return False;
23 // the correctness of the algorithm is ensured by Theorem 1

```

**Algorithm 1:** Algorithm of Theorem 2

## 4 An FPT algorithm for the PTMC problem

We need the following definitions and results before we are ready to prove the main result of this section.

Given a plane graph  $\Gamma$  and a non-negative integer  $k$ , we say that a graph  $\Gamma'$  is a  $k$ -face completion of  $\Gamma$  if it can be obtained from  $\Gamma$  in the following way; for every  $f \in F(\Gamma)$  we add a set  $E_f$  of at most  $k$  edges to  $\Gamma$  such that the endpoints of the edges in  $E_f$  are vertices of  $\Gamma$  that belong to the boundary of  $f$ , all the edges  $E_f$  lie inside  $f$ , they do not intersect  $\Gamma$  in any points other than their endpoints, and finally they do not intersect each other.

Let  $r$  and  $q$  be integers such that  $r \in \mathbb{N}_{\geq 3}, q \in \mathbb{N}_{\geq 1}$ . A  $(r, q)$ -cylinder, denoted by  $C_{r,q}$ , is the Cartesian product of a cycle on  $r$  vertices and a path on  $q$  vertices. We will refer to  $r$  as the *length* and  $q$  as the *width* of  $C_{r,q}$ . Note here that  $C_{r,q}$  is a 3-connected graph and thus, by Whitney's Theorem, it is uniquely embeddable (up to homeomorphism) in the sphere. Furthermore,  $C_{r,q}$  has exactly two non-square faces  $f_1$  and  $f_2$  that are incident only with vertices of degree 3. We call one of the faces  $f_1$  and  $f_2$  the *interior* of  $C_{r,q}$  and the other the *exterior* of  $C_{r,q}$ . We call the vertices incident to the interior (exterior) of  $C_{r,q}$  *base* (*roof*) of  $C_{r,q}$ .

Let  $\Gamma$  be a plane graph. We give the definition of the graph  $\Gamma_{r,q}$  for  $r \in \mathbb{N}_{\geq 3}$  and  $q \in \mathbb{N}_{\geq 3}$ . Let  $f_i \in F(\Gamma)$  and let  $\Theta_1^i, \dots, \Theta_{\rho_i}^i$  be the connected components of  $B_\Gamma(f_i)$ . For each  $\Theta_j^i$ , we denote by  $\sigma_{j,i}$  the length of a facial walk of  $\Theta_j^i$ . We then add a copy  $C_j^i$  of  $(\sigma_{j,i} \cdot r, q)$ -cylinder in the embedding of  $\Gamma$  such that  $\Theta_j^i$  is contained in the interior of  $C_j^i$  and all  $\Theta_1^i, \dots, \Theta_{j-1}^i, \dots, \Theta_{j+1}^i, \dots, \Theta_{\rho_i}^i$  are contained in the exterior of  $C_j^i$ . Then we partition the base of  $C_j^i$  into  $\sigma_{j,i}$  parts  $Q_l, l \in \sigma_{j,i}$

each consisting of  $r$  consecutive base vertices. Let  $(u_{j,i}^1, u_{j,i}^2, \dots, u_{j,i}^{\sigma_{j,i}}, u_{j,i}^1)$  be a facial walk of  $\Theta_{j,i}$ . We join by  $r$  edges the vertex  $u_{j,i}^x$  to all the vertices of the set  $Q_l$ ,  $l \in \sigma_{j,i}$ . We apply this enhancement for each connected component of the boundary of each face of  $\Gamma$  and we denote the resulting graph by  $\widehat{\Gamma}_{r,q}$ .

We call a face  $f_i$  of  $\widehat{\Gamma}_{r,q}$  non-trivial if  $B_{\widehat{\Gamma}_{r,q}}(f_i)$  has more than one connected components  $\Theta_1^i, \dots, \Theta_{\rho_i}^i$ . Notice that if  $f_i$  is non-trivial, each  $\Theta_j^i$  is the roof of some previously added cylinder. For each such cylinder, let  $J_j^i$  be a set of  $r$  consecutive vertices of its roof. We add inside  $f_i$  a copy  $C_{f_i}$  of  $C_{\rho_i, r, q}$  such that its base is a subset of  $f_i$  and let  $\{I_1, \dots, I_{\rho_i}\}$  be a partition of its roof in  $\rho_i$  parts, each consisting of  $r$  consecutive base vertices. For each  $x \in \{1, \dots, \rho_i\}$  we add  $r$  edges each connecting a vertex of  $J_j^i$  with some vertex of  $I_x$  in a way that the resulting embedding remains plane (there is a unique way for this to be done). We apply this enhancement for each non-trivial face of  $\widehat{\Gamma}_{r,q}$  and the resulting graph is the graph  $\Gamma_{r,q}$ . Notice that  $\Gamma_{r,q}$  is not uniquely defined as its definition depends on the choice of the sets  $J_x$ . From now on, we always consider an arbitrary choice for  $\Gamma_{r,q}$  and we call  $\Gamma_{r,q}$  the  $(r, q)$ -cylindrical enhancement of  $\Gamma$ . Finally, given a plane graph  $\Gamma$  and  $r, q \in \mathbb{N}_{\geq 3}$ . Let  $V_{\Gamma, r, q}^0 = V(\Gamma)$  and  $V_{\Gamma, r, q}^n = V(\Gamma_{r,q}) \setminus V(\Gamma)$  and notice that  $\deg_{\Gamma_{r,q}}(v) \leq 4$ , for every  $v \in V_{\Gamma, r, q}^n$ . (For an example, see Figure 3.) Given a positive integer  $k$ , we denote by  $\widetilde{\Gamma}_k$  the graph  $\Gamma_{2, k, 8 \cdot k}$ .




Figure 3: This figure depicts the construction of  $\Gamma_{3,2}$  from  $\Gamma$  ( $\Gamma$  is the graph on the left).

**Lemma 7** ([2]). *Let  $\Gamma$  be a plane graph and let  $f$  be a face of  $\Gamma$ . Let also  $M$  be a plane graph such that  $M \subseteq \mathbf{cl}(f)$ ,  $E(M) \cap E(B_{\Gamma}(f)) = \emptyset$  and  $V(M) \subseteq V(B_{\Gamma}(f))$ . Then there is a closed curve  $K$  in  $f$  meeting each edge of  $M$  twice.*

We define the *dual* of the plane graph  $\Gamma$ , denoted by  $\Gamma' = (V', A')$  in the following way:

1. for every  $f \in F(\Gamma)$ , we add a vertex  $v_f \in V'$  mapped to a point of  $f$ ,
2. for every two faces  $f, f' \in F(\Gamma)$  and every edge  $e \in \mathbf{bd}(f) \cap \mathbf{bd}(f')$  we add an arc  $e_{f, f'} \in A'$  joining  $v_f, v_{f'}$  and meeting  $\Gamma$  only at a single point of  $e$ .

A planar graph  $G$  is *outerplanar* if there exists an embedding  $\Gamma$  in  $\Sigma$  and a face  $f \in F(\Gamma)$  such that every vertex of  $\Gamma$  belongs to  $\mathbf{bd}(f)$ . We define the *weak dual* of such an embedding  $\Gamma$  of an outerplanar graph by removing from its dual the vertex that is contained in the interior of its unique face  $f$  that contains all vertices of  $\Gamma$  in its border.

**Lemma 8.** *Let  $\Gamma$  be a cycle with  $n$  vertices and  $k$  chords such that no two chords share a common endpoint. Let also  $\Gamma$  be embedded on  $\Sigma$  in such a way that if  $K$  is the curve defined by the vertices and the edges of  $\Gamma$  then all the chords of  $\Gamma$  are contained in its interior. Then  $\Gamma$  is an embedded topological minor of the  $(n, k+1)$ -cylinder  $C$  where the vertices of  $\Gamma$  are mapped on vertices of the roof of  $C$  and the paths joining the edges of the cycle contain only vertices and edges of the roof of  $C$ .*



*Proof.* We prove this lemma by induction on  $k$ . Notice that if  $\Gamma$  has at most one chord then the statement holds trivially.

Assume that the lemma holds for every cycle with  $l < k$  chords and let  $\Gamma$  be a cycle with  $k$  chords. Notice that  $\Gamma$  is an outerplanar graph and consider its weak-dual, which by definition is a tree, say  $T$ . Let  $e_i$ ,  $i \in [k]$  be the chords that bound the faces of  $\Gamma$  that correspond to leaves of  $T$ . We call these chords simplicial. From the induction hypothesis  $\Gamma \setminus \{e_i \mid i \in [k]\}$  is a topological minor of the  $(n, k - i + 1)$ -cylinder, say  $C'$ , and the paths joining the edges of  $\Gamma$  contain only vertices and edges of the roof of  $C'$ . Let  $C''$  be the graph obtained from the  $C'$  by subdividing each edge of  $C'$  that has exactly one endpoint in its roof exactly once and add edges between the subdivided edges so as to obtain the  $(n, k - i + 2)$ -cylinder, say  $C$ . Notice then that  $\Gamma \setminus \{e_i \mid i \in [k]\}$  is an embedded topological minor of  $C$  such that the paths joining the edges of  $\Gamma$  contain only vertices and edges of the roof of  $C$ . Finally, since no two chords of  $\Gamma$  share a common endpoint observe that we may find paths joining the simplicial edges  $\{e_i \mid i \in [k]\}$  by using the newly added vertices and edges of  $C$ . By the construction, it is easy to verify that the paths joining the edges of  $\Gamma$  in  $C$  form indeed the model of an embedded topological minor of  $\Gamma$ .  $\square$

**Observation 4.** Let  $r \in \mathbb{N}_{\geq 3}$  and  $q \in \mathbb{N}_{\geq 2}$ . For every plane graph  $\Gamma$ ,  $|V(\Gamma_{r,q})| = \mathcal{O}(|V(\Gamma)|)$  and  $|E(\Gamma_{r,q})| = \mathcal{O}(|E(\Gamma)|)$ .

Let  $G$  be a graph. We say that two paths in  $G$  are *disjoint* if they don't share common internal vertices. Given a collection  $\mathcal{P}$  of pairwise disjoint paths of  $G$ , we define  $L(\mathcal{P}) = \{\{x, y\} \mid x \text{ and } y \text{ are the endpoints of a path in } \mathcal{P}\}$ .

Given a plane graph  $\Gamma$  and an open set  $\Lambda$  of  $\Sigma$ , we define  $\Gamma\langle\Lambda\rangle$  as the graph whose edge set consists of the edges of  $\Gamma$  that are subsets of  $\Lambda$  and whose vertex set consists of their endpoints.

**Proposition 2.** Let  $\Gamma$  be a plane graph,  $k \in \mathbb{N}_{\geq 1}$  and let  $\Gamma^+$  be a  $k$ -face completion of  $\Gamma$ . For every face  $f \in F(\Gamma)$ , there is a collection  $\mathcal{P}$  of  $k$  disjoint paths in the graph  $\tilde{\Gamma}_k\langle f \rangle$  such that  $L(\mathcal{P}) = E(\Gamma^+\langle f \rangle)$ .

*Proof.* Let  $f_i \in F(\Gamma)$  be a face of  $\Gamma$ . By the assumption, there are at most  $k$  edges of  $\Gamma^+$  contained in  $f_i$ . Let  $\{s_q, t_q\}$ ,  $q \in \{1, \dots, k\}$  be those  $k$  edges. From now on we will call them completion edges. Let also  $\Theta_1^i, \Theta_2^i, \dots, \Theta_{\rho_i}^i$  be the connected components of  $B_\Gamma(f_i)$ . Notice that for every completion edge  $e$  either there exists a  $\Theta_j^i$  such that both endpoints of  $e$  are vertices of  $\Theta_j^i$  or there exist disjoint  $\Theta_j^i$  and  $\Theta_{j'}^i$  where each of them contains exactly one of the endpoints of  $e$ . For every edge  $e$  for which there exists an  $\Theta_j^i$  that contains both endpoints of  $e$ , we say that  $e$  is *internal* to  $\Theta_j^i$ . For every edge  $e$  for which there exist disjoint  $\Theta_j^i$  and  $\Theta_{j'}^i$  where each of them contains exactly one of the endpoints of  $e$  we say that  $e$  is an *outgoing* edge of  $\Theta_j^i$  and  $\Theta_{j'}^i$ .

We first consider the case where  $\rho_i \geq 2$ . From Lemma 7, there exists a closed curve  $K_{f_i}$  in  $f_i$  meeting each completion edge twice such that  $\mathbf{int}(K_{f_i})$  contains only points of  $f_i$ . Moreover, notice that for every connected component  $\Theta_j^i$  there exists a closed curve  $K_j^i \subseteq \Sigma \setminus \mathbf{cl}(K_{f_i})$  in  $f$  such that (1)  $\Theta_j^i \subseteq \mathbf{int}(K_j^i)$ , (2)  $K_j^i$  meets each internal edge of  $\Theta_j^i$  exactly twice, (3)  $K_j^i$  meets each outgoing edge of  $\Theta_j^i$  exactly once and (4) for  $j \neq j'$ ,  $K_j^i \cap K_{j'}^i = \emptyset$ . Finally, observe that we may choose the curves  $K_j^i$ ,  $j \in [\rho_i]$ , and  $K_{f_i}$  in such a way that  $K_{f_i} \subseteq \Sigma \setminus \mathbf{cl}(K_j^i)$ ,  $j \in [\rho_i]$ .

At every point in  $\Sigma$  where a completion edge meets a curve  $K_{f_i}$  or  $K_j^i$ ,  $j \in [\rho_i]$  we add a new vertex. For every  $j \in [\rho_i]$ , we denote the set of new vertices on  $K_j^i$  by  $V_j^i$ . We also denote by  $V_{f_i}^i$  the vertices on  $K_{f_i}$ . Note here that by definition  $|V_{f_i}^i|, |V_j^i| \leq 2 \cdot k$ . Notice that every vertex  $v \in V_{f_i}^i$  has exactly one neighbor in  $\bigcup_{j \in [\rho_i]} V_j^i$ . That is, for every vertex

$v \in V_{f_i}^i$  there exists exactly one  $j \in [\rho_i]$  such that  $N(v) \cap V_j^i \neq \emptyset$ . We partition  $V_{f_i}^i$  to the sets  $\tilde{V}_j^i = \{v \in V_{f_i}^i \mid N(v) \cap V_j^i \neq \emptyset\}$ . Finally observe that, due to the planarity of the embedding, for every  $j \in [\rho_i]$  the vertices of  $\tilde{V}_j^i$  appear consecutively in  $K_{f_i}$ .

Recall that for every  $j \in [\rho_i]$ ,  $\Theta_j^i$  defines a closed curve in  $\Sigma$  and let  $A_j = A[\Theta_j^i, K_j^i]$  denote the annulus between the curve defined by  $\Theta_j^i$  and  $K_j^i$ . Without loss of generality we may assume that the cylinder  $C_j^i$  of  $\Gamma_{2 \cdot k, 8 \cdot k}$  is embedded inside  $A_j$  in such a way that  $V_j^i$  are consecutive vertices of the roof of  $C_j^i$  and that  $V_j^i \subseteq J_j^i$ . It is then easy to see that for every edge inside the annulus  $A_j$  (which is part of a completion edge) we may find a path in  $C_j^i$  joining its endpoints in  $\Gamma_k$ .

Let now  $\mathbf{int}(K_{f_i})$  be the open disk of  $\Sigma \setminus K_{f_i}$  that contains only points of the completion edges. Without loss of generality we may also assume that the cylinder  $C_{f_i}$  is embedded inside  $\mathbf{int}(K_{f_i})$  in such a way that  $\tilde{V}_j^i \subseteq I_j$ , for every  $j \in [\rho_i]$ . Observe that the curve  $K_{f_i}$  together with the edges in its interior forms a cycle  $Q$  on  $2 \cdot k$  vertices with  $k$  chords where no two chords share a common endpoint. From Lemma 8,  $Q$  is an embedded topological minor of the  $(2 \cdot k, k + 1)$ -cylinder  $C$  such that all vertices of  $Q$  are mapped on vertices of the roof of  $C$  and the paths joining the edges of the cycle contain only vertices and edges of the roof of  $C$ . Thus, by appropriately subdividing  $C$  we may see that  $Q$  is also a topological minor of  $C_{f_i}$ . Therefore, for the edges in  $\mathbf{int}(K_{f_i})$  there exist vertex disjoint paths in  $C_{f_i}$  joining their endpoints. This concludes the proof that there exists a collection  $\mathcal{P}$  of  $k$  disjoint paths in the graph  $\tilde{\Gamma}_k \langle f_i \rangle$  such that  $L(\mathcal{P}) = E(\Gamma^+ \langle f_i \rangle)$ .

In the special case where  $\rho_i = 1$ , let us consider only the closed curve  $K_{f_i}$  in  $f_i$  that meets each completion edge twice and add a vertex at every point where a completion edge meets a curve  $K_{f_i}$ . Then as above the curve  $K_{f_i}$  together with the edges in its interior forms a cycle  $Q$  on  $2 \cdot k$  vertices with  $k$  chords where no two chords share a common endpoint. Again, from Lemma 8,  $Q$  is an embedded topological minor of the  $(2 \cdot k, k + 1)$ -cylinder  $C$  such that all vertices of  $Q$  are mapped on vertices of the roof of  $C$  and the paths joining the edges of the cycle contain only vertices and edges of the roof of  $C$ . Thus, by appropriately subdividing  $C$  we may see that  $Q$  is also a topological minor of  $C_1^i$ . Therefore, for the edges in  $\mathbf{int}(K_{f_i})$  there exist vertex disjoint paths in  $C_1^i$  joining their endpoints. This concludes the proof that there exists a collection  $\mathcal{P}$  of  $k$  disjoint paths in the graph  $\tilde{\Gamma}_k \langle f_i \rangle$  such that  $L(\mathcal{P}) = E(\Gamma^+ \langle f_i \rangle)$ .  $\square$

From the proof of [2, Theorem 2] we obtain the following lemma.

**Lemma 9.** *Let  $k$  be a positive integer. Let  $\Gamma$  be a plane graph and  $\{(s_i, t_i) \mid i \in [k]\}$  be  $k$  pairs of terminals of  $\Gamma$  such that the graph with vertex set  $\bigcup_{i \in [k]} \{s_i, t_i\}$  and edge set  $\{\{s_i, t_i\} \mid i \in [k]\}$  is connected. Let also  $f$  be a face of  $\Gamma$  and  $\Gamma^+$  be a face completion of  $f$ . If  $\Gamma^+$  contains  $k$  internally vertex disjoint paths joining the terminals  $(s_i, t_i)$ , then there exists a face completion  $\Gamma'$  of  $f$  with at most  $k^{2^k}$  edges.*

We are now ready to prove one of the main results of this section.

**Theorem 3.** *Let  $\Gamma$  and  $\Delta$  be plane graphs where  $\Delta$  is connected and  $k = |E(\Delta)|^{2^{|E(\Delta)|}}$ . There exists a  $k$ -face completion  $\Gamma^+$  of  $\Gamma$  such that  $\Delta \leq_{etm} \Gamma^+$  if and only if  $\Delta \leq_{etm}^S \tilde{\Gamma}_k$  where  $S = V(\tilde{\Gamma}_k) \setminus V(\Gamma) = V_{\Gamma, 2 \cdot k, 8 \cdot k}^n$ .*

*Proof.* Suppose that  $\Delta \leq_{etm} \tilde{\Gamma}_k$ , no vertex of  $\Delta$  is mapped to a vertex of  $V(\tilde{\Gamma}_k) \setminus V(\Gamma)$ , and that  $\mathcal{P}$  is the collection of disjoint paths in  $\tilde{\Gamma}_k$ , corresponding to edges of  $\Delta$ . Let  $\mathcal{P}'$  be family of the maximal (according to the number of vertices) subpaths of the paths in  $\mathcal{P}$  that consist only of edges of  $E(\tilde{\Gamma}_k) \setminus E(\Gamma)$ . Let  $\Gamma^+$  be the graph obtained from  $\tilde{\Gamma}_k$  by removing all vertices

of  $(\tilde{\Gamma}_k \setminus \Gamma)$  that do not belong to any path of  $\mathcal{P}'$  and contracting all paths in  $\mathcal{P}'$  to a single edge. Then by construction,  $\Gamma^+$  is a completion of  $\Gamma$  such that  $\Delta \leq_{etm} \Gamma^+$ .

We now show that if there exists a completion  $\Gamma^+$  of  $\Gamma$  such that  $\Delta \leq_{etm} \Gamma^+$  then  $\Delta \leq_{etm} \tilde{\Gamma}_k$  and no vertex of  $\Delta$  is mapped to a vertex of  $V(\tilde{\Gamma}_k) \setminus V(\Gamma)$ . Let  $\Gamma^+$  be a completion of  $\Gamma$  such that  $\Delta \leq_{etm} \Gamma^+$  where  $|E(\Gamma^+) \setminus E(\Gamma)|$  is minimal. From Lemma 9 we may assume that in each face  $f \in F(\Gamma)$  at most  $k$  edges have been added in  $\Gamma^+$ . Suppose not and let  $f \in F(\Gamma)$  be a face of  $\Gamma$  where more than  $k$  edges have been added in  $\Gamma^+$ . Let also  $\{\{s_i, t_i\} \mid i \in |E(\Delta)|\}$  be the pairs of vertices of  $\Gamma$  corresponding to the endpoints of the edges of  $\Delta$ . Clearly, in  $\Gamma^+$  there exist internally vertex disjoint paths joining the pairs  $\{s_i, t_i\}$ . Let  $\hat{\Gamma}$  be the graph obtained from  $\Gamma^+$  after we remove the completion edges that have been added to the face  $f$  of  $\Gamma$  in it. As  $\Gamma^+$  and  $\hat{\Gamma}$  only differ in the edges of  $f$  and  $f$  is a face of  $\hat{\Gamma}$  there clearly exists a completion of the face  $f$  of  $\hat{\Gamma}$  (that is,  $\Gamma^+$ ) such that there exist  $|E(\Delta)|$  internally vertex disjoint paths joining the pairs  $\{s_i, t_i\}$ . From Lemma 9, there exists a face completion  $\Gamma'$  of  $f$  in  $\hat{\Gamma}$  with at most  $k$  edges. However,  $\Gamma'$  is also a completion of  $\Gamma$  that contains  $|E(\Delta)|$  internally vertex disjoint paths joining the pairs  $\{s_i, t_i\}$ . Moreover, in every face of  $\Gamma$  except  $f$  the same amount of edges as in  $\Gamma^+$  has been added and strictly less edges have been added in  $f$ , a contradiction to the minimality of  $\Gamma^+$ .

Proposition 2 yields that for every face  $f \in F(\Gamma)$  there exist a collection  $\mathcal{P}$  of  $k$  disjoint paths in  $\tilde{\Gamma}_k \setminus f$  such that  $L(\mathcal{P}) = E(\Gamma^+ \setminus f)$ . Therefore, we conclude that  $\Delta \leq_{etm} \tilde{\Gamma}_k$  and by construction no vertex of  $\Delta$  is mapped to a vertex of  $V(\tilde{\Gamma}_k) \setminus V(\Gamma)$ .  $\square$

**Proposition 3** ([11, 10]). *There exists an algorithm  $A_1$  that given a planar graph  $G$  and a non-negative integer  $q$  outputs either a tree decomposition of  $G$  of width at most  $18q$  or a subdivided wall  $W$  of  $G$  of height  $q$  and a tree decomposition  $\mathcal{D}$  of the compass  $K_W$  of  $W$  of width at most  $18q$  in time  $O_q(|E(G)|)$ .*

**Proposition 4** ([11]). *There exists a function  $f : \mathbb{N}_{\geq 1} \rightarrow \mathbb{N}_{\geq 0}$  and an algorithm  $A_2$  that, given a planar graph  $\Gamma$ , an  $k$ -vertex planar graph  $\Delta$ , and a subdivided wall  $W$  as a subgraph of  $\Gamma$  of height  $f(k)$  whose compass has treewidth at most  $\ell$ , outputs an edge  $e = \{u, v\}$  of  $\Gamma$  where  $\deg_{\Gamma}(v), \deg_{\Gamma}(u) \geq 2$  such that  $\Delta \leq_{etm} \Gamma$  if and only if  $\Delta \leq_{etm} \Gamma \setminus \{e\}$ . This algorithm runs in  $O_{k,\ell}(|E(\Gamma)|)$  steps.*

**Lemma 10.** *Let  $\Gamma$  and  $\Delta$  be two plane graphs and  $V \subseteq V(\Gamma)$ , where  $l = \max\{\deg_{\Gamma}(v) \mid v \in V\}$ . There exist two plane graphs  $\Gamma'$  and  $\Delta'$  such that*

- $\deg_{\Gamma'}(v) = 1$ , for every  $v \in V(\Gamma') \setminus V(\Gamma)$ , every edge in  $E(\Gamma') \setminus E(\Gamma)$  has one endpoint in  $\Gamma' \setminus \Gamma$ ,
- $|E(\Gamma')| \leq (2l + 3)|E(\Gamma)|$ ,  $|E(\Delta')| \leq (2l + 3)|E(\Delta)|$ ,  $\mathbf{tw}(\Gamma') = \mathbf{tw}(\Gamma)$ , and
- $\Delta \leq_{etm}^V \Gamma$  if and only if  $\Delta' \leq_{etm}^{\emptyset} \Gamma'$ .

Moreover if  $\tilde{\Gamma}$  is a spanning subgraph of  $\Gamma'$  such that  $\Delta' \leq_{etm} \tilde{\Gamma}$  then  $\Delta \leq_{etm}^V \Gamma \setminus E$ , where  $E = E(\Gamma') \setminus E(\tilde{\Gamma})$ .

*Proof.* For every vertex  $v \in V(\Gamma) \setminus V$  let  $(v_1, v_2, \dots, v_k)$  be the cyclic neighborhood of  $v$ . We add  $(l + 1)k$  new pendant neighbors to  $v$  in such a way that in its new cyclic neighborhood exactly  $(l + 1)$  vertices appear between  $v_i$  and  $v_{i+1}$ ,  $i \in [k]$ , where  $v_{k+1} = v_1$ . Let  $\Gamma'$  denote the new graph.

For every vertex  $u \in V(\Delta)$  let  $(u_1, u_2, \dots, u_p)$  be the cyclic ordering of the neighborhood of  $u$ . We add  $(l + 1)p$  new pendant neighbors to  $u$  in such a way that in the new cyclic ordering exactly  $(l + 1)$  vertices appear between  $u_i$  and  $u_{i+1}$ ,  $i \in [p]$ , where  $u_{p+1} = u_1$ . Let  $\Delta'$  denote the new graph.

Let  $f$  be an embedded topological minor model of  $\Delta$  in  $\Gamma$  where every vertex of  $\Delta$  is mapped to a vertex of  $V(\Gamma) \setminus V$ . Then by using the newly added pendant vertices in  $\Gamma'$  we may extend it to an embedded topological minor model of  $\Delta'$  in  $\Gamma'$ .

Conversely, let  $f$  be an embedded topological minor model of  $\Delta'$  in  $\Gamma'$ . Notice that all vertices of  $V(\Delta)$  in  $\Delta'$  have degree greater than  $l$ . Observe also that by construction all vertices of  $V(\Gamma')$  that are not vertices of  $V(\Gamma) \setminus V$  are the vertices of  $V$  and the newly added pendant vertices and thus, have degree at most  $l$ . This implies that no vertex of  $\Delta$  can be mapped to a vertex of  $V$  in  $\Gamma'$ . It follows that if  $e \in E(\Delta)$  then the path joining its endpoints in the model of  $\Delta'$  in  $\Gamma'$  does not contain any of the newly added pendant edges. Therefore, the restriction of  $f$  to vertices of  $\Delta$  and paths that model edges of  $\Delta$  leads to an embedded topological minor model of  $\Delta$  in  $\Gamma$  where no vertex of  $\Delta$  has been mapped to a vertex of  $V$ .

Let now  $\tilde{\Gamma}$  be a spanning subgraph of  $\Gamma'$ , where  $E = E(\Gamma') \setminus E(\tilde{\Gamma})$ , such that  $\Delta'$  is an embedded topological minor of  $\tilde{\Gamma}$ . Since the degree of every vertex in  $\tilde{\Gamma}$  does not increase after removing edges, as above we obtain that if  $f$  is an embedded topological minor model of  $\Delta$  in  $\tilde{\Gamma}$  no vertex of  $\Delta$  has been mapped to a vertex of  $V$ . Therefore, again, the restriction of  $f$  to vertices of  $\Delta$  and paths that model edges of  $\Delta$  leads to an embedded topological minor model of  $\Delta$  in  $\Gamma \setminus E$  where no vertex of  $\Delta$  has been mapped to a vertex of  $V$ .

Observe that by construction  $|E(\Gamma')| \leq (2l+3)|E(\Gamma)|$  and  $|E(\Delta')| \leq (2l+3)|E(\Delta)|$ . Finally, the fact that  $\mathbf{tw}(\Gamma') = \mathbf{tw}(\Gamma)$  follows from the folklore observation that the addition of pendant vertices on a graph does not increase its treewidth.  $\square$

**Proposition 5.** *Let  $c$  be a fixed positive integer. Let  $\Delta$  be a  $k$ -edge plane graph and  $\Gamma$  be a plane graph with  $\mathbf{tw}(\Gamma) > 18f((2c+3) \cdot k)$ , where  $f : \mathbb{N}_{\geq 1} \rightarrow \mathbb{N}_{\geq 0}$  is the function of Proposition 4 and  $V \subseteq V(\Gamma)$  where  $\deg_{\Gamma}(z) \leq c$ , for every  $z \in V$ . Then, there exists an algorithm  $A$  that, given  $\Gamma$  and  $\Delta$ , outputs an edge  $e = \{u, v\} \in E(\Gamma)$  such that  $\deg_{\Gamma}(v), \deg_{\Gamma}(u) \geq 2$  and  $\Delta \leq_{etm}^V \Gamma$  if and only if  $\Delta \setminus \{e\} \leq_{etm}^V \Gamma$ . This algorithm runs in  $O_k(|E(\Gamma)|)$  steps.*

*Proof.* Observe first that for every edge  $e \in E(\Gamma)$ , if  $\Delta \leq_{etm}^V \Gamma \setminus \{e\}$  then clearly  $\Delta \leq_{etm}^V \Gamma$ . Let  $\Gamma'$  and  $\Delta'$  be the graphs of Lemma 10. Run algorithm  $A_1$  of Proposition 3 with input  $\Gamma'$  and  $f((2c+3) \cdot k)$ . As  $\mathbf{tw}(\Gamma) > 18f((2c+3) \cdot k)$ ,  $A_1$  outputs a subdivided wall  $W$  of height  $f((2c+3) \cdot k)$  that is a subgraph of  $\Gamma'$  and whose compass  $K_W$  has treewidth at most  $18f((2c+3) \cdot k)$ .

Run algorithm  $A_2$  of Proposition 4 with input  $\Gamma', \Delta'$  and  $W$ . The output of  $A_2$  is an edge  $e = \{u, v\} \in E(\Gamma')$  such that  $\Delta' \leq_{etm} \Gamma'$  if and only if  $\Delta' \leq \Gamma' \setminus \{e\}$ . Recall that all edges of  $E(\Gamma') \setminus E(\Gamma)$  have one endpoint of degree 1. Thus,  $e$  is an edge of  $\Gamma$ . We will prove that  $\Delta \leq_{etm}^V \Gamma$  if and only if  $\Delta \leq_{etm}^V \Gamma \setminus \{e\}$ . Observe first that for every edge  $e' \in E(\Gamma)$ , if  $\Delta \leq_{etm}^V \Gamma \setminus \{e'\}$  then clearly  $\Delta \leq_{etm}^V \Gamma$ . Thus, let us assume that  $\Delta \leq_{etm}^V \Gamma$ . Then from Lemma 10,  $\Delta' \leq_{etm} \Gamma'$ . Therefore, from the choice of  $e$ ,  $\Delta' \leq_{etm} \Gamma' \setminus \{e\}$ . Since  $\Gamma' \setminus \{e\}$  is a spanning subgraph of  $\Gamma'$ , again from Lemma 10,  $\Delta \leq_{etm}^V \Gamma \setminus \{e\}$ . Since  $|E(\Gamma')| \leq (2c+3) \cdot |E(\Gamma)|$ , it is clear that the overall algorithm runs in  $O_k(|E(\Gamma)|)$  steps.  $\square$

The proof of the following theorem follows by repetitive applications of the algorithm of Proposition 5.

**Theorem 4.** *There exists an algorithm that given two plane graphs  $\Gamma$  and  $\Delta$  and a set  $V \subseteq V(\Gamma)$  with  $\deg_{\Gamma}(z) \leq c$ , for every  $z \in V$  outputs a graph  $\Gamma'$ , with  $\Gamma' \subseteq_{sp} \Gamma$  and  $\mathbf{tw}(\Gamma') = \mathcal{O}(f(|E(\Delta)|))$ , for some computable function  $f$  such that  $\Delta \leq_{etm}^V \Gamma$  if and only if  $\Delta \leq_{etm}^V \Gamma'$ . This algorithm runs in  $\mathcal{O}_{|E(\Delta)|}(|E(\Gamma)|)$  steps.*

Let  $\Gamma$  be a connected plane graph and  $Z \subseteq V(\Gamma)$ , we define the following pair of vertex and edge structures:

$$\mathbf{G}_{\Gamma,Z} := (\mathbf{d}(\mathbf{p}(\Gamma, R(\Gamma)), Z), \mathbf{e}(\Gamma, R(\Gamma))).$$

Given two connected plane graphs  $\Delta$  and  $\Gamma$  and  $Z \subseteq V(\Gamma)$  we say that  $\mathbf{G}_{\Delta,\emptyset}$  is a *restricted topological minor* of  $\mathbf{G}_{\Gamma,Z}$ , denoted by  $\mathbf{G}_{\Delta,\emptyset} \leq_{rtm} \mathbf{G}_{\Gamma,Z}$ , if and only if there exist two functions  $f_1 : V(R^3(\Delta)) \rightarrow 2^{V(R^3(\Gamma))}$  and  $f_2 : E(R^3(\Delta)) \rightarrow 2^{E(R^3(\Gamma))}$  satisfying the following:

1. for every  $x \in V(\Delta)$ ,  $f_1(x) \in V(\Gamma) \setminus Z$  and  $|f_1(x)| = 1$ ,
2. for every  $x \in \cup_{i \in [3]} V(R_s^i(\Delta))$ ,  $f_1(x) \notin \cup_{i \in [3]} (V(R_r^i(\Gamma)))$  and  $|f_1(x)| = 1$ ,
3. for every  $x, y \in \cup_{i \in [3]} V(R_r^i(\Delta))$  is connected and  $f_1(x) \cap f_1(y) = \emptyset$ ,
4. for every  $xy \in E_s^3(\Delta)$ ,  $G[f_2(xy)]$  is a path between  $f_1(x)$  and  $f_1(y)$  and  $f_2(xy) \subseteq E_s^3(\Gamma)$ , and
5. for every  $xy \in E_r^3(\Delta)$ ,  $G[f_2(xy)]$  is a path between some vertex of  $f_1(x)$  and some vertex of  $f_1(y)$ .

We will use the following three observations:

**Observation 5.** *Let  $\Delta, \Gamma$  and  $\Theta$  be three plane graphs,  $\Theta_s$  be a subdivision of  $\Theta$ , and  $Z \subseteq V(\Gamma)$ . If  $\Delta$  is topologically isomorphic to  $\Theta$  and  $\Theta_s \leq_{es}^Z \Gamma$ , then  $\Delta \leq_{etm}^Z \Gamma$ .*

**Observation 6.** *Let  $\Theta$  be a connected plane graph and  $\Gamma$  be a plane graph. If  $\Theta \leq_{es}^Z \Gamma$ , then  $\mathbf{G}_{\Theta,\emptyset} \leq_{rtm} \mathbf{G}_{\Gamma,Z}$ .*

**Observation 7.** *If a connected plane graph  $\Gamma$  is a subdivision of a connected plane graph  $\Theta$ , then  $\mathbf{G}_{\Theta,\emptyset} \leq_{rtm} \mathbf{G}_{\Gamma,\emptyset}$ .*

**Theorem 5.** *Let  $\Gamma, \Delta$  be two connected plane graphs and  $Z \subseteq V(\Gamma)$ . Then  $\Delta \leq_{etm}^Z \Gamma$  if and only if  $\mathbf{G}_{\Delta,\emptyset} \leq_{rtm} \mathbf{G}_{\Gamma,Z}$ .*

*Proof.* Let us make clear what kind of information about  $\Gamma$  and  $\Delta$  is encoded in the structures  $\mathbf{G}_{\Gamma,Z}$  and  $\mathbf{G}_{\Delta,\emptyset}$ .

We first prove that  $\Delta \leq_{etm}^Z \Gamma$  implies that  $\mathbf{G}_{\Delta,\emptyset} \leq_{rtm} \mathbf{G}_{\Gamma,Z}$ . Suppose that  $\Delta \leq_{etm}^Z \Gamma$ . Then, there exist functions  $\rho_1 : V(\Delta) \rightarrow V(\Gamma)$  and  $\rho_2 : E(\Delta) \rightarrow \mathcal{P}(\Gamma)$  that satisfy conditions (1)-(3) of the definition of embedded topological minors and let  $\Theta$  be the union of all paths in  $\rho_2(E(\Delta))$ . It is clear that  $\Theta$  is connected and is a subdivision of  $\Gamma \langle \rho_2 \rangle$ , that does not contain any vertex in  $Z$ , thus we get, from Lemma 7, that  $\mathbf{G}_{\Gamma \langle \rho_2 \rangle, \emptyset} \leq_{rtm} \mathbf{G}_{\Theta,Z}$ . We also have that  $\Theta \leq_{es}^Z \Gamma$  (we just delete every vertex of  $\Gamma$  that does not belong to any path of  $\rho_2(E(\Delta))$ ) and thus from Lemma 6 that  $\mathbf{G}_{\Gamma \langle \rho_2 \rangle, \emptyset} \leq_{rtm} \mathbf{G}_{\Gamma,Z}$ . Finally, as  $\Delta$  is topologically isomorphic to  $\Gamma \langle \rho_2 \rangle$ , we get that  $\mathbf{G}_{\Delta \langle \rho_2 \rangle, \emptyset} \leq_{rtm} \mathbf{G}_{\Gamma,Z}$ , which is what we wanted to prove.

Now we prove that  $\mathbf{G}_{\Delta \langle \rho_2 \rangle, \emptyset} \leq_{rtm} \mathbf{G}_{\Gamma,Z}$  implies that  $\Delta \leq_{etm}^Z \Gamma$ . For any connected plane graph  $\Lambda$ , let  $V^r(\Lambda) = V_r^1(\Lambda) \cup V_r^2(\Lambda) \cup V_r^3(\Lambda)$  and  $V^s(\Lambda) = V_s^1(\Lambda) \cup V_s^2(\Lambda) \cup V_s^3(\Lambda)$ . Suppose that  $\mathbf{G}_{\Delta \langle \rho_2 \rangle, \emptyset} \leq_{rtm} \mathbf{G}_{\Gamma,Z}$ . Then, there exist two functions  $f_1 : V(H) \rightarrow 2^{V(G)}$  and  $f_2 : E(H) \rightarrow 2^{E(G)}$  that satisfy conditions (1)-(5) of the definition of restricted topological minors, where  $H = R^3(\Delta)$  and  $G = R^3(\Gamma)$ . Let  $\Xi$  be the connected graph obtained by the union of all paths in  $f_2(E(H))$ . Observe that by deleting from  $H$  all vertices in  $V^r(\Delta)$  and by dissolving all vertices in  $V^s(\Delta)$  we obtain the initial plane graph  $\Delta$ . Let  $G'$  be the connected plane graph obtained by deleting from  $G \langle f_2 \rangle$  all vertices in  $V^r(\Gamma)$  and  $G^*$  the connected plane graph obtained by dissolving all vertices of  $V^s(\Gamma)$  in  $G'$ . As  $H$  and  $G \langle f_2 \rangle$  are 3-connected

and thus uniquely embedded, we finally obtain that  $\Delta$  and  $G^*$  are topologically isomorphic. Moreover,  $G'$  is a subdivision of  $G^*$  and  $G' \leq_{es}^Z \Gamma$ , as it is obtained from  $\Gamma$  by deleting every vertex that does not belong to any path of  $f_2(E(\Delta))$ . Thus, from Lemma 5, we get that  $\Delta \leq_{etm}^Z \Gamma$ , which is what we want.  $\square$

**Our algorithm for PTMC.** Let  $\Gamma$  and  $\Delta$  be two plane graphs, where  $\Delta$  is connected. From Theorem 3 we construct a cylindrical enhancement  $\tilde{\Gamma}_k$  of  $\Gamma$ , where the vertices of the set  $S = V_{\Gamma, 2k, 8k}^n$  have degree bounded by a constant such that  $\Delta, \Gamma$  are a yes instance if and only if  $\Delta \leq_{etm}^S \tilde{\Gamma}_k$ . Then, the algorithm of Theorem 3 with inputs  $\tilde{\Gamma}_k, \Delta, S$  outputs a graph  $\Gamma'$  with  $\Gamma' \subseteq_{sp} \Gamma$  and  $\mathbf{tw}(\Gamma') = \mathcal{O}(f(|E(\Delta)|))$ . Moreover, Theorem 5 translates  $\Gamma', \Delta$ , and  $S$  to two structures  $\mathbf{G}_{\Delta, \emptyset}$  and  $\mathbf{G}_{\Gamma, S}$ , for which  $\Delta \leq_{etm}^S \Gamma$  if and only if  $\mathbf{G}_{\Delta, \emptyset} \leq_{rtm} \mathbf{G}_{\Gamma, S}$ . Notice that the relation  $\mathbf{G}_{\Delta, \emptyset} \leq_{rtm} \mathbf{G}_{\Gamma, S}$  can be expressed in Monadic Second Order Logic. Finally, by observing that  $\mathbf{tw}(R^3(\Gamma)) = \mathcal{O}(f(|E(\Delta)|))$  we can employ Courcelle's Theorem [6] to obtain an  $f(|E(\Delta)|) \cdot m^2$  time algorithm, for some computable function  $f$ .

## 5 Extensions

Our approach for the PSC problem can also solve the PLANE INDUCED SUBGRAPH COMPLETION problem, with the same running time, where instead of an embedded subgraph we ask for an embedded *induced subgraph*. The only modification would be at the definition of a facial extension of  $\Delta$  where we would additionally require that every connected graph  $\Delta^+$  contains  $\Delta$  as an *induced* subgraph.

In the PTMC problem the connectivity of  $\Delta$  is only required in the proof of Proposition 3 (that has been moved to the appendix). We would like to remark here that if we disregard the embedding of  $\Delta$  then the Proposition holds for disconnected graphs as well. In this case by modifying the algorithm for PTMC we may obtain an FPT algorithm that given a plane graph  $\Gamma$  and a *planar* graph  $D$  decides whether there exists a face completion of  $\Gamma$ , say  $\Gamma^+$ , such that  $D$  is a rooted topological minor of  $\Gamma$ . That is, each vertex of  $D$  is mapped to a specified vertex of  $\Gamma$ . Notice that this approach also permits us to solve the PLANAR DISJOINT PATHS COMPLETION problem where we allow edge additions inside all faces of  $\Gamma$  (in contrast to [2] where edge additions are allowed only inside a specified face of  $\Gamma$ ).

Finally, with the same cylindrical enhancement that we apply for PTMC and the extra restriction that the sets of vertices of the enhanced graph that are contracted to a vertex of the pattern graph  $\Delta$  contain only vertices of the initial graph we can solve the PLANE MINOR COMPLETION problem. In these last two cases, however, only the existence of an FPT algorithm is verified (since both would be derived by Courcelle's Theorem).

## References

- [1] Isolde Adler, Frederic Dorn, Fedor V. Fomin, Ignasi Sau, and Dimitrios M. Thilikos. Fast minor testing in planar graphs. *Algorithmica*, 64(1):69–84, 2012.
- [2] Isolde Adler, Stavros G. Kolliopoulos, and Dimitrios M. Thilikos. Planar disjoint-paths completion. In *Parameterized and Exact Computation - 6th International Symposium, IPEC 2011, Saarbrücken, Germany, September 6-8, 2011. Revised Selected Papers*, pages 80–93, 2011.
- [3] Ivan Bliznets, Fedor V. Fomin, Marcin Pilipczuk, and Michal Pilipczuk. A subexponential parameterized algorithm for proper interval completion. In *Algorithms - ESA 2014 - 22th*

- Annual European Symposium, Wroclaw, Poland, September 8-10, 2014. Proceedings*, pages 173–184, 2014.
- [4] H.L. Bodlaender. Planar graphs with bounded treewidth. Technical Report RUU-CS-88-14, Department of Information and Computing Sciences, Utrecht University, 1988.
- [5] V. Chvatal. Hamiltonian cycles. In E.L. Lawler, J.K. Lenstra, A.H.G.Rinnooy Kan, and D.B. Shmoys (Eds.), editors, *The Traveling Salesman Problem*, pages 403–429. Wiley, New York, 1985.
- [6] Bruno Courcelle. The monadic second-order logic of graphs III: tree-decompositions, minor and complexity issues. *ITA*, 26:257–286, 1992.
- [7] Reinhard Diestel. *Graph Theory, 4th Edition*, volume 173 of *Graduate texts in mathematics*. Springer, 2012.
- [8] Michael B. Dillencourt. Finding hamiltonian cycles in delaunay triangulations is np-complete. *Discrete Applied Mathematics*, 64(3):207 – 217, 1996.
- [9] Pål Grønås Drange, Fedor V. Fomin, Michal Pilipczuk, and Yngve Villanger. Exploring subexponential parameterized complexity of completion problems. In *31st International Symposium on Theoretical Aspects of Computer Science (STACS 2014), STACS 2014, March 5-8, 2014, Lyon, France*, pages 288–299, 2014.
- [10] Petr A. Golovach, Marcin Jakub Kaminski, Spyridon Maniatis, and Dimitrios M. Thilikos. The parameterized complexity of graph cyclability. In *Algorithms - ESA 2014 - 22th Annual European Symposium, Wroclaw, Poland, September 8-10, 2014. Proceedings*, pages 492–504, 2014.
- [11] Marcin Kaminski and Dimitrios M. Thilikos. Contraction checking in graphs on surfaces. In *29th International Symposium on Theoretical Aspects of Computer Science, STACS 2012, February 29th - March 3rd, 2012, Paris, France*, pages 182–193, 2012.
- [12] Haim Kaplan, Ron Shamir, and Robert Endre Tarjan. Tractability of parameterized completion problems on chordal, strongly chordal, and proper interval graphs. *SIAM J. Comput.*, 28(5):1906–1922, 1999.
- [13] Bojan Mohar and Carsten Thomassen. *Graphs on Surfaces*. Johns Hopkins series in the mathematical sciences. Johns Hopkins University Press, 2001.
- [14] Yngve Villanger, Pinar Heggernes, Christophe Paul, and Jan Arne Telle.
- [15] Hassler Whitney. Congruent graphs and the connectivity of graphs. *American Journal of Mathematics*, 54(1):pp. 150–168, 1932.
- [16] M. Yannakakis. Computing the minimum fill-in is np-complete. *SIAM Journal on Algebraic Discrete Methods*, 2(1):77–79, 1981.