THE CONNECTING SOLUTION OF THE PAINLEVÉ PHASE TRANSITION MODEL

MARCEL G. CLERC, MICHAŁ KOWALCZYK, AND PANAYOTIS SMYRNELIS

ABSTRACT. The second Painlevé O.D.E. $y'' - xy - 2y^3 = 0$, $x \in \mathbb{R}$, is known to play an important role in the theory of integrable systems, random matrices, Bose-Einstein condensates and other problems. The generalized second Painlevé equation $\Delta y - x_1y - 2y^3 = 0$, $(x_1, x_2) \in \mathbb{R}^2$, is obtained by multiplying by $-x_1$ the linear term u of the Allen-Cahn equation $\Delta u = u^3 - u$. It involves a non autonomous potential $H(x_1, y)$ which is bistable for every fixed $x_1 < 0$, and thus describes as the Allen-Cahn equation a phase transition model. The scope of this paper is to construct a solution y connecting along the vertical direction x_2 , the two branches of minima of H parametrized by x_1 . This solution plays a similar role that the heteroclinic orbit for the Allen-Cahn equation. It is the the first to our knowledge solution of the Painlevé P.D.E. both relevant from the applications point of view (liquid crystals), and mathematically interesting.

1. The Allen-Cahn and Painlevé phase transition models

A standard phase transition model is given by the Allen-Cahn O.D.E.:

(1.1)
$$u'' = u^3 - u, \qquad \text{in } \mathbb{R},$$

that can alternatively be written u'' = W'(u), where $W(u) = \frac{1}{4}(u^2 - 1)^2$ is a double well potential. In this model, u describes the mass fraction of the two phases of a substance (e.g. an alloy), and takes values approximately +1 or -1 for the pure phases. Equation (1.1) has variational structure. Let

$$E_{\rm AC}(u,(a,b)) := \int_a^b \left(\frac{1}{2}|u'|^2 + \frac{1}{4}(u^2 - 1)^2\right)$$

be the Allen-Cahn energy associated to (1.1). To minimize E_{AC} the right balance between the contributions of the kinetic energy $\frac{1}{2}|u'|^2$ and the potential should be achieved. On the one hand the term $\frac{1}{2}|u'|^2$ penalizes high variations of u, while on the other hand the potential term W forces the minimizer to be close to its global minima ± 1 . It is clear that the trivial solutions ± 1 are the two global minimizers of E_{AC} . Thus, it is more relevant to investigate instead, the existence of *local minimizers* which are also called *minimal* solutions. While solutions of (1.1) are critical points of E_{AC} , a minimal solution u of (1.1) satisfies the stronger condition:

$$E_{\rm AC}(u, \operatorname{supp} \phi) \le E_{\rm AC}(u + \phi, \operatorname{supp} \phi),$$

for all $\phi \in C_0^{\infty}(\mathbb{R})$ (i.e. any perturbation with compact support of u has greater or equal energy). It turns out that up to translations and change of x by -x, the only minimal solution of (1.1) is the heteroclinic orbit $\eta(x) = \tanh(x/\sqrt{2})$, connecting at $\pm \infty$ the two phases ± 1 .

A much more challenging problem is the description of all bounded solutions of the Allen-Cahn P.D.E.:

(1.2)
$$\Delta u = u^3 - u, \quad \text{in } \mathbb{R}^n,$$

which is associated to the functional $E_{AC}(u, \Omega) := \int_{\Omega} \left(\frac{1}{2} |\nabla u|^2 + \frac{1}{4} (u^2 - 1)^2\right)$ (where $\Omega \subset \mathbb{R}^n$ is bounded). De Giorgi in 1978 [17] suggested a striking analogy with minimal surface theory that led to significant developments in P.D.E. and the Calculus of Variations, by stating the following conjecture about bounded solutions on \mathbb{R}^n :

Conjecture (De Giorgi). Let $u \in C^2(\mathbb{R}^n)$ be a solution to (1.2) such that

- (i) |u| < 1,
- (ii) $\frac{\partial u}{\partial x_n} > 0$ for all $x \in \mathbb{R}^n$.

Is it true that all the level sets of u are hyperplanes, at least for $n \leq 8$?

¹⁹⁹¹ Mathematics Subject Classification. Primary 35J91; 35J20; Secondary 35B40; 35B06; 35B25.

The relationship with the Bernstein problem for minimal graphs is the reason why $n \leq 8$ appears in the conjecture. We refer to the expository papers of Farina and Valdinoci [18], and Savin [28] for a detailed account. The conjecture was proved by Ghoussoub and Gui in [19] for n = 2, for n = 3 by Ambrosio and Cabré in [7] and for $4 \leq n \leq 8$ by Savin in [27] under the extra requirement that

(1.3)
$$\lim_{x_n \to \pm \infty} u(x_1, \dots, x_n) = \pm 1.$$

If we drop the monotonicity requirement as well as (1.3) and simply ask about the structure of minimal solutions¹ of (1.2), then we know from [27] that, for $n \leq 7$ any minimal solution u of (1.2) is either trivial i.e. $u \equiv \pm 1$ or one dimensional i.e. $u(x) = \eta((x - x_0) \cdot \nu)$, for some $x_0 \in \mathbb{R}^n$, and some unit vector $\nu \in \mathbb{R}^n$. Thus the heteroclinic orbit η of O.D.E. (1.1) plays a crucial role for entire solutions of P.D.E. (1.2).

In order to construct other connecting solutions of (1.2), one shall impose some additional requirements. For instance, when n = 2, (1.2) admits a unique *saddle* solution u (cf. [16]) satisfying the following properties:

- $u(x_1, x_2)$ has the same sign as the product $x_1 x_2$,
- u is odd with respect to x_1 and x_2 ,
- $\lim_{x_1\to\infty} u(x_1, x_2) = \eta(x_2)$, and $u_{x_1}(x_1, x_2) > 0$, $\forall x_2 > 0$,
- $\lim_{\lambda \to \infty} u(\lambda \cos \theta, \lambda \sin \theta) = 1, \forall \theta \in (0, \pi/2).$

This example also outlines the hierarchical structure of (1.2), since by taking the limit of a solution along certain directions at infinity, lower dimensional solutions are obtained (cf. [6, Chapter 8]). For more examples of connecting maps under symmetry assumptions or in the vector case, we refer to [6, Chapters 6, 7, 9] and the references therein (in particular [4] and [29]).

The second Painlevé O.D.E.:

(1.4)
$$y'' - xy - 2y^3 = 0, \qquad x \in \mathbb{R},$$

is basically obtained by multiplying the linear term in the right hand side of (1.1) by -x. Alternatively, we can write (1.4) as

(1.5)
$$y'' = H_y(x, y), \qquad x \in \mathbb{R}$$

with $H(x, y) := \frac{1}{2}xy^2 + \frac{1}{2}y^4$. In contrast with W (defined below (1.1)), the potential H is non autonomous i.e. it depends both on x and y.

Equation (1.4) is known to play an important role in the theory of integrable systems [1], random matrices [15, 20, 11], Bose-Einstein condensates [2, 3, 24, 30] and other problems [5, 23, 25]. Recently [12] it has been shown that when the right hand side of (1.4) is allowed to be a constant $\alpha \in \mathbb{R}$ then it describes local profiles of the so-called *shadow kink* in the theory of light-matter interaction of nematic liquid crystals (cf. also [32]. [31]). In [8, 13, 14] further relation between other types of non topological defects (*shadow vortices, shadow domain walls*) and the generalized Painlevé equation

(1.6)
$$\Delta y - x_1 y - 2y^3 = 0, \qquad \forall x = (x_1, x_2) \in \mathbb{R}^2$$

was established showing that their local structure is determined by special solutions of (1.6). One of the characteristics of these solutions is that they should be entire, another is that they should be *minimal*. To explain what this means, let $\Omega \in \mathbb{R}^2$ be a bounded subset of \mathbb{R}^2 and

$$E_{\mathrm{P}_{\mathrm{II}}}(u,\Omega) = \int_{\Omega} \left[\frac{1}{2} |\nabla u|^2 + \frac{1}{2} x_1 u^2 + \frac{1}{2} u^4 \right],$$

be the functional associated to the generalized second Painlevé equation. By definition a solution of (1.6) is minimal if

(1.7)
$$E_{\mathrm{P}_{\mathrm{II}}}(y, \operatorname{supp} \phi) \le E_{\mathrm{P}_{\mathrm{II}}}(y + \phi, \operatorname{supp} \phi)$$

for all $\phi \in C_0^{\infty}(\mathbb{R}^2)$. This notion of minimality is standard for many problems in which the energy of a localized solution is actually infinite due to non compactness of the domain. The study of minimal solutions of (1.4) has been recently initiated in [12] where we have showed that the Hastings-McLeod solution, denoted

¹Again, we say that the solution u is minimal if $E_{AC}(u, \operatorname{supp} \phi) \leq E_{AC}(u + \phi, \operatorname{supp} \phi)$, for all $\phi \in C_0^{\infty}(\mathbb{R}^n)$.

in this paper by h, is, up to the sign change, the only minimal solution which is bounded at $+\infty$. We recall (cf. [22]) that $h : \mathbb{R} \to \mathbb{R}$ is positive, strictly decreasing (h' < 0) and such that

(1.8)
$$h(x) \sim Ai(x), \qquad x \to \infty,$$
$$h(x) \sim \sqrt{|x|/2}, \qquad x \to -\infty$$

Clearly, the asymptoic behaviour of h is determined by the location of the global minima of the potential H(x, y) associated to the equation (1.4). Indeed for x fixed, H attains its global minimum equal to 0 when y = 0 and $x \ge 0$, and equal to $-\frac{x^2}{8}$ when $y = \pm \sqrt{|x|/2}$ and x < 0. Thus, the global minima of H bifurcate from the origin, and the two minimal solutions $\pm h$ of (1.4) interpolate these two branches of minima.

Equation (1.6) or equivalently $\Delta y = H_y(x_1, y)$, with $x = (x_1, x_2) \in \mathbb{R}^2$ and $H(x_1, y) := \frac{1}{2}x_1y^2 + \frac{1}{2}y^4$ (cf. the expression of $E_{\text{P}_{\text{II}}}$), involves a non autonomous potential which is bistable for every fixed $x_1 < 0$. Hence the Painlevé generalized equation (1.6) describes as the Allen-Cahn equation a phase transition model. For the later the phase transition connects the two minima ± 1 while for the former the phase transition connects the two minima ± 1 while for the former the phase transition connects the two branches $\pm \sqrt{(-x_1)^+/2}$ of minima of H parametrized by x_1 . Note that in the Painlevé model the phase transition occurs only in the P.D.E. case, i.e. when the domain is $\mathbb{R}^{n+1} = \mathbb{R} \times \mathbb{R}^n$ with $n \geq 1$. The scope of this paper is to construct a solution y of (1.6) connecting as $x_2 \to \pm \infty$ and x_1 is fixed, the two branches of minima of H (cf. Theorem 1.1 below). It is the first to our knowledge example of solution of the generalized Painlevé equation both relevant from the applications point of view and mathematically interesting. The solution y has similar properties as the heteroclinic orbit η : it is odd and monotonous with respect to x_2 , and as $x_1 \to -\infty$ its rescaled profile is actually given by η . After the statement of Theorem 1.1, we will further discuss its similarities with the heteroclinic orbit η .

Theorem 1.1. There exists a solution $y : \mathbb{R}^2 \to \mathbb{R}$ to

(1.9)
$$\Delta y - x_1 y - 2y^3 = 0, \qquad \forall x = (x_1, x_2) \in \mathbb{R}^2,$$

such that

- (i) y is positive in the upper-half plane and odd with respect to x_2 i.e. $y(x_1, x_2) = -y(x_1, -x_2)$.
- (ii) y and its derivatives are bounded in the half-planes $[s_0, \infty) \times \mathbb{R}, \forall s_0 \in \mathbb{R}$.
- (iii) y is minimal with respect to perturbations $\phi \in C_0^{\infty}(\mathbb{R}^2)$ such that $\phi(x_1, x_2) = -\phi(x_1, -x_2)$.
- (iv) $\frac{|y(x_1,x_2)|}{Ai(x_1)} = O(1)$, as $x_1 \to \infty$ (uniformly in x_2).

(v) For every
$$x_2 \in \mathbb{R}$$
 fixed, let $\tilde{y}(t_1, t_2) := \frac{\sqrt{2}}{(-\frac{3}{2}t_1)^{\frac{1}{3}}} y \Big(-(-\frac{3}{2}t_1)^{\frac{2}{3}}, x_2 + t_2(-\frac{3}{2}t_1)^{-\frac{1}{3}} \Big)$. Then
 $\Big(\tanh(t_2/\sqrt{2}) \quad \text{when } x_2 = 0.$

(1.10)
$$\lim_{l \to -\infty} \tilde{y}(t_1 + l, t_2) = \begin{cases} \tanh(t_2/\sqrt{2}) & \text{when } x_2 = 0, \\ 1 & \text{when } x_2 > 0, \\ -1 & \text{when } x_2 < 0, \end{cases}$$

for the $C^1_{\text{loc}}(\mathbb{R}^2)$ convergence.

- (vi) $y_{x_1}(x_1, x_2) < 0, \forall x_1 \in \mathbb{R}, \forall x_2 > 0.$
- (vii) $y_{x_2}(x_1, x_2) > 0$, $\forall x_1, x_2 \in \mathbb{R}$, and $\lim_{l \to \pm \infty} y(x_1, x_2 + l) = \pm h(x_1)$ in $C^2_{\text{loc}}(\mathbb{R}^2)$, where h is the Hastings-McLeod solution of (1.4).

The solution provided by Theorem 1.1 has a form of a quadruple connection between the Airy function Ai, the two one dimensional Hastings-McLeod solutions $\pm h$, and the heteroclinic orbit η of the one dimensional Allen-Cahn equation. Comparing (iv) with (1.8) we see that as $x_1 \to \infty$ the function $y(x_1, x_2)$ behaves similarly as the Hastings-McLeod solution $h(x_1)$. At the same time, as $x_2 \to \pm \infty$ we have $y(x_1, x_2) \to \pm h(x_1)$, $x_2 \to \pm \infty$. Perhaps the most interesting aspect of the above solution y is stated in property (v), since after rescaling we obtain as $x_1 \to -\infty$, the convergence to the heteroclinic orbit $\eta(x) = \tanh(x/\sqrt{2})$ of the Allen-Cahn O.D.E. (1.1). In the proof of Theorem 1.1 it is shown that a minimal solution of (1.9) rescaled as in (v), converges as $x_1 \to -\infty$ to a minimal solution of (1.2). This deep connection of the structure of the Painlevé equation with the Allen-Cahn P.D.E., suggests that several properties of the Allen-Cahn equation should be transfered to the Painlevé equation. Although by construction the solution y is only minimal for odd perturbations, we expect that y is actually minimal for general perturbations, and plays a similar role that the heteroclinic orbit for the Allen-Cahn equation. What's more the two global minimizers ± 1 of the functional $E_{\rm AC}$ have their counterparts in the two minimal solutions $\pm h$ of the Painlevé equation. Indeed, property (vii) establishes that y connects monotonically along the vertical direction x_2 , the two minimal solutions $\pm h(x_1)$, in the same way that η connects monotonically the two global minimizers ± 1 . While η is a one dimensional object, the solution $y(x_1, x_2)$ is two dimensional, since x_1 parametrizes the branches of minima of the potential H, and only x_2 is involved in the phase transition.

We believe that in higher dimension $y : \mathbb{R}^{n+1} \to \mathbb{R}, (n \ge 1)$ the structure of solutions of (1.9) exactly mirrors that of (1.2), and going further, one may ask: is it true that that in dimension $n \le 7$, any minimal solution $Y : \mathbb{R}^{n+1} \to \mathbb{R}$ of (1.9) is either $Y(x_1, x_2, \ldots, x_{n+1}) = \pm h(x_1)$ or $Y(x_1, x_2, \ldots, x_{n+1}) = y(x_1, (x_2, \ldots, x_{n+1}) \cdot \mathbf{n} + b)$, for some constant $b \in \mathbb{R}$, and some unit vector $\mathbf{n} \in \mathbb{S}^{n-1}$?

Acknowledgments. M. G. Clerc was partially supported by Fondecyt 1180903. M. Kowalczyk was partially supported by Chilean research grants Fondecyt 1130126 and 1170164, Fondo Basal AFB170001 CMM-Chile. P. Smyrnelis was partially supported by Fondo Basal AFB170001 CMM-Chile and Fondecyt postdoctoral grant 3160055.

2. Odd minimizers of the Ginzburg-Landau type energy

We consider the energy functional

(2.1)
$$E(u) = \int_{\mathbb{R}^2} \frac{\epsilon}{2} |\nabla u|^2 - \frac{1}{2\epsilon} \mu(x) u^2 + \frac{1}{4\epsilon} u^4,$$

where $u \in H^1(\mathbb{R}^2)$ and $\epsilon > 0$ is a small parameters. We suppose that $\mu \in C^{\infty}(\mathbb{R}^2)$ is radial i.e. $\mu(x) = \mu_{rad}(|x|)$, with $\mu_{rad} \in C^{\infty}(\mathbb{R})$ an even function. In addition we assume that

(2.2)
$$\mu \in L^{\infty}(\mathbb{R}^2), \ \mu'_{\text{rad}} < 0 \text{ in } (0,\infty), \text{ and } \mu_{\text{rad}}(\rho) = 0 \text{ for a unique } \rho > 0,$$

In the physical context described in [8] the function μ is specific

$$\mu(x) = e^{-|x|^2} - \chi$$
, with some $\chi \in (0, 1)$, $f(x) = -\frac{1}{2} \nabla \mu(x)$,

but this particular form is irrelevant here. The Euler-Lagrange equation of E is

(2.3)
$$\epsilon^2 \Delta u + \mu(x)u - u^3 = 0, \qquad x = (x_1, x_2) \in \mathbb{R}$$

and we also write its weak formulation:

(2.4)
$$\int_{\mathbb{R}^2} -\epsilon^2 \nabla u \cdot \nabla \psi + \mu u \psi - u^3 \psi = 0, \qquad \forall \psi \in H^1(\mathbb{R}^2)$$

where \cdot denotes the inner product in \mathbb{R}^2 . Note that due to the radial symmetry of μ the energy (2.1) and equation (2.3) are invariant under orthogonal transformations in the domain, and sign change in the range. Our strategy to construct the solution of (1.9) enjoying the properties described in Theorem 1.1 is to find first an *odd* with respect to x_2 minimizer u_{ϵ} of E and then scaling and passing to the limit $\epsilon \searrow 0$ recover y - this gives us existence. Second, in section 3 we show all the properties of y stated in Theorem 1.1.

We explain, formally at the moment, the relation between (1.9) and the energy E. Looking at the energy density of E it is evident that as $\epsilon \to 0$ the modulus of the global or odd minimizer u_{ϵ} should approach a nonnegative root of the polynomial

$$-\mu(x)z + z^3 = 0,$$

or in other words, $|u_{\epsilon}| \to \sqrt{\mu^+}$ as $\epsilon \to 0$ in some, perhaps weak, sense. This function, called the Thomas-Fermi limit of the minimizer is not in $H^1(\mathbb{R}^2)$ and therefore the transition near the set $\mu(x) = 0$ has to be mediated somehow. To see this let us consider a point ξ such that $\mu(\xi) = 0$. By (2.2) $\xi = \rho e^{i\theta}$. At ξ introduce local orthogonal frame $(e^{i\theta}, ie^{i\theta})$ and coordinates $s = (s_1, s_2)$ associated with it. Let u_{ϵ} be any solution of (2.3) and

$$z(s) = e^{-1/3}u(\xi + e^{2/3}s).$$

Noting that $\mu(\xi + \epsilon^{2/3}s) = \epsilon^{2/3}s_1\mu_1 + \dots$ with $\mu_1 < 0$ we get that z satisfies

$$\Delta_s z + s_1 \mu_1 z - z^3 = o(1), \quad \text{as } \epsilon \searrow 0.$$

The equation on the left becomes the second Painlevé equation after passing to the limit and suitable scaling. Now, suppose that u_{ϵ} is the odd minimizer of E, i.e. $u_{\epsilon}(x_1, x_2) = -u_{\epsilon}(x_1, -x_2)$. Except for the points $\bar{x} = (\pm \rho, 0)$ the limiting function z could be one of the Hastings-McLeod one dimensional solutions. However, at $(\pm \rho, 0)$ we should have $z(s_1, s_2) = -z(s_1, -s_2)$, which means that z genuinely depends on both variables. This is the idea behind the proof of the existence part in Theorem 1.1. Showing properties of the solution is a different story and depends on rather tricky application of the moving plane method.

Our first purpose in this paper is to study qualitative properties of the global minimizers of E as $\epsilon \searrow 0$. In our previous work [12] we studied the following energy

$$E(u,\mathbb{R}) = \int_{\mathbb{R}} \frac{\epsilon}{2} |u_x|^2 - \frac{1}{2\epsilon} \mu(x)u^2 + \frac{1}{4\epsilon} |u|^4 - af(x)u, \quad u \colon \mathbb{R} \to \mathbb{R},$$

where $a \ge 0$ is a parameter and $f = -\frac{1}{2}\mu'$, and in [13] we studied its analog for maps $u: \mathbb{R}^2 \to \mathbb{R}^2$.

By proceeding as in [13], one can see that under the above assumptions there exists a global minimizer v of E in $H^1(\mathbb{R}^2)$, namely that $E(v) = \min_{H^1(\mathbb{R}^2)} E$. In addition, we show that v is a classical solution of (2.3), and v is radial. Similarly, in the class $H^1_{\text{odd}}(\mathbb{R}^2) := \{u \in H^1(\mathbb{R}^2) : u(x_1, x_2) = -u(x_1, -x_2)\}$ of odd functions with respect to x_2 , there exists an odd minimizer u which also solves (2.3) and satisfies $u(x_1, x_2) = u(-x_1, x_2)$. Although in the sequel we will focus on the odd minimizer for completeness we chose to present our next result in a slightly more general framework.

Theorem 2.1. For $\epsilon \ll 1$ let u_{ϵ} be a solution of (2.3) converging to 0 as $|x| \to \infty$ (which may be the odd or global minimizer). Let $\rho > 0$ be the zero of μ_{rad} and let $\mu_1 := \mu'_{rad}(\rho) < 0$. For every $\xi = \rho e^{i\theta}$, we consider the local coordinates $s = (s_1, s_2)$ in the basis ($e^{i\theta}$, $ie^{i\theta}$), and the rescaled functions:

(2.5)
$$w_{\epsilon}(s) = 2^{-1/2} (-\mu_1 \epsilon)^{-1/3} u_{\epsilon} \left(\xi + \epsilon^{2/3} \frac{s}{(-\mu_1)^{1/3}}\right).$$

As $\epsilon \to 0$, the function w_{ϵ} converges in $C^2_{\text{loc}}(\mathbb{R}^2)$ up to subsequence, to a function y bounded in the half-planes $[s_0, \infty) \times \mathbb{R}$, for every $s_0 \in \mathbb{R}$, which is a solution of

(2.6)
$$\Delta y(s) - s_1 y(s) - 2y^3(s) = 0, \qquad \forall s = (s_1, s_2) \in \mathbb{R}^2.$$

In particular, if we take u_{ϵ} to be the odd minimizer of E and $\xi = (\pm \rho, 0)$, then the solution y satisfies $y(s_1, s_2) = -y(s_1, -s_2)$, and is minimal with respect to perturbations $\phi \in C_0^{\infty}(\mathbb{R}^2)$, $\phi(s_1, s_2) = -\phi(s_1, -s_2)$. On the other hand, if we take u_{ϵ} to be the global minimizer then $y(s_1, s_2) = h(s_1)$ or $y(s_1, s_2) = -h(s_1)$.

We observe that as a corollary of [14, Theorem 1.1.] it can be proven that $|v_{\epsilon}| \rightarrow \sqrt{\mu^+}$ in $C_{\text{loc}}^0(D(0;\rho))$. Because of the analogy between the functional E and the Gross-Pitaevskii functional in the theory of Bose-Einstein condensates we will call $\sqrt{\mu^+}$ the Thomas-Fermi limit of v_{ϵ} . Theorem 2.1 gives account on how non smoothness of the limit of v_{ϵ} is mediated near the circumference $|x| = \rho$, where μ changes sign, through the solution of (2.6). We should mention here that detailed description of the minimizers for yet more general setting of the energy can be found in [13, 14].

Before proving the theorem we gather general results for minimizers and solutions that are valid for any values of the parameters $\epsilon > 0$. For the rest of this paper v or v_{ϵ} will be the global minimizer and u or u_{ϵ} will be the odd minimizer or a critical point of E. We first prove the existence of global and odd minimizers.

Lemma 2.2. For every $\epsilon > 0$ there exists $v \in H^1(\mathbb{R}^2)$ such that $E(v) = \min_{H^1(\mathbb{R}^2)} E$. As a consequence, v is a C^{∞} classical solution of (2.3). Moreover, for $\epsilon \ll 1$ the global minimizer v is unique up to change of v by -v, and it can be written as $v(x) = v_{rad}(|x|)$, with $v_{rad} \in C^{\infty}(\mathbb{R})$, positive, even, and such that $\lim_{\infty} v_{rad} = 0$.

Proof. We proceed as in [13, Lemma 2.1] to establish that the global minimizer exists and is a smooth solution of (2.3) converging to 0 as $|x| \to \infty$. Next, we notice that $v \neq 0$ for $\epsilon \ll 1$. Indeed, by choosing a test function $\psi \neq 0$ supported in $D(0; \rho) \cap \{x_2 > 0\}$, and such that $\psi^2 < 2\mu$, one can see that

$$E(\psi) = \frac{\epsilon}{2} \int_{\mathbb{R}^2} |\nabla \psi|^2 + \frac{1}{4\epsilon} \int_{\mathbb{R}^2} \psi^2(\psi^2 - 2\mu) < 0, \qquad \epsilon \ll 1.$$

Let $x_0 \in \mathbb{R}^2$ be such that $v(x_0) \neq 0$. Without loss of generality we may assume that $v(x_0) > 0$. Next, consider $\tilde{v} = |v|$ which is another global minimizer and thus another solution. Clearly, in a neighborhood of x_0 we

have v = |v|, and as a consequence of the unique continuation principle (cf. [26]) we deduce that $v \equiv \tilde{v} \geq 0$ on \mathbb{R}^2 . Furthermore, the maximum principle implies that v > 0, since $v \neq 0$. To prove that v is radial we consider the reflection with respect to the line $x_1 = 0$. We can check that $E(v, \{x_1 > 0\}) = E(v, \{x_1 < 0\})$, since otherwise by even reflection we can construct a map in H^1 with energy smaller than v. Thus, the map $\tilde{v}(x) = v(|x_1|, x_2)$ is also a minimizer, and since $\tilde{v} = v$ on $\{x_1 > 0\}$, it follows by unique continuation that $\tilde{v} \equiv v$ on \mathbb{R}^2 . Repeating the same argument for any line of reflection, we deduce that v is radial. To complete the proof, it remains to show the uniqueness of v up to change of v by -v. Let \tilde{v} be another global minimizer such that $\tilde{v} > 0$, and $\tilde{v} \neq v$. Choosing $\psi = u$ in (2.4), we find for any solution $u \in H^1(\mathbb{R}^2)$ of (2.3) the following alternative expression of the energy:

(2.7)
$$E(u) = -\int_{\mathbb{R}^2} \frac{u^4}{4\epsilon}.$$

Formula (2.7) implies that v and \tilde{v} intersect for |x| = r > 0. However, setting

$$w(x) = \begin{cases} v(x) & \text{for } |x| \le r\\ \tilde{v}(x) & \text{for } |x| \ge r, \end{cases}$$

we can see that w is another global minimizer, and again by the unique continuation principle we have $w \equiv v \equiv \tilde{v}$. This completes the proof of Lemma 2.2.

On the other hand, in the class $H^1_{\text{odd}}(\mathbb{R}^2) := \{u \in H^1(\mathbb{R}^2) : u(x_1, x_2) = -u(x_1, -x_2)\}$ of odd functions with respect to x_2 , there exists an odd minimizer with the following properties:

Lemma 2.3. For every $\epsilon > 0$ there exists $u \in H^1_{\text{odd}}(\mathbb{R}^2)$ such that $E(u) = \min_{H^1_{\text{odd}}(\mathbb{R}^2)} E$. As a consequence, u is a C^{∞} classical solution of (2.3). Moreover

- (i) $u(x) \to 0$ as $|x| \to \infty$,
- (ii) $u(x_1, x_2) = u(-x_1, x_2),$
- (iii) up to transformation $u \mapsto -u$ we have $u(x_1, x_2) > 0$, $\forall (x_1, x_2) \in \mathbb{R} \times (0, \infty)$, provided that $\epsilon \ll 1$.

Proof. The existence of $u \in H^1_{\text{odd}}(\mathbb{R}^2)$ such that $E(u) = \min_{H^1_{\text{odd}}(\mathbb{R}^2)} E$, follows as in [13, Lemma 2.1], and clearly u is a critical point of E in the subspace $H^1_{\text{odd}}(\mathbb{R}^2)$. In view of the radial symmetry of μ it is easy to see that the Euler-Lagrange equation (2.4) holds also for every $\phi \in H^1(\mathbb{R}^2)$, such that $\phi(x_1, x_2) = \phi(x_1, -x_2)$. As a consequence, u is a C^{∞} classical solution of (2.3).

For the proof of (i) we refer to [13, Lemma 2.1]. To show that $u(x_1, x_2) = u(-x_1, x_2)$, we first note that $E(u, [0, \infty) \times \mathbb{R}) = E(u, (-\infty, 0] \times \mathbb{R})$. Indeed, if we assume without loss of generality that $E(u, [0, \infty) \times \mathbb{R}) < E(u, (-\infty, 0] \times \mathbb{R})$, the function

(2.8)
$$\tilde{u}(x_1, x_2) = \begin{cases} u(x_1, x_2) & \text{when } x_1 \ge 0, \\ u(-x_1, x_2) & \text{when } x_1 \le 0, \end{cases}$$

has strictly less energy than u, which is a contradiction. Thus, $E(u, [0, \infty) \times \mathbb{R}) = E(u, (-\infty, 0] \times \mathbb{R})$, and as a consequence the function \tilde{u} is also an odd minimizer and a solution. It follows by unique continuation [26] that $\tilde{u} \equiv u$, that is, $u(x_1, x_2) = u(-x_1, x_2)$.

Now, it remains to establish the uniqueness of the odd minimizer u, when $\epsilon \ll 1$. Proceeding as in Lemma 2.2, we can see that $u \neq 0$ for $\epsilon \ll 1$, and that either u > 0 or u < 0 on $\mathbb{R} \times (0, \infty)$. Assume that u_1 and u_2 are two minimizers of E in $H^1_{\text{odd}}(\mathbb{R}^2)$ such that $u_1 > 0$ and $u_2 > 0$ on $\mathbb{R} \times (0, \infty)$. Next, define the maps

(2.9)
$$u_*(x_1, x_2) = \begin{cases} \min(u_1(x_1, x_2), u_2(x_1, x_2)) & \text{when } x_2 \ge 0, \\ \max(u_1(x_1, x_2), u_2(x_1, x_2)) & \text{when } x_2 \le 0, \end{cases}$$

(2.10)
$$u^*(x_1, x_2) = \begin{cases} \max(u_1(x_1, x_2), u_2(x_1, x_2)) & \text{when } x_2 \ge 0, \\ \min(u_1(x_1, x_2), u_2(x_1, x_2)) & \text{when } x_2 \le 0, \end{cases}$$

and the set $A := \{(x_1, x_2) \in \mathbb{R} \times (0, \infty) : u_1(x_1, x_2) < u_2(x_1, x_2)\}$. We can see that $E(u_1, A) = E(u_2, A)$ since otherwise we have either $E(u_*) < E(u_2)$ or $E(u^*) < E(u_1)$, which contradicts the minimality of u_1 and u_2 . As a consequence, $E(u_*) = E(u_2) = E(u_1) = E(u^*)$, and it follows that u_* and u^* are also minimizers and

solutions. Next, by unique continuation [26], we obtain that either $u_1 \equiv u_*$ or $u_1 \equiv u^*$, i.e. we have either $0 \leq u_1 \leq u_2$ or $u_1 \geq u_2 \geq 0$ on $\mathbb{R} \times [0, \infty)$. Finally, applying (2.7) to $E(u_1) = E(u_2)$, we conclude in view of the ordering of u_1 and u_2 that $u_1 \equiv u_2$. This completes the proof.

To study the limit of solutions as $\epsilon \to 0$, we need uniform bounds. Modifying slightly the arguments in [13, Section 2], we obtain:

Lemma 2.4. For every $\epsilon > 0$ let u_{ϵ} be a solution of (2.3) converging to 0 as $|x| \to \infty$. Then, u_{ϵ} are uniformly bounded.

Proof. We drop the index and write $u := u_{\epsilon}$. Since μ is bounded, the roots of the cubic equation $u^3 - \mu(x)u = 0$ belong to a bounded interval, for all values of x. If u takes positive values, then it attains its maximum $0 \le \max_{\mathbb{R}^2} u = u(x_0)$, at a point $x_0 \in \mathbb{R}^2$. In view of (2.3):

$$0 \ge \epsilon^2 \Delta u(x_0) = u^3(x_0) - \mu(x_0)u(x_0),$$

thus it follows that $u(x_0)$ is uniformly bounded above. In the same way, we prove the uniform lower bound for u.

Lemma 2.5. For $\epsilon \ll 1$ let u_{ϵ} be a solution of (2.3) converging to 0 as $|x| \to \infty$. Then, there exist a constant K > 0 such that

(2.11)
$$|u_{\epsilon}(x)| \le K(\sqrt{\max(\mu(x), 0)} + \epsilon^{1/3}), \quad \forall x \in \mathbb{R}^2.$$

As a consequence, if for every $\xi = \rho e^{i\theta}$ we consider the local coordinates $s = (s_1, s_2)$ in the basis $(e^{i\theta}, ie^{i\theta})$, then the rescaled functions $w_{\epsilon}(s)$ defined in (2.5) are uniformly bounded on the half-planes $[s_0, \infty) \times \mathbb{R}$, $\forall s_0 \in \mathbb{R}$.

Proof. As above we write $u := u_{\epsilon}$. Let us define the following constants

- M > 0 is the uniform bound of $|u_{\epsilon}|$ (cf. Lemma 2.4),
- $\lambda > 0$ is such that $3\mu_{\rm rad}(\rho h) \leq 2\lambda h, \forall h \in [0, \rho],$
- $\kappa > 0$ is such that $\kappa^4 \ge 6\lambda$.

Next, we construct the following comparison function

(2.12)
$$\chi(x) = \begin{cases} \lambda \left(\rho - |x| + \frac{\epsilon^{2/3}}{2} \right) & \text{for } |x| \le \rho, \\ \frac{\lambda}{2\epsilon^{2/3}} (|x| - \rho - \epsilon^{2/3})^2 & \text{for } \rho \le |x| \le \rho + \epsilon^{2/3}, \\ 0 & \text{for } |x| \ge \rho + \epsilon^{2/3}. \end{cases}$$

One can check that $\chi \in C^1(\mathbb{R}^2 \setminus \{0\}) \cap H^1(\mathbb{R}^2)$ satisfies $\Delta \chi \leq \frac{2\lambda}{\epsilon^{2/3}}$ in $H^1(\mathbb{R}^2)$. Finally, we define the function $\psi := \frac{|u|^2}{2} - \chi - \kappa^2 \epsilon^{2/3}$, and compute:

(2.13)

$$\epsilon^{2}\Delta\psi = \epsilon^{2}(|\nabla u|^{2} + u\Delta u - \Delta\chi)$$

$$\geq -\mu|u|^{2} + |u|^{4} - \epsilon^{2}\Delta\chi$$

$$\geq -\mu|u|^{2} + |u|^{4} - 2\epsilon^{4/3}\lambda.$$

Now, one can see that when $x \in \omega := \{x \in \mathbb{R}^2 : \psi(x) > 0\}$, we have $\frac{|u|^4}{3} - \mu |u|^2 \ge 0$, since

$$x \in \omega \cap \overline{D(0;\rho)} \Rightarrow \frac{|u|^4}{3} \ge \frac{2\lambda}{3} \left(\rho - |x| + \frac{\epsilon^{2/3}}{2}\right) |u|^2 \ge \mu |u|^2.$$

In the open set ω we also have: $\frac{|u|^4}{3} \ge \frac{\kappa^4}{3} \epsilon^{4/3} \ge 2\epsilon^{4/3}\lambda$, thus $\Delta \psi \ge 0$ in ω in the H^1 sense. To conclude, we apply Kato's inequality that gives: $\Delta \psi^+ \ge 0$ on \mathbb{R}^2 in the H^1 sense. Since ψ^+ is subharmonic with compact support, we obtain by the maximum principle that $\psi^+ \equiv 0$ or equivalently $\psi \le 0$ in \mathbb{R}^2 . The statement of the lemma follows by adjusting the constant K.

After this preparation we are ready to prove the main result of this section.

Proof Theorem 2.1. For every $\xi = \rho e^{i\theta}$ we consider the local coordinates $s = (s_1, s_2)$ in the basis $(e^{i\theta}, ie^{i\theta})$, and we rescale the solutions by setting $\tilde{u}(s) = \frac{u_{\epsilon}(\xi + s\epsilon^{2/3})}{\epsilon^{1/3}}$. Clearly $\Delta \tilde{u}(s) = \epsilon \Delta u(\xi + s\epsilon^{2/3})$, thus,

$$\Delta \tilde{u}(s) + \frac{\mu(\xi + s\epsilon^{2/3})}{\epsilon^{2/3}}\tilde{u}(s) - \tilde{u}^3(s) = 0, \qquad \forall s \in \mathbb{R}^2.$$

Writing $\mu(\xi + h) = \mu_1 h_1 + h \cdot A(h)$, with $\mu_1 := \mu'_{rad}(\rho) < 0$, $A \in C^{\infty}(\mathbb{R}^2, \mathbb{R}^2)$, and A(0) = 0, we obtain

(2.14)
$$\Delta \tilde{u}(s) + (\mu_1 s_1 + A(s\epsilon^{2/3}) \cdot s)\tilde{u}(s) - \tilde{u}^3(s) = 0, \qquad \forall s \in \mathbb{R}^2.$$

Next, we define the rescaled energy by

(2.15)
$$\tilde{E}(\tilde{u}) = \int_{\mathbb{R}^2} \left(\frac{1}{2} |\nabla \tilde{u}(s)|^2 - \frac{\mu(\xi + s\epsilon^{2/3})}{2\epsilon^{2/3}} \tilde{u}^2(s) + \frac{1}{4} \tilde{u}^4(s) \right) \mathrm{d}s$$

With this definition $\tilde{E}(\tilde{u}) = \frac{1}{\epsilon^{5/3}} E(u)$. From Lemma 2.5 and (2.14), it follows that $\Delta \tilde{u}$, and also $\nabla \tilde{u}$, are uniformly bounded on compact sets. Moreover, by differentiating (2.14) we also obtain the boundedness of the second derivatives of \tilde{u} . Thanks to these uniform bounds, we can apply the theorem of Ascoli via a diagonal argument to obtain the convergence of \tilde{u} in $C^2_{\text{loc}}(\mathbb{R}^2)$ (up to a subsequence) to a solution \tilde{z} of

(2.16)
$$\Delta \tilde{z}(s) + \mu_1 s_1 \tilde{z}(s) - \tilde{z}^3(s) = 0, \ \forall s \in \mathbb{R}^2,$$

which is associated to the functional

(2.17)
$$\tilde{E}_0(\phi, J) = \int_J \left(\frac{1}{2}|\nabla\phi(s)|^2 - \frac{\mu_1}{2}s_1\phi^2(s) + \frac{1}{4}\phi^4(s)\right) \mathrm{d}s.$$

Given $\tilde{\psi}(s)$ a test function supported in the compact set K, let $\psi(x) := \epsilon^{1/3} \tilde{\psi}(\frac{x-\xi}{\epsilon^{2/3}}) \Leftrightarrow \tilde{\psi}(s) = \frac{\psi(\xi+s\epsilon^{2/3})}{\epsilon^{1/3}}$. In the case where we take u to be the global minimizer v, since $E(v_{\epsilon} + \psi, \operatorname{supp} \psi) \geq E(v_{\epsilon}, \operatorname{supp} \psi)$, we have $\tilde{E}(\tilde{v}_{\epsilon} + \tilde{\psi}, K) \geq \tilde{E}(\tilde{v}_{\epsilon}, K)$, and at the limit $\tilde{E}_0(\tilde{z} + \tilde{\psi}, K) \geq \tilde{E}_0(\tilde{z}, K)$. Thus, \tilde{z} is a minimal solution of (2.16). In addition, the radial symmetry of v, implies that \tilde{z} depends only on the variable s_1 . Indeed, noticing that $\lim_{\epsilon \to 0} \frac{|\xi+\epsilon^{\frac{2}{3}}(s_1,s_2)|-\rho}{\epsilon^{\frac{2}{3}}} = s_1$, it follows that $\tilde{v}_{\epsilon}(s_1,s_2) = \tilde{v}_{\epsilon}(s_1 + o(1),0)$, and $\tilde{z}(s_1,s_2) = \tilde{z}(s_1,0)$. Similarly, in the case where we take u to be the odd minimizer and $\xi = (\pm \rho, 0)$, we can see that \tilde{z} is a minimal solution of (2.16) for perturbations such that $\tilde{\psi}(s_1,s_2) = -\tilde{\psi}(s_1,-s_2)$. Finally, setting $y(s) := \frac{1}{\sqrt{2}(-\mu_1)^{1/3}}\tilde{z}(\frac{s}{(-\mu_1)^{1/3}})$, (2.16) reduces to (2.6), that is, y solves (2.6). In the case where we take u to be the global minimizer v, we can see that either $y(s_1,s_2) = h(s_1)$ or $y(s_1,s_2) = -h(s_1)$, since $\pm h$ are the only minimal solutions of (1.4) (cf. [12, Theorem 1.3]). On the other hand, in the case where we take u to be the odd minimizer and $\xi = (\pm \rho, 0)$, it is clear that y is odd with respect to s_2 , and minimal for perturbations such that $\tilde{\psi}(s_1,s_2) = -\tilde{\psi}(s_1,-s_2)$.

3. Proof of Theorem 1.1

We will proceed in few steps. The proof of (i), (ii) and (iii) follows from Theorem 2.1, Lemma 2.5, and the fact that a minimal solution of 1.9 cannot be identically zero. To establish (v) we proceed as in Theorem 2.1. After rescaling appropriately y as $x_1 \to -\infty$, we compute uniform bounds of the rescaled functions. Then, by the theorem of Ascoli, we obtain at the limit a minimal solution of the Allen-Cahn equation (1.2). The proof of (vi) and (vii) is based on the moving plane method applied in a sector contained in the upper half-plane. The main difficulty is due to the unboundedness of the domain and to the availability of boundary conditions only on the x_1 axis where $y(x_1, 0) = 0$. We also utilize the asymptotic behaviour of y, as $x_1 \to \pm\infty$, provided respectively by (v) and Lemma 3.2. Our main tool is a version of the maximum principle in unbounded domains (cf. Lemma 3.1), that allows us to compute bounds for y_{x_1} and y_{x_2} when x_1 is large enough and $x_2 > 0$ (cf. Lemmas 3.3 and 3.4). Next, these bounds are extended to the whole half-plane $x_2 > 0$ by applying the sliding method (cf. Lemma 3.5).

Proof of (i), (ii) and (iii). By applying Theorem 2.1 in a neighborhood of the point $\xi = (\rho, 0)$ to the odd minimizer u, such that u > 0 on $\mathbb{R} \times (0, \infty)$, it is clear that we obtain a solution y of (2.6) which is odd with respect to the second variable s_2 , and such that $y \ge 0$, on $\mathbb{R} \times (0, \infty)$. For the sake of convenience in what follows we substitute the variables (s_1, s_2) by (x_1, x_2) . The properties (ii) and (iii) are also straightforward by Theorem 2.1 and Lemma 2.5. Thus, it remains to show that $y(x_1, x_2) > 0$, $\forall x \in \mathbb{R} \times (0, \infty)$. Assume by

contradiction that $y(x_1, x_2) = 0$, for some $x \in \mathbb{R} \times (0, \infty)$, then it follows from the maximum principle that $y \equiv 0$. To conclude we are going to show that a solution y of (1.9) which is minimal for odd perturbations, cannot be identically zero. Indeed, the minimality of y implies that the second variation of the energy $E_{P_{II}}$ is nonnegative:

(3.1)
$$\int_{\mathbb{R}^2} (|\nabla \phi(x)|^2 + (6y^2(x) + x_1)\phi^2(x)) dx \ge 0, \quad \forall \phi \in C_0^1(\mathbb{R}^2), \text{ such that } \phi(x_1, x_2) = -\phi(x_1, -x_2).$$

Clearly (3.1) does not hold when $y \equiv 0$, if we take $\phi(x) = \phi_0(x_1 + l, x_2)$, with $l \to \infty$, and $\phi_0 \in C_0^1(\mathbb{R}^2)$ fixed, such that $\phi_0(x_1, x_2) = -\phi_0(x_1, -x_2)$, and $\phi_0 \neq 0$.

Next we recall a useful version of the maximum principle in unbounded domains [9, Lemma 2.1].

Lemma 3.1. Let D be a domain (open connected set) in \mathbb{R}^n , possibly unbounded. Assume that \overline{D} is disjoint from the closure of an infinite open connected cone Σ . Suppose there is a function z in $C(\overline{D})$ that is bounded above and satisfies for some continuous function c(x)

$$\Delta z - c(x)z \ge 0 \text{ in } D \text{ with } c(x) \ge 0$$
$$z \le 0 \text{ on } \partial D.$$

Then $z \leq 0$ in D.

As a first application of Lemma 3.1 we prove the exponential convergence of y to 0, as $x_1 \to \infty$.

Lemma 3.2. $|y(x_1, x_2)| = O(e^{-\frac{2}{3}x_1^{3/2}})$, as $x_1 \to \infty$ (uniformly in x_2).

Proof. We define $\psi(x_1, x_2) := Me^{-\frac{2}{3}x_1^{3/2}}$, in the domain $D := \{(x_1, x_2) : x_1 > 1, x_2 > 0\}$, where $M \ge e^{\frac{2}{3}} \sup_{x_2 \ge 0} y(1, x_2)$ is a constant. It is easy to see that $\Delta \psi \le x_1 \psi$ in D, and $\Delta(y - \psi) \ge x_1(y - \psi)$ in D. Since $y - \psi \le 0$ on ∂D , it follows from Lemma 3.1 that $y \le \psi$ in D.

Proof of (v). We set $(t_1, t_2) := \left(-\frac{2}{3}(-x_1)^{\frac{3}{2}}, (-x_1)^{\frac{1}{2}}r\right)$, where $x_1 \leq -1$ and $r \in \mathbb{R}$. Equivalently we have $(x_1, r) = \left(-(-\frac{3}{2}t_1)^{\frac{2}{3}}, t_2(-\frac{3}{2}t_1)^{-\frac{1}{3}}\right)$. Next we define $\tilde{y}(t_1, t_2) := \frac{\sqrt{2}}{(-\frac{3}{2}t_1)^{\frac{1}{3}}} y(x_1, r + x_2)$, for every $x_2 \in \mathbb{R}$ fixed, or equivalently

(3.2)
$$y(x_1, r+x_2) = \frac{(-x_1)^{\frac{1}{2}}}{\sqrt{2}}\tilde{y}(t_1, t_2).$$

We are going to show that $\tilde{y}(t_1, t_2)$ is uniformly bounded up to the second derivatives, when t_2 belongs to a compact interval and $t_1 \to -\infty$. By differentiating (3.2) with respect to s_1 and r we obtain

(3.3a)
$$\sqrt{2}y_{x_2}(x_1, r + x_2) = (-x_1)\tilde{y}_{t_2}(t_1, t_2),$$

(3.3b)
$$\sqrt{2}y_{x_2x_2}(x_1, r + x_2) = (-x_1)^{\frac{3}{2}}\tilde{y}_{t_2t_2}(t_1, t_2),$$

(3.3c)
$$\sqrt{2}y_{x_1}(x_1, r+x_2) = -\frac{1}{2}(-x_1)^{-\frac{1}{2}}\tilde{y}(t_1, t_2) + (-x_1)\tilde{y}_{t_1}(t_1, t_2) - \frac{r}{2}\tilde{y}_{t_2}(t_1, t_2),$$

(3.3d)
$$\sqrt{2}y_{x_1x_2} = -\tilde{y}_{t_2} + (-x_1)^{\frac{3}{2}}\tilde{y}_{t_1t_2} - \frac{r}{2}(-x_1)^{\frac{1}{2}}\tilde{y}_{t_2,t_2},$$

(3.3e)
$$\sqrt{2}y_{x_1x_1} = -\frac{1}{4}(-x_1)^{-\frac{3}{2}}\tilde{y} - \frac{3}{2}\tilde{y}_{t_1} + \frac{r}{4}(-x_1)^{-1}\tilde{y}_{t_2} + (-x_1)^{\frac{3}{2}}\tilde{y}_{t_1t_1} - r(-x_1)^{\frac{1}{2}}\tilde{y}_{t_1t_2} + \frac{r^2}{4}(-x_1)^{-\frac{1}{2}}\tilde{y}_{t_2,t_2}.$$

Since by construction (cf. (2.11) in Lemma 2.5) y satisfies $|y(x_1, x_2)| = O(|-x_1|^{\frac{1}{2}})$ as $x_1 \to -\infty$ (i.e. \tilde{y} is bounded), we obtain by (1.9) and standard elliptic estimates [21, §3.4 p. 37] that

(3.4)
$$|\nabla y(x_1, x_2)| = O(|-x_1|^{\frac{3}{2}}) \text{ and } |D^2 y(x_1, x_2)| = O(|-x_1|^{\frac{5}{2}}), \text{ as } x_1 \to -\infty$$

From (3.4) and (3.3) it follows that

(3.5)
$$|\nabla \tilde{y}(t_1, t_2)| = O(|-x_1|^{\frac{1}{2}}) \text{ and } |D^2 \tilde{y}(t_1, t_2)| = O(|-x_1|), \text{ as } x_1 \to -\infty,$$

provided that $(t_1, t_2) \in \Sigma_{t_0, r_0} := \{(t_1, t_2) : t_1 \le t_0, |t_2| \le r_0(-\frac{3}{2}t_1)^{\frac{1}{3}}\}$, where $t_0 < 0$ and $r_0 > 0$ are arbitrary constants. In particular, we have $\sqrt{2}\Delta y(x_1, x_2) = (-x_1)^{\frac{3}{2}}\Delta \tilde{y}(t_1, t_2) + O(|-x_1|^{\frac{3}{2}})$, for $(t_1, t_2) \in \Sigma_{t_0, r_0}$. On the other hand it is clear by (1.9) that $\sqrt{2}\Delta y(x_1, x_2) = (-x_1)^{\frac{3}{2}}(\tilde{y}^3(t_1, t_2) - \tilde{y}(t_1, t_2))$, thus

(3.6)
$$|\Delta \tilde{y}(t_1, t_2)|$$
 and $|\nabla \tilde{y}(t_1, t_2)|$ are bounded, $\forall (t_1, t_2) \in \Sigma_{t_0, r_0}$.

Similarly, by differentiating once more equations (3.3) with respect to x_1 and r, one can show that

(3.7)
$$|D^2 \tilde{y}(t_1, t_2)| \text{ is bounded, } \forall (t_1, t_2) \in \Sigma_{t_0, r_0}$$

Next, we apply the theorem of Ascoli to the sequence $\tilde{y}(t_1+l,t_2)$ as $l \to -\infty$. Up to a subsequence $l_n \to -\infty$, we obtain via a diagonal argument, the convergence in $C^1_{\text{loc}}(\mathbb{R}^2)$ of $\tilde{y}_n(t_1,t_2) := \tilde{y}(t_1+l_n,t_2)$ to a bounded function $\tilde{z}(t_1,t_2)$ that we are going to determine. Our claim is that the limit \tilde{z} is a minimal solution of the Allen-Cahn equation (1.2), which is independent of the subsequence l_n . The proof of this property is based on the following energy considerations. Let (e_1, e_2) be the canonical basis of \mathbb{R}^2 . The energy functional

(3.8)
$$E_{\mathrm{P}_{\mathrm{II}}}(y,A) = \int_{A-x_2e_2} \left[\frac{1}{2} |\nabla y(x_1,r+x_2)|^2 + \frac{1}{2} x_1 y^2(x_1,r+x_2) + \frac{1}{2} y^4(x_1,r+x_2) \right] \mathrm{d}x_1 \mathrm{d}r,$$

associated to (1.9), becomes after changing variables as in (3.2)

(3.9)
$$E_{\mathrm{P}_{\mathrm{II}}}(y,A) = \tilde{E}_{\mathrm{P}_{\mathrm{II}}}(\tilde{y},\tilde{A}) = \tilde{F}(\tilde{y},\tilde{A}) + \tilde{R}(\tilde{y},\tilde{A}),$$

where

(3.10)
$$\tilde{A} := \{ (t_1(x_1), t_2(x_1, r)) : (x_1, r) \in A - x_2 e_2 \},\$$

(3.11)
$$\tilde{F}(\tilde{y},\tilde{A}) := \int_{\tilde{A}} \frac{1}{2} \left(-\frac{3}{2} t_1 \right)^{\frac{2}{3}} \left[\frac{1}{2} |\nabla \tilde{y}(t_1,t_2)|^2 - \frac{\tilde{y}^2(t_1,t_2)}{2} + \frac{\tilde{y}^4(t_1,t_2)}{4} \right] \mathrm{d}t_1 \mathrm{d}t_2,$$

and

(3.12)
$$\tilde{R}(\tilde{y},\tilde{A}) := \int_{\tilde{A}} \left[\frac{(\tilde{y} + t_2 \tilde{y}_{t_2})^2}{16(-\frac{3}{2}t_1)^{\frac{4}{3}}} - \frac{(\tilde{y} + t_2 \tilde{y}_{t_2})\tilde{y}_{t_1}}{4(-\frac{3}{2}t_1)^{\frac{1}{3}}} \right] \mathrm{d}t_1 \mathrm{d}t_2$$

Let $\tilde{\phi}(t_1,t_2) \in C_0^{\infty}(\mathbb{R}^2)$ be a test function such that $\tilde{B} := \operatorname{supp} \tilde{\phi} \subset \{(t_1,t_2) : c-d \leq t_1 \leq c\}$, for some constants $c \in \mathbb{R}$ and d > 0. Given $l \in \mathbb{R}$, we consider the translated functions $\tilde{\phi}^{-l}(t_1,t_2) := \tilde{\phi}(t_1-l,t_2)$, and $\tilde{y}^l(t_1,t_2) = \tilde{y}(t_1+l,t_2)$. Note that $\tilde{B}^l := \operatorname{supp} \tilde{\phi}^{-l} = \tilde{B} + le_1$, and $\operatorname{supp} \tilde{\phi}^{-l} \subset \{(t_1,t_2) : t_1 < -1\}$ when l < 1-c. Thus, for l < 1-c, we can define $\phi^{-l} \in C_0^{\infty}(\mathbb{R}^2)$ by $\phi^{-l}(x_1,r+x_2) = \frac{(-x_1)^{\frac{1}{2}}}{\sqrt{2}} \tilde{\phi}^{-l}(t_1,t_2)$ as in(3.2). Let $B^l := \{(x_1(t_1), r(t_1,t_2) + x_2) : (t_1,t_2) \in \tilde{B}^l\}$.

We first examine the case where $x_2 = 0$, and assume that $\tilde{\phi}(t_1, t_2) = -\tilde{\phi}(t_1, -t_2)$. In view of the minimality of y and (3.9), we have

(3.13)
$$\tilde{E}_{\mathrm{P}_{\mathrm{II}}}(\tilde{y}+\tilde{\phi}^{-l},\tilde{B}^{l})=E_{\mathrm{P}_{\mathrm{II}}}(y+\phi^{-l},B^{l})\geq E_{\mathrm{P}_{\mathrm{II}}}(y,B^{l})=\tilde{E}_{\mathrm{P}_{\mathrm{II}}}(\tilde{y},\tilde{B}^{l}).$$

On the one hand, it is clear that the boundedness of \tilde{y} and (3.6) imply that $\lim_{l\to\infty} \tilde{R}(\tilde{y} + \tilde{\phi}^{-l}, \tilde{B}^l) = 0$ and $\lim_{l\to\infty} \tilde{R}(\tilde{y}, \tilde{B}^l) = 0$. Next, setting $t_0 := c + l$, we have

(3.14)
$$\left(-\frac{3}{2}t_1\right)^{\frac{2}{3}} \le \left(-\frac{3}{2}t_0\right)^{\frac{2}{3}} + d\left(-\frac{3}{2}t_0\right)^{-\frac{1}{3}}, \forall t_1 \in [t_0 - d, t_0].$$

Thus, we obtain

(3.15)
$$\tilde{F}(\tilde{y}, \tilde{B}^l) = \frac{1}{2} \left(-\frac{3}{2} t_0 \right)^{\frac{2}{3}} \tilde{G}(\tilde{y}, \tilde{B}^l) + O(|t_0|^{-\frac{1}{3}}) = \frac{1}{2} \left(-\frac{3}{2} t_0 \right)^{\frac{2}{3}} \tilde{G}(\tilde{y}^l, \tilde{B}) + O(|t_0|^{-\frac{1}{3}}),$$

and

$$(3.16) \quad \tilde{F}(\tilde{y} + \tilde{\phi}^{-l}, \tilde{B}^{l}) = \frac{1}{2} \left(-\frac{3}{2} t_{0} \right)^{\frac{2}{3}} \tilde{G}(\tilde{y} + \tilde{\phi}^{-l}, \tilde{B}^{l}) + O(|t_{0}|^{-\frac{1}{3}}) = \frac{1}{2} \left(-\frac{3}{2} t_{0} \right)^{\frac{2}{3}} \tilde{G}(\tilde{y}^{l} + \tilde{\phi}, \tilde{B}) + O(|t_{0}|^{-\frac{1}{3}}),$$

$$10$$

where we have set $\tilde{G}(\tilde{z},\tilde{B}) := \int_{\tilde{B}} (\frac{1}{2} |\nabla \tilde{z}|^2 - \frac{\tilde{z}^2}{2} + \frac{\tilde{z}^4}{4}) dt$. Finally, since $\tilde{y}^{l_n}(t_1,t_2) \to \tilde{z}(t_1,t_2)$ in $C^1_{\text{loc}}(\mathbb{R}^2)$, as $n \to \infty$, we conclude that

$$(3.17) \quad \tilde{G}(\tilde{z}+\tilde{\phi},\tilde{B}) = \lim_{n \to \infty} \frac{2}{(-\frac{3}{2}(c+l_n))^{\frac{2}{3}}} \tilde{E}_{\mathrm{PII}}(\tilde{y}^{l_n}+\tilde{\phi},\tilde{B}) \ge \lim_{n \to \infty} \frac{2}{(-\frac{3}{2}(c+l_n))^{\frac{2}{3}}} \tilde{E}_{\mathrm{PII}}(\tilde{y}^{l_n}+\tilde{\phi},\tilde{B}) = \tilde{G}(\tilde{z},\tilde{B}),$$

or equivalently $E_{AC}(\tilde{z} + \tilde{\phi}, \tilde{B}) \geq E_{AC}(\tilde{z}, \tilde{B})$. This means that \tilde{z} is a minimal solution of the Allen-Cahn equation (1.2) for odd perturbations $\tilde{\phi}$. In particular $\tilde{z} \neq 0$, and as a consequence of the maximum principle, $\tilde{z}(t_1, 0) = 0, \forall t_1 \in \mathbb{R}, \text{ and } \tilde{z}(t_1, t_2) \geq 0, \forall (t_1, t_2) \in \mathbb{R} \times (0, \infty)$, imply that $\tilde{z}(t_1, t_2) > 0, \forall (t_1, t_2) \in \mathbb{R} \times (0, \infty)$. Thus, from [10, Theorem 1.5], it follows that \tilde{z} is a function of only t_2 , which is the heteroclinic connection $\tilde{z}(t_1, t_2) = \eta(t_2) = \tanh(t_2/\sqrt{2})$. Furthermore, since the limit \tilde{z} is uniquely determined, the convergence $\tilde{y}^l(t_1, t_2) \to \tilde{z}(t_1, t_2)$ holds as $l \to -\infty$.

It remains to examine the case where $x_2 \neq 0$. Without loss of generality we assume that $x_2 > 0$. Now (3.13) holds for arbitrary test functions $\tilde{\phi}(t_1, t_2) \in C_0^{\infty}(\mathbb{R}^2)$, since $B^l \subset \{(x_1, x_2) : x_2 > 0\}$ as $l \to -\infty$. Repeating the previous arguments we find that \tilde{z} is a nonnegative minimal solution of (1.2). Applying [6, Corollary 5.2], we deduce that $\tilde{z} \equiv 1$. This completes the proof of (v).

Proof of (vi) and (vii). The proofs of (vi) and (vii) which are based on the moving plane method, follow from the next lemmas.

Lemma 3.3. We have $y_{x_1}(x_1, x_2) < 0$, $\forall x_1 \ge 0$, $\forall x_2 > 0$. In addition, for every d > 0, there holds $\sup_{x_2 > d} y_{x_1}(1, x_2) < 0$, and $\inf_{x_2 \ge d} y(1, x_2) > 0$.

Proof. Given $\lambda \geq 0$, we define the function $\psi_{\lambda}(x_1, x_2) := y(x_1, x_2) - y(-x_1 + 2\lambda, x_2)$ for $(x_1, x_2) \in D_{\lambda} := \{(x_1, x_2) : x_1 > \lambda, x_2 > 0\}$. One can check that $\psi_{\lambda} = 0$ on ∂D_{λ} , and

$$\Delta \psi_{\lambda} - c(x_1, x_2)\psi_{\lambda} = 2(x_1 - \lambda)y(-x_1 + 2\lambda, x_2) \ge 0,$$

with $c(x_1, x_2) = x_1 + 2(y^2(x_1, x_2) + y(x_1, x_2)y(-x_1 + 2\lambda, x_2) + y^2(-x_1 + 2\lambda, x_2)) \geq 0$. Furthermore, ψ_{λ} is bounded above by Theorem 1.1 (ii), and not identically zero by Theorem 1.1 (v). As a consequence of Lemma 3.1, it follows that $\psi_{\lambda}(x_1, x_2) < 0$, $\forall x_1 > \lambda$, $\forall x_2 > 0$, and thus by Hopf's Lemma $\frac{\partial \psi_{\lambda}}{\partial x_1}(\lambda, x_2) = 2y_{x_1}(\lambda, x_2) < 0$, $\forall x_2 > 0$. To establish that $\sup_{x_2 \geq d} y_{x_1}(1, x_2) < 0$, we proceed by contradiction and assume the existence of a sequence $\{l_n\}$ such that $\lim_{n\to\infty} l_n = \infty$ and $\lim_{n\to\infty} y_{x_1}(1, l_n) = 0$. Next, we set $\tilde{y}_n(x_1, x_2) = y(x_1, x_2 + l_n)$. In view of the bounds provided in Theorem 1.1 (ii), we obtain by the theorem of Ascoli that (up to subsequence) \tilde{y}_n converges in C^1_{loc} to a nonnegative minimal solution \tilde{y} of (1.9). Since $\tilde{y}_{x_1}(1, 0) = \lim_{n\to\infty} y_{x_1}(1, l_n) = 0$, and $\tilde{y}_{x_1}(x_1, x_2) \leq 0$, $\forall x_1 \geq 0$, $\forall x_2 \in \mathbb{R}$, the maximum principle applied to

(3.18)
$$\Delta \tilde{y}_{x_1} = \tilde{y} + (x_1 + 6\tilde{y}^2)\tilde{y}_{x_1} \ge (x_1 + 6\tilde{y}^2)\tilde{y}_{x_1},$$

implies that $\tilde{y}_{x_1}(x_1, x_2) = 0$, $\forall x_1 \geq 0$, $\forall x_2 \in \mathbb{R}$. Then, since $\lim_{x_1 \to \infty} \tilde{y}(x_1, x_2) = 0$, $\forall x_2 \in \mathbb{R}$, it follows that $\tilde{y} \equiv 0$ in the half-plane $x_1 \geq 0$. Finally, we deduce by unique continuation that $\tilde{y} \equiv 0$ in \mathbb{R}^2 , which is a contradiction since \tilde{y} is minimal. Thus we have established that $\sup_{x_2 \geq d} y_{x_1}(1, x_2) < 0$. The proof that $\inf_{x_2 \geq d} y(1, x_2) > 0$ is similar. \Box

Lemma 3.4. For every vector $\mathbf{n} = e^{i(\theta + \frac{\pi}{2})} \in \mathbb{C} \sim \mathbb{R}^2$, with $\theta \in (0, \frac{\pi}{2})$, there exists $s_{\mathbf{n}} > 0$ such that $\nabla y(x_1, x_2) \cdot \mathbf{n} > 0$, $\forall x_1 > s_{\mathbf{n}}$, $\forall x_2 > 0$.

Proof. Our first claim is that there is a constant $k_1 > 0$, such that $k_1y_{x_1}(x_1, x_2) \leq -\sqrt{x_1}y(x_1, x_2)$, $\forall x_1 \geq 1$, $\forall x_2 \geq 0$. Indeed, let $\psi(x_1, x_2) = k_1y_{x_1}(x_1, x_2) + \sqrt{x_1}y(x_1, x_2)$ for $(x_1, x_2) \in D := \{x_1 > 1, x_2 > 0\}$, where the constant k_1 will be adjusted later. It is clear that $\psi(x_1, 0) = 0$, $\forall x_1 \geq 1$. We also note that $y_{x_1x_2}(1, 0) < 0$, since the function y_{x_1} vanishes at (1, 0), is negative in $\{x_1 > 0, x_2 > 0\}$, and satisfies (3.18). This and $\sup_{x_2 \geq d} y_{x_1}(1, x_2) < 0$, $\forall d > 0$, imply that when k_1 is large enough, we have $\psi(1, x_2) \leq 0$, $\forall x_2 \geq 0$. Next, we compute

$$\Delta \psi = \left(x_1 + 6y^2 + \frac{1}{k_1\sqrt{x_1}}\right)k_1y_{x_1} + \left(x_1 + 2y^2 + \frac{k_1}{\sqrt{x_1}} - \frac{1}{4x_1^2}\right)\sqrt{x_1}y$$
$$= \left(x_1 + 2y^2 + \frac{k_1}{\sqrt{x_1}} - \frac{1}{4x_1^2}\right)\psi + \left(4y^2 + \frac{1}{k_1\sqrt{x_1}} - \frac{k_1}{\sqrt{x_1}} + \frac{1}{4x_1^2}\right)k_1y_{x_1}.$$

By choosing k_1 large enough we can ensure that $\left(x_1 + 2y^2 + \frac{k_1}{\sqrt{x_1}} - \frac{1}{4x_1^2}\right) \ge 0$ and $\left(4y^2 + \frac{1}{k_1\sqrt{x_1}} - \frac{k_1}{\sqrt{x_1}} + \frac{1}{4x_1^2}\right) \le 0$, when $x_1 \ge 1$ and $x_2 \ge 0$. Thus, by applying Lemma 3.1, our claim follows.

Similarly, we are going to establish that there is a constant $k_2 > 0$, such that $y_{x_2}(x_1, x_2) \ge -k_2y(x_1, x_2)$, $\forall x_1 \ge 1, \forall x_2 \ge 0$. To do this we let $\psi(x_1, x_2) = -y_{x_2}(x_1, x_2) - k_2y(x_1, x_2)$ for $(x_1, x_2) \in D$, where the constant k_2 will again be adjusted later. We first note that $y_{x_2}(x_1, 0) > 0, \forall x_1 \in \mathbb{R}$, since the function y vanishes at $(x_1, 0)$, is positive in $\{x_2 > 0\}$, and satisfies (1.9). This and $\inf_{x_2 \ge d} y(1, x_2) > 0, \forall d > 0$, imply that when k_2 is large enough, we have $\psi(1, x_2) \le 0, \forall x_2 \ge 0$. On the other hand, it is clear that $\psi(x_1, 0) < 0, \forall x_1 \ge 1$. Next, we compute

$$\Delta \psi = (x_1 + 6y^2)(-y_{x_2}) + (x_1 + 2y^2)(-k_2y) \ge (x_1 + 6y^2)\psi.$$

Thus, by applying Lemma 3.1, it follows that $\psi \leq 0$ in D.

Finally, setting $\frac{\nabla y}{|\nabla y|} = e^{i\phi}$ when $(x_1, x_2) \in D$ (with $\phi \in (\frac{\pi}{2}, \frac{3\pi}{2})$), we find that

$$\tan \phi = \frac{y_{x_2}}{y_{x_1}} \le \frac{k_1 k_2}{\sqrt{x_1}} \Longrightarrow \phi \le \pi + \arctan\left(\frac{k_1 k_2}{\sqrt{x_1}}\right).$$

As a consequence, we have $\nabla y(x_1, x_2) \cdot \boldsymbol{n} > 0$ if $\theta \in \left(\arctan\left(\frac{k_1k_2}{\sqrt{x_1}}\right), \frac{\pi}{2}\right)$, that is, if $x_1 > s_n := \left(\frac{k_1k_2}{\tan\theta}\right)^2$. \Box

Lemma 3.5. Let $\theta \in (0, \frac{\pi}{2})$ be fixed, and consider for every $\lambda \in \mathbb{R}$ the reflection σ_{λ} with respect to the line $\Gamma_{\lambda} := \{(x_1, x_2) : x_2 = \tan \theta(x_1 - \lambda)\}$, and the domain $D_{\lambda} := \{(x_1, x_2) : 0 < x_2 < \tan \theta(x_1 - \lambda)\}$. Then, the function $\psi_{\lambda}(x_1, x_2) := y(x_1, x_2) - y(\sigma_{\lambda}(x_1, x_2))$ is negative in D_{λ} , for every $\lambda \in \mathbb{R}$.

Proof. We set $\mathbf{n} = e^{i(\theta + \frac{\pi}{2})}$ as in Lemma 3.4, and denote by (p', q') the image by σ_{λ} of a point $(p, q) \in D_{\lambda}$, and by D'_{λ} the set $\sigma_{\lambda}(D_{\lambda})$. It is obvious that $\psi_{\lambda}(x_1, 0) < 0$, $\forall x_1 > \lambda$, and that $\psi_{\lambda}(x_1, x_2) = 0$, $\forall (x_1, x_2) \in \Gamma_{\lambda}$. Moreover, ψ_{λ} satisfies

$$\Delta \psi_{\lambda}(p,q) - c(p,q)\psi_{\lambda} = (p-p')y(p',q') \ge 0, \quad \forall (p,q) \in D_{\lambda},$$

with $c(p,q) = p + 2(y^2(p,q) + y(p,q)y(p',q') + y^2(p',q'))$. For each $\lambda \in \mathbb{R}$ we consider the statement

(3.19)
$$\psi_{\lambda}(p,q) < 0, \quad \forall (p,q) \in D_{\lambda}.$$

We shall first establish Lemma 3.5 in the case where $\theta \in (0, \frac{\pi}{4})$. According to Lemma 3.4, (3.19) is valid for each $\lambda \geq s_n$. Set $\lambda_0 = \inf\{\lambda \in \mathbb{R} : \psi_{\mu} < 0 \text{ holds in } D_{\mu}, \text{ for each } \mu \geq \lambda\}$. We will prove $\lambda_0 = -\infty$. Assume instead $\lambda_0 \in \mathbb{R}$. Then, there exist a sequence $\lambda_k < \lambda_0$ such that $\lim_{k\to\infty} \lambda_k = \lambda_0$, and a sequence $(p_k, q_k) \in D_{\lambda_k}$, such that $y(p_k, q_k) \geq y(p'_k, q'_k)$. According to Lemma 3.4, we have $p'_k \leq s_n$, thus the sequence (p_k, q_k) is bounded. Up to subsequence we may assume that $\lim_{k\to\infty}(p_k, q_k) = (p_0, q_0) \in \overline{D_{\lambda_0}}$, with $p'_0 \leq s_n$. By definition of λ_0 , we have $\psi_{\lambda_0} \leq 0$ in D_{λ_0} , and $\psi_{\lambda_0}(p_0, q_0) = 0$ i.e. $y(p_0, q_0) = y(p'_0, q'_0)$. Now we distinguish the following cases. If $(p_0, q_0) \in D_{\lambda_0}$, the maximum principle implies that $\psi_{\lambda_0} \equiv 0$ in D_{λ_0} . Clearly, this situation is excluded, since y is positive in the half-plane $\{x_2 > 0\}$. On the other hand, the maximum principle also implies that $\frac{\partial \psi_{\lambda_0}}{\partial n}(p,q) = 2\frac{\partial y}{\partial n}(p,q) > 0$, provided that $(p,q) \in \Gamma_{\lambda_0}$ and q > 0. Furthermore, the previous inequality still holds at the vertex $(p,q) = (\lambda_0,0)$, since $y_{x_2}(x_1,0) > 0$ and $y_{x_1}(x_1,0) = 0, \forall x_1 \in \mathbb{R}$ (cf. the proof of Lemma 3.4). As a consequence, in a neighborhood of the line segment $\{(x_1, x_2) : x_2 = \tan \theta(x_1 - \lambda), 0 \leq x_1 \leq s_n\}$, we have that $\frac{\partial y}{\partial n} > 0$, and it follows that (p_0, q_0) cannot belong to Γ_{λ_0} . Finally, since the case where $p_0 > \lambda_0$ and $q_0 = 0$ is ruled out (because y is positive in the half-plane $\{x_2 > 0\}$), we have reached a contradiction.

Next, we establish Lemma 3.5 in the case where $\theta \in [\frac{\pi}{4}, \frac{\pi}{2})$, which is a little bit more involved. When $\theta = \frac{\pi}{4}$, it is clear that (3.19) is valid for each $\lambda \geq s_n$. Otherwise, when $\theta \in (\frac{\pi}{4}, \frac{\pi}{2})$, let $A'_{\lambda} := \{(p', q') \in D'_{\lambda} : p' \leq s_n\}$, and let $A_{\lambda} = \sigma_{\lambda}(A'_{\lambda})$. Our first claim is that $m := \inf_{A'_{s_n+1}} y > 0$. Indeed, proceeding as in the proof of Theorem 1.1 (v), one can see that

$$\lim_{(x_1,x_2)\in A'_{x_{n+1}}, x_1\to -\infty} \frac{\sqrt{2}}{\sqrt{-x_1}} y(x_1,x_2) = 1.$$

In addition, proceeding as in the proof of Lemma 3.4, we obtain that $\inf\{y(x_1, x_2) : (x_1, x_2) \in A'_{s_n+1}, s_n - l \le x_1 \le s_n\} > 0$, for every constant l > 0. Thus, m > 0. On the other hand, we have $\lim_{\lambda \to \infty} \sup\{y(x_1, x_2) : (x_1, x_2) : (x_1, x_2) \in A'_{s_n+1}, s_n - l \le x_1 \le s_n\}$



FIGURE 1. The sets A_{λ} , A'_{λ} , $B_{\lambda,\nu}$, $B'_{\lambda,\nu}$, and the lines Γ_{λ} , Δ_{ν} , in the case where $\lambda > s_n$ and $\lambda < s_n$.

 $(x_1, x_2) \in A_{\lambda} = 0$, since $\lim_{\lambda \to \infty} \inf\{x_1 : (x_1, x_2) \in A_{\lambda}\} = 0$ (cf. Lemma 3.2). As a consequence when $\lambda \geq s_n + 1$ is large enough, we have $y(p', q') \geq m > y(p, q)$, $\forall(p, q) \in A_{\lambda}$, and also y(p', q') > y(p, q), $\forall(p, q) \in D_{\lambda} \setminus A_{\lambda}$, by definition of s_n . This establishes that (3.19) holds for λ large enough. Then, defining λ_0 as previously, we assume by contradiction that $\lambda_0 \in \mathbb{R}$, and deduce in a similar way the existence of the sequences λ_k and $(p_k, q_k) \in D_{\lambda_k}$. We need to show that (p_k, q_k) is bounded. For $\nu > \lambda$, let $M_{\nu} := (\nu, \tan \theta(\nu - \lambda)) \in \Gamma_{\lambda}$, and let $\Delta_{\nu} := \{(x_1, x_2) : x_2 = \tan(\theta + \frac{\pi}{2})(x_1 - \nu) + \tan \theta(\nu - \lambda)\}$ be the line parallel to n and passing through M_{ν} . Let also $B'_{\lambda,\nu} := \{(p',q') \in A'_{\lambda} : q' \geq \tan(\theta + \frac{\pi}{2})(p' - \nu) + \tan \theta(\nu - \lambda)\}$ be the subset of A'_{λ} which is above Δ_{ν} , and $B_{\lambda,\nu} := \sigma_{\lambda}(B'_{\lambda,\nu})$. Proceeding as previously, we can see that $\forall \nu > \lambda_0 + 2$, $\forall \lambda > \lambda_0 - 1$, we have $\inf_{B'_{\lambda,\nu}} y > m$ for some constant m > 0, while $\lim_{\nu \to \infty} \sup\{y(x_1, x_2) : (x_1, x_2) \in B_{\lambda,\nu}\} = 0$. As a consequence, for ν large enough and $\lambda > \lambda_0 - 1$, we have $y(p', q') \geq m > y(p, q)$, $\forall(p, q) \in B_{\lambda,\nu}$, and thus $(p_k, q_k) \notin B_{\lambda,\mu}$. Furthermore, since $p'_k \leq s_n$ by Lemma 3.4, we have established the boundedness of (p_k, q_k) .

Lemma 3.5 implies that $\forall \theta \in (0, \frac{\pi}{2}), \forall \lambda \in \mathbb{R}$, and $(p, q) \in \Gamma_{\lambda}$ with q > 0, we have $\frac{\partial \psi_{\lambda}}{\partial n}(p, q) = 2\frac{\partial y}{\partial n}(p, q) > 0$, where $\mathbf{n} = e^{i(\theta + \frac{\pi}{2})}$. It follows that $y_{x_1}(x_1, x_2) \leq 0$, and $y_{x_2}(x_1, x_2) \geq 0$, $\forall x_1 \in \mathbb{R}, \forall x_2 \geq 0$. Moreover, in the half-plane $x_2 \geq 0, y_{x_1}$ and y_{x_2} satisfy respectively $\Delta y_{x_1} \geq (x_1 + 6y^2)y_{x_1}$, and $\Delta y_{x_2} = (x_1 + 6y^2)y_{x_2}$, thus y_{x_1} (resp. y_{x_2}) cannot vanish in the open half-plane $x_2 > 0$, since otherwise we would obtain by the maximum principle $y_{x_1} \equiv 0$ (resp. $y_{x_2} \equiv 0$). These situations are excluded by the fact that y > 0 in the open half-plane $x_2 > 0$, and $y_{x_2}(x_1, 0) > 0, \forall x_1 \in \mathbb{R}$. Therefore we have proved that $y_{x_1}(x_1, x_2) < 0, \forall x_1 \in \mathbb{R}, \forall x_2 > 0$, and $y_{x_2}(x_1, x_2) > 0, \forall x_1, x_2 \in \mathbb{R}$. Finally, setting $\tilde{y}_l(x_1, x_2) = y(x_1, x_2 + l)$, we obtain by the Theorem of Ascoli, that up to a subsequence $l_k \to \infty$, \tilde{y}_{l_k} converges in C_{loc}^2 to a nonnegative minimal solution \tilde{y}_{∞} of (1.9). Furthermore, the monotonicity of y along the x_2 direction implies that \tilde{y}_{∞} is independent of x_2 . Thus, since h is the only nonnegative minimal solution of (1.9) (cf. [12, Theorem 1.3]), we deduce that $\tilde{y}_{\infty}(x_1, x_2) = h(x_1)$, and that $\lim_{l\to\infty} y(x_1, x_2 + l) = h(x_1)$ is independent of the subsequence l_k . We also note that $|y(x_1, x_2)| < h(x_1), \forall (x_1, x_2) \in \mathbb{R}^2$, from which Theorem 1.1 (iv) follows. This completes the proof of Theorem 1.1.

References

- M. J. Ablowitz and P. A. Clarkson, Solitons, nonlinear evolution equations and inverse scattering, London Mathematical Society Lecture Note Series, vol. 149, Cambridge University Press, Cambridge, 1991. MR 1149378 (93g:35108)
- [2] Amandine Aftalion and Xavier Blanc, Existence of vortex-free solutions in the Painlevé boundary layer of a Bose-Einstein condensate, J. Math. Pures Appl. (9) 83 (2004), no. 6, 765–801. MR 2062641
- [3] Amandine Aftalion, Robert L. Jerrard, and Jimena Royo-Letelier, Non-existence of vortices in the small density region of a condensate, J. Funct. Anal. 260 (2011), no. 8, 2387–2406. MR 2772375
- [4] Alama, S., Bronsard, L., Gui, C.: Stationary layered solutions in R² for an Allen–Cahn system with multiple well potential. Calc. Var. 5 No. 4, 359–390 (1997)
- [5] Nicholas D. Alikakos, Peter W. Bates, Cahn John W., Paul C. Fife, and Fusco G.and Tanoglu G. B., Analysis of a corner layer problem in anisotropic interfaces, Discrete Contin. Dyn. Syst. Ser. B 6 (2006), no. 2, 237–255.
- [6] Alikakos, N. D., Fusco, G., Smyrnelis, P.: Elliptic systems of phase transition type. Progress in Nonlinear Differential Equations and Their Applications 91, Springer-Birkhäuser (2018).
- [7] Ambrosio, L., Cabré, X.: Entire solutions of semilinear elliptic equations in R³ and a conjecture of De Giorgi. J. Am. Math. Soc. 13, 725–739 (2000)
- [8] R. Barboza, U. Bortolozzo, J. D. Davila, M. Kowalczyk, S. Residori, and E. Vidal Henriquez, Light-matter interaction induces a shadow vortex, Phys. Rev. E 93 (2016), no. 5, 050201.
- Berestycki, H., Caffarelli, L., Nirenberg, L.: Monotonicity for elliptic equations in an unbounded Lipschitz domain. Comm. Pure Appl. Math. 50, 1089-1112 (1997)
- [10] Berestycki, H., Caffarelli, L., Nirenberg, L.: Further qualitative properties for elliptic equations in unbounded domains. Ann. Scuola Norm. Sup. Pisa Cl. Sci. No. 4 25, 69-94 (1997)
- [11] T. Claeys, A. B. J. Kuijlaars, and M. Vanlessen, Multi-critical unitary random matrix ensembles and the general Painleve II equation, Ann. of Math. (2) 167 (2008), 601–641.
- [12] Marcel G. Clerc, Juan Diego Dávila, Michał Kowalczyk, Panayotis Smyrnelis, and Estefania Vidal-Henriquez, Theory of light-matter interaction in nematic liquid crystals and the second Painlevé equation, Calculus of Variations and Partial Differential Equations 56 (2017), no. 4, 93.
- [13] Marcel G. Clerc, Michał Kowalczyk, and Panayotis Smyrnelis, Symmetry breaking and restoration in the ginzburg-landau model of nematic liquid crystals, Journal of Nonlinear Science 28 (2018), no. 3, 1079–1107.
- [14] Marcel Clerc, Michał Kowalczyk, and Panayotis Smyrnelis, Gradient theory of domain walls in thin, nematic liquid crystals films, Preprint 2018.
- [15] P. Deift, Universality for mathematical and physical systems, ArXiv Mathematical Physics e-prints (2006).
- [16] Dang, H., Fife, P. C., Peletier, L. A.: Saddle solutions of the bistable diffusion equation. Z. Angew. Math. Phys., 43 (6), 984–998 (1992)
- [17] De Giorgi, E.: Convergence problems for functionals and operators. In: Proceedings of the International Meeting on Recent Methods in Nonlinear Analysis, Rome (1978), 131–188. Pitagora, Bologna (1979)
- [18] Farina, A., Valdinoci, E.: The state of art for a conjecture of De Giorgi and related questions. Reaction-diffusion systems and viscosity solutions. World Scientific, Singapore (2008)
- [19] Ghoussoub, N., Gui, C.: On a conjecture of De Giorgi and some related problems. Math. Ann. 311, 481-491 (1998)
- [20] Hermann Flaschka and Alan C. Newell, Monodromy and spectrum-preserving deformations I, Communications in Mathematical Physics 76 (1980), no. 1, 65–116.
- [21] D. Gilbarg and N. S. Trudinger, Elliptic Partial Differential Equations of Second Order, (1983).
- [22] S. P. Hastings and J. B. McLeod, A boundary value problem associated with the second Painlevé transcendent and the Korteweg-de Vries equation, Arch. Rational Mech. Anal. 73 (1980), no. 1, 31–51. MR 555581
- [23] Bernard Helffer and Fred B. Weissler, On a family of solutions of the second Painlevé equation related to superconductivity, European Journal of Applied Mathematics 9 (1998), no. 3, 223–243.
- [24] Georgia Karali and Christos Sourdis, The ground state of a Gross-Pitaevskii energy with general potential in the Thomas-Fermi limit, Arch. Ration. Mech. Anal. 217 (2015), no. 2, 439–523. MR 3355003
- [25] Nicolai A. Kudryashov, The second painlevé equation as a model for the electric field in a semiconductor, Physics Letters A 233 (1997), no. 4, 397 – 400.
- [26] M. Sanada, Strong unique continuation property for some second order elliptic systems, Proc. Japan Acad. 83, Ser. A (2007).
- [27] Ovidiu Savin, Regularity of flat level sets in phase transitions, Ann. of Math. (2) 169 (2009), no. 1, 41–78. MR 2480601 (2009m:58025)
- [28] Savin, O.: Minimal Surfaces and Minimizers of the Ginzburg Landau energy. Cont. Math. Mech. Analysis AMS 526, 43–58 (2010)
- [29] Schatzman, M.: Asymmetric heteroclinic double layers. Control Optim. Calc. Var. 8 (A tribute to J. L. Lions), 965–1005 (electronic) (2002)
- [30] Sourdis, C. and Paul C. Fife, Existence of heteroclinic orbits for a corner layer problem in anisotropic interfaces, Adv. Differential Equations 12 (2007), no. 623–668.
- [31] Sourdis, C.: Painlevé-II profile of the shadow kink in the theory of light-matter interaction in nematic liquid crystals. Preprint.

[32] Troy, W.: The role of Painlevé II in predicting new liquid crystal self-assembly mechanism, Arch Rational Mech Anal (2018) 227:367.

DEPARTAMENTO DE FÍSICA, FCFM, UNIVERSIDAD DE CHILE, CASILLA 487-3, SANTIAGO, CHILE. *E-mail address*: marcel@dfi.uchile.cl

Departamento de Ingeniería Matemática and Centro de Modelamiento Matemático (UMI 2807 CNRS), Universidad de Chile, Casilla 170 Correo 3, Santiago, Chile.

 $E\text{-}mail\ address:$ kowalczy@dim.uchile.cl

Centro de Modelamiento Matemático (UMI 2807 CNRS), Universidad de Chile, Casilla 170 Correo 3, Santiago, Chile.

E-mail address: psmyrnelis@dim.uchile.cl