

MULTIPHASE SOLUTIONS TO THE VECTOR ALLEN-CAHN EQUATION: CRYSTALLINE AND OTHER COMPLEX SYMMETRIC STRUCTURES

PETER W. BATES, GIORGIO FUSCO, AND PANAYOTIS SMYRNELIS

ABSTRACT. We present a systematic study of entire symmetric solutions $u : \mathbb{R}^n \rightarrow \mathbb{R}^m$ of the vector Allen-Cahn equation $\Delta u - W_u(u) = 0$, $x \in \mathbb{R}^n$, where $W : \mathbb{R}^m \rightarrow \mathbb{R}$ is smooth, symmetric, nonnegative with a finite number of zeros, and $W_u := (\partial W / \partial u_1, \dots, \partial W / \partial u_m)^\top$. We introduce a general notion of equivariance with respect to a homomorphism $f : G \rightarrow \Gamma$ (G, Γ reflection groups) and prove two abstract results, concerning the cases of G finite and G discrete, for the existence of equivariant solutions. Our approach is variational and based on a mapping property of the parabolic vector Allen-Cahn equation and on a pointwise estimate for vector minimizers.

1. INTRODUCTION

We study bounded solutions $u : \mathbb{R}^n \rightarrow \mathbb{R}^m$ to the vector Allen-Cahn equation

$$(1.1) \quad \Delta u - W_u(u) = 0, \quad x \in \mathbb{R}^n,$$

where $W : \mathbb{R}^m \rightarrow \mathbb{R}$ is a smooth nonnegative multi-well potential that vanishes at $N \geq 1$ distinct points $a_1, a_2, \dots, a_N \in \mathbb{R}^m$, and $W_u := (\partial W / \partial u_1, \dots, \partial W / \partial u_m)^\top$. The stationary Allen-Cahn equation is formally the L^2 Euler-Lagrange equation associated to the functional

$$(1.2) \quad J_\Omega(u) := \int_\Omega \left(\frac{1}{2} |\nabla u|^2 + W(u) \right) dx,$$

defined on each bounded domain $\Omega \subset \mathbb{R}^n$ for $u \in W^{1,2}(\Omega, \mathbb{R}^m)$.

A complete description of the set of entire solutions of (1.1), even in the scalar case $m = 1$ in spite of many interesting and deep results (see for example [8] [18] [9] and the references therein), appears to be an impossible task. In this paper we are mostly interested in the vector case $m > 1$. We focus on *symmetric* potentials and proceed to a systematic discussions of symmetric solutions that can be determined by minimization of the energy (1.2).

The restriction to the symmetric setting has both physical and mathematical motivations. From the physical point of view we observe that symmetry is ubiquitous in nature and that the modeling of systems (e.g. materials) that can exist in different symmetric crystalline phases requires the use of vector *order parameters* [7]. Symmetric structures are observed at the junction of three or four coexisting phases in physical space. Similar structures appear at the singularities of soap films and compounds of soap bubbles.

2000 *Mathematics Subject Classification.* 35J91, 35J50.

Key words and phrases. vector Allen-Cahn equation, reflection groups, homomorphism, equivariant, positive map.

PWB was supported in part by NSF DMS-0908348, DMS-1413060, and the IMA. GF was partially supported by the IMA. PS was partially supported through the project PDEGE Partial Differential Equations Motivated by Geometric Evolution, co-financed by the European Union European Social Fund (ESF) and national resources, in the framework of the program Aristeia of the Operational Program Education and Lifelong Learning of the National Strategic Reference Framework (NSRF).

From the mathematical point of view, as we discuss in Sec.2, symmetry plays an essential role in the derivation of pointwise estimate for solutions of (1.1). Indeed by exploiting the symmetry we show, as in [1] and [3], the existence of minimizers of J_Ω that map *fundamental domains* in the domain x -space into *fundamental domains* in the target u -space. A basic consequence of this is the existence of minimizers that in certain sub-domains avoid all the minima of W but one. It follows that in each such sub-domain the potential W can be considered to have a unique global minimum. This is a key point since, for potentials with two or more global minima, it is very difficult to determine *a priori* in which subregions a minimizer u of J_Ω is near to one or another of the minima of W . Even in the scalar case, $m = 1$, for a potential with two minima this has proved to be a very difficult question. For instance, this is one of the main difficulties in the problem of characterizing the level sets of solutions $u : \mathbb{R}^n \rightarrow \mathbb{R}$ of the equation

$$(1.3) \quad \Delta u = u^3 - u, \quad x \in \mathbb{R}^n,$$

that satisfy the bound $|u| \leq 1$ and the conditions

$$(1.4) \quad \frac{\partial u}{\partial x_n} > 0,$$

a problem related to a celebrated conjecture of De Giorgi [10].

The study of equation (1.1) under symmetry hypotheses on the potential W was initiated in [6] and [5] where the existence of a solution $u : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ of (1.1) with the symmetry of the equilateral triangle was established. Existence of solutions with the symmetry of the regular tetrahedron was later proved in [14] for $n = m = 3$. These solutions are known as the triple and the quadruple *junction* states, respectively and, as indicated before with reference to soap films, are related to minimal surface complexes and in particular to the local structure of singularities where three sheets meet along a line or where four of these lines meet at a point [20]. Existence of solutions $u : \mathbb{R}^n \rightarrow \mathbb{R}^m$ equivariant with respect to a generic finite reflection group G acting both on the domain and on the target space was studied in [1], [2] and [11].

We introduce an abstract notion of equivariance of maps that includes as a special case the notion considered in [5], [14] and [1]. We assume there is a finite or discrete (infinite) reflection group G acting on \mathbb{R}^n and a finite reflection group Γ acting on \mathbb{R}^m and assume there exists a homomorphism $f : G \rightarrow \Gamma$ between G and Γ . For the concept of fundamental domain and for the general theory of reflection groups we refer to [13] and [16]. We define a map $u : \mathbb{R}^n \rightarrow \mathbb{R}^m$ to be f -equivariant if

$$(1.5) \quad u(gx) = f(g)u(x), \quad \text{for } g \in G, \quad x \in \mathbb{R}^n.$$

We characterize the homomorphisms which allow for the existence of f -equivariant maps that map a fundamental domain F for the action of G on \mathbb{R}^n into a fundamental domain Φ for the action of Γ on \mathbb{R}^m :

$$(1.6) \quad u(\overline{F}) \subset \overline{\Phi}.$$

We refer to these homomorphism and to the maps that satisfy (1.6) as *positive*. Positive homomorphisms (see Definition 1 below) have certain mapping properties that relate the projections associated to the walls of a fundamental domain F to the projections associated to the walls of a corresponding region Φ . These properties are instrumental to show that minimizing in the class of f -equivariant maps that satisfy (1.6) does not affect the Euler-Lagrange equation and renders a smooth solution of (1.1). The proof of this fact is based on a quite sophisticated use of the maximum principle for parabolic equations that was first introduced in [19] and [3]. We prove (see Lemma 4.2) that, provided f is a positive homomorphism, the L^2 gradient flow associated to the functional

(1.2) preserves the positivity condition (1.6). By a careful choice of certain scalar projections of the vector parabolic equation that describes the above mentioned gradient flow, we show that this fact is indeed a consequence of the maximum principle. Based on this and on a pointwise estimate from [11] and [12] we prove two abstract existence results: Theorem 3.1 which concerns the case where G is a finite reflection group and Theorem 3.2 that treats the case of a discrete (infinite) group G .

From (1.6) and the f -equivariance of u it follows

$$(1.7) \quad u(g\bar{F}) \subset f(g)\bar{\Phi}, \text{ for } g \in G.$$

Therefore, besides its importance for the proofs of Theorem 3.1 and Theorem 3.2, the mapping property (1.6) is a source of information on the geometric structure of the vector valued map u . The fact that (1.7) holds true in general in the abstract setting of the present analysis can perhaps be regarded as a major result of our work. Indeed due to the variety of choices for n and m , the dimensions of domain and target space, of the possible choices of the reflection groups G and Γ , and of the homomorphism $f : G \rightarrow \Gamma$, we will deduce from Theorem 3.1 and Theorem 3.2 the existence of various complex multi-phase solutions of (1.1) including several types of lattice solutions. A characterization of all homomorphisms between reflection groups in general dimensions is not known. For the special case $n = m = 2$, in the Appendix, we determine all positive homomorphisms between finite reflection groups and the corresponding solutions of (1.1).

The paper is organized as follows. In Sec.2 we discuss the notion of f -equivariance, the concept of *positive* homomorphism $f : G \rightarrow \Gamma$ and give a few examples. In Sec.3 we list the hypotheses on the potential W and state the main results, Theorem 3.1 and Theorem 3.2, that we prove in Sec.4. In Sec.5 and in Sec.6 we apply the abstract theorems proved in Sec.4 to specific situations and present a series of solutions of (1.1).

We denote by $\langle x, y \rangle$ the standard inner product of $x, y \in \mathbb{R}^k$, $k \geq 1$, by $|x| = \sqrt{\langle x, x \rangle}$ the norm of $x \in \mathbb{R}^k$ and by $d(x, A) = \inf_{y \in A} |y - x|$ the distance of x from $A \subset \mathbb{R}^k$.

2. f -EQUIVARIANCE AND THE NOTION OF POSITIVE HOMOMORPHISM

Here and in the following section, together with the abstract concepts and general proofs, we present simple but significant examples and special cases to help the reader to build a deep understanding of the paper. We begin with some examples of f -equivariant maps. We let I_k the identity map of \mathbb{R}^k , $k \geq 1$. As a first example we observe that, in the particular case where $\Gamma = G$ and the homomorphism f is the identity, f -equivariance reduces to the notion considered in [5], [14], [1], [2], and [11]:

$$u(gx) = gu(x), \text{ for } g \in G, x \in \mathbb{R}^n.$$

The next example is a genuine f -equivariant map. In [4], under the assumption that W is invariant under the group Γ of the equilateral triangle, we constructed a solution $u : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ to system (1.1) such that

- (i) $u(\gamma x) = \gamma u(x)$ for all $\gamma \in \Gamma$ (which is the dihedral group D_3).
- (ii) $u(-x) = u(x)$ for all $x \in \mathbb{R}^2$.

If we incorporate the additional symmetry (ii) in a group structure, this solution can be seen as an f -equivariant map. Indeed, the regular hexagon reflection group $G = D_6$ contains $\Gamma = D_3$, and the antipodal map $\sigma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by $\sigma(x) = -x$. Since σ commutes with the elements of D_3 , G is isomorphic to the group product $\{I_2, \sigma\} \times D_3$. Furthermore, we can define a homomorphism $f : D_6 = \{I_2, \sigma\} \times D_3 \rightarrow D_3$, by setting $f(\gamma) = \gamma$ and $f(\sigma\gamma) = \gamma$, for every $\gamma \in D_3$. Then, the above conditions (i) and (ii) express the f -equivariance of the solution u in [4].

Similarly, we can consider the action on \mathbb{R}^2 of the discrete reflection group G' generated by the reflections s_1 , s_2 and s_3 with respect to the corresponding lines $P_1 := \{x_2 = 0\}$, $P_2 := \{x_2 = x_1/\sqrt{3}\}$ and $P_3 := \{x_2 = -\sqrt{3}(x_1 - 1)\}$ (the dashed lines in Figure 1). These three lines bound a triangle F' with angles 30, 60 and 90 degrees, which is a fundamental domain of G' . The discrete group G' contains also all the reflections with respect to the lines drawn in Figure 1, which partition the plane into triangles congruent to F' .




FIGURE 1. The discrete reflection group G' on the left and the finite reflection group $\Gamma = D_3$ on the right.

The point group of G' , that is the stabilizer of the origin: $\{g \in G' : g(0) = 0\}$, is the group $G = D_6$, and we have $G' = TG$, where T is the translation group of G' . T is generated by the translations t^\pm by the vectors $(\frac{3}{2}, \pm\frac{\sqrt{3}}{2})$. Now, if we compose the canonical homomorphism $p : G' \rightarrow G$ such that $p(tg) = g$ for every $t \in T$ and $g \in G$, with the homomorphism $f : D_6 \rightarrow D_3$ defined in the previous paragraph, we obtain a homomorphism $f' : G' \rightarrow D_3$. We have in particular

$$\begin{aligned}
 f'(s_1) &= f(p(s_1)) = s_1, \\
 f'(s_3) &= f(p(s_3)) = p(s_3), \\
 f'(s_2) &= f'(\sigma p(s_3)) = f(\sigma p(s_3)) = p(s_3),
 \end{aligned}
 \tag{2.1}$$

where $p(s_3)$ is the reflection in the line $\Pi_2 = \{u_2 = -\sqrt{3}u_1\}$. We note that the image of the line $P_1 = \{x_2 = 0\}$ by an f' -equivariant map $u : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is contained in the line $\Pi_1 := \{u_2 = 0\}$ while the images of the lines $P_2 = \{x_2 = x_1/\sqrt{3}\}$ and $P_3 = \{x_2 = -\sqrt{3}(x_1 - 1)\}$ are contained in the line $\Pi_2 := \{u_2 = -\sqrt{3}u_1\}$. Indeed

$$\begin{aligned}
 x = s_1x &\Rightarrow u(x) = u(s_1x) = f'(s_1)u(x) = s_1u(x), \\
 x = s_2x &\Rightarrow u(x) = u(s_2x) = f'(s_2)u(x) = p(s_3)u(x), \\
 x = s_3x &\Rightarrow u(x) = u(s_3x) = f'(s_3)u(x) = p(s_3)u(x).
 \end{aligned}
 \tag{2.2}$$

The lines Π_1 and Π_2 define a 60 degree sector Φ which is a fundamental domain of the finite reflection group D_3 . At a later stage, we will prove the existence of a solution to (1.1) that maps the triangle F' in this sector.

Now we return to the general setting and discuss the notion of positive homomorphism $f : G \rightarrow \Gamma$ between reflection groups G and Γ . Before giving the definition we observe that if $s \in G$ is a reflection we have $I_m = f(I_n) = f(s)f(s)$. This and the fact that $f(s)$ is an orthogonal transformation imply that $f(s)$ is symmetric. Thus $f(s)$ has m ortho-normal eigenvectors ν_1, \dots, ν_m and $\nu_j = f(s)f(s)\nu_j = \mu_j^2\nu_j$ implies that $|\mu_j| = 1$ for the corresponding eigenvalues μ_j , $j = 1, \dots, m$. Therefore if we let $E \subset \mathbb{R}^m$ be the span of the eigenvectors corresponding to the eigenvalue $\mu = 1$, that is, $E = \ker(f(s) - I_m)$ the set of the points fixed by $f(s)$, we have $\mathbb{R}^m = E \oplus E^\perp$ and

$$(2.3) \quad f(s)u = f(s)(u_E + (u - u_E)) = u_E - (u - u_E) = -u + 2u_E,$$

where we have used the decomposition $u = u_E + (u - u_E)$ with $u_E \in E$ and $u - u_E \in E^\perp$. We can interpret (2.3) by saying that $f(s)$ is a projection with respect to the subspace E or that $f(s)$ coincides with I_m on E and with the antipodal map on E^\perp .

Definition 1. Let F be a fundamental domain of G , bounded by the hyperplanes P_1, \dots, P_l , corresponding to the reflections s_1, \dots, s_l . We say that a homomorphism $f : G \rightarrow \Gamma$ is *positive* if there exists a fundamental domain Φ of Γ , bounded by the hyperplanes Π_1, \dots, Π_k , such that for every $i = 1, \dots, l$, there is $1 \leq k_i \leq k$ and $\tilde{\Pi}_1, \dots, \tilde{\Pi}_{k_i} \in \{\Pi_1, \dots, \Pi_k\}$ such that

$$(2.4) \quad \ker(f(s_i) - I_m) = \cap_{j=1}^{k_i} \tilde{\Pi}_j.$$

That is, the set of points fixed by the orthogonal map $f(s_i)$ is one of the hyperplanes Π_j , or the intersection of several of them.

The property of being positive for a homomorphism f is independent of the choice of F . Indeed, if we take $\hat{F} = gF$, with $g \in G$, then \hat{F} is bounded by the hyperplanes gP_1, \dots, gP_l , corresponding to the reflections $gs_1g^{-1}, \dots, gs_lg^{-1}$. In addition $\ker(f(gs_ig^{-1}) - I_m) = f(g)\ker(f(s_i) - I_m)$, thus, the fundamental domain $\hat{\Phi} = f(g)\Phi$ can be associated with \hat{F} , in accordance with the definition.

Note that the choice of Φ is not unique, since the homomorphism f can associate F to Φ , or to $-\Phi$ indifferently.

The homomorphism $f' : G' \rightarrow D_3$ defined above is an example of a positive homomorphism. Indeed, if we identify F with the triangle F' and Φ with the 60 degree sector $\{-\sqrt{3}u_1 < u_2 < 0, u_1 > 0\}$ bounded by the lines Π_1 and Π_2 , then (2.1) expresses the positivity of f' .

It is not true in general that a homomorphism $f : G \rightarrow \Gamma$ between reflection groups G and Γ is positive. For example the canonical projection p of a discrete reflection group G' onto its point group G does not, in general, fulfill this requirement. To see this, let us revisit the discrete reflection group G' depicted in Figure 1. We have

$$\begin{aligned} p(s_1) &= \text{reflection in the line } \Pi_1, \\ p(s_2) &= \text{reflection in the line } \{u_2 = u_1/\sqrt{3}\}, \\ p(s_3) &= f(s_3) = \text{reflection in the line } \{u_2 = -\sqrt{3}u_1\}, \end{aligned}$$

then $p(s_i), i = 1, 2, 3$ are reflections with respect to three distinct lines intersecting at the origin. Thus, the canonical projection $p : G' \rightarrow G = D_6$ cannot associate F' to any fundamental domain of D_6 (a 30 degree sector).

3. THE THEOREMS

We assume:

Hypothesis 1 (Homomorphism). *There exist: a finite (or discrete) reflection group G acting on \mathbb{R}^n , a finite reflection group Γ acting on \mathbb{R}^m , and a homomorphism $f : G \rightarrow \Gamma$ which is positive in the sense of Definition 1. We denote by Φ the fundamental domain of Γ that f associates with the fundamental domain F of G .*

Hypothesis 2. *The potential $W : \mathbb{R}^m \rightarrow [0, \infty)$, of class C^3 , is invariant under the finite reflection group Γ , that is,*

$$(3.1) \quad W(\gamma u) = W(u), \text{ for all } \gamma \in \Gamma \text{ and } u \in \mathbb{R}^m.$$

Moreover, we assume that there exists $M > 0$ such that $W(su) \geq W(u)$, for $s \geq 1$ and $|u| = M$.

Hypothesis 3. *There exist $a \in \overline{\Phi}$, the closure of Φ , and $q^* > 0$ such that:*

- (i) $0 = W(a) < W(u)$, for $u \in \overline{\Phi} \setminus \{a\}$ and
- (ii) for each $\nu \in \mathbb{S}^{m-1}$, $\mathbb{S}^{m-1} \subset \mathbb{R}^m$ the unit sphere, the map $(0, q^*] \ni q \rightarrow W(a + q\nu)$ has a strictly positive second derivative.

Hypothesis 2 and 3 determine the number N of minima of W . From Hypothesis 2 we have

$$W(\gamma a) = 0, \text{ for } \gamma \in \Gamma.$$

Therefore, if $a \in \Phi$, that is, a is in the interior of Φ , from the fact that $\gamma\Phi \neq \Phi$ for $\gamma \in \Gamma \setminus \{I_m\}$ it follows that W has exactly $N = |\Gamma|$ distinct minima, where $|\mathcal{G}|$ denotes the order of a group \mathcal{G} . If $a \in \partial\Phi$ then the stabilizer $\Gamma_a = \{\gamma \in \Gamma : \gamma a = a\}$ of a is nontrivial and we have $N = |\Gamma|/|\Gamma_a| < |\Gamma|$ and a is the unique minimum of W in the cone $\mathcal{D} \subset \mathbb{R}^m$ defined by

$$\mathcal{D} = \text{Int } \cup_{\gamma \in \Gamma_a} \gamma \overline{\Phi}.$$

The set \mathcal{D} satisfies

$$(3.2) \quad \text{for } \gamma \in \Gamma : \text{ either } \gamma\mathcal{D} \cap \mathcal{D} = \emptyset \text{ or } \gamma\mathcal{D} = \mathcal{D}.$$

It follows that

$$\mathbb{R}^m = \cup_{\gamma \in \Gamma} \gamma \overline{\mathcal{D}},$$

that is, \mathbb{R}^m is partitioned into $N = |\Gamma|/|\Gamma_a|$ cones congruent to \mathcal{D} . The cone $\mathcal{D} \subset \mathbb{R}^m$ has its counterpart in the set $D \subset \mathbb{R}^n$ given by

$$(3.3) \quad D = \text{Int } (\cup_{g \in f^{-1}(\Gamma_a)} g \overline{F})$$

which is mapped into $\overline{\mathcal{D}}$ by any positive f -equivariant map $u : \mathbb{R}^n \rightarrow \mathbb{R}^m$. Indeed, for such a map (1.7) implies that $u(g\overline{F}) \subset \overline{\mathcal{D}}$ if and only if $f(g) \in \Gamma_a$ or equivalently $g \in f^{-1}(\Gamma_a)$.

In the case $G = \Gamma$ and $f = I$, the expression for D reduces to $D = \text{Int } (\cup_{g \in \Gamma_a} g \overline{F})$, considered in [1], [2], and [11].

The set D satisfies the analog of (3.2). Therefore also the domain space \mathbb{R}^n is partitioned into $N = |\Gamma|/|\Gamma_a|$ sets congruent to D .

Let us consider a few examples

- (i) if $G = \Gamma$, f is the identity and a is in the interior of Φ , then $\mathcal{D} = \Phi$ and $D = F$.

- (ii) if $m = n = 2$, $G = \Gamma = D_3$, $\Phi = \{u : 0 < u_2 < \sqrt{3}u_1, u_1 > 0\}$, f is the identity and $a = (1, 0)$, then $\Gamma_a = \{I_2, g_1\}$ where g_1 is the reflection in the line $\{u_2 = 0\}$. Therefore (3.3) yields $D = \text{Int}(\overline{F} \cup g_1 \overline{F}) = \{x : |x_2| < \sqrt{3}x_1, x_1 > 0\}$.
- (iii) if $n = m = 2$, $G = D_6 = \{I_2, \sigma\} \times D_3$, $\Gamma = D_3$, $f(\gamma) = \gamma$ and $f(\sigma\gamma) = \gamma$ for every $\gamma \in D_3$, and $a \in \Phi = \{u : 0 < u_2 < \sqrt{3}u_1, u_1 > 0\}$, then $\Gamma_a = \{I_2\}$ and $f^{-1}(\Gamma_a) = \{I_2, \sigma\}$. Therefore $\mathcal{D} = \Phi$ and $D = \text{Int}(\overline{F} \cup \sigma \overline{F}) = \{x : 0 < -\frac{x_1 x_2}{|x_1|} < \frac{|x_1|}{\sqrt{3}}, x_1 \neq 0\}$. ($F = \{x : 0 < -x_2 < \frac{x_1}{\sqrt{3}}, x_1 > 0\}$)
- (iv) If in the previous example we take $a = (1, 0) \in \overline{\Phi}$ we have $\Gamma_a = \{I_2, g_1\}$ and $f^{-1}(\Gamma_a) = \{I_2, \sigma, g_1, \sigma g_1\}$. It follows $\mathcal{D} = \text{Int}(\overline{\Phi} \cup g_1 \overline{\Phi}) = \{0 < |u_2| < \sqrt{3}u_1, u_1 > 0\}$ and $D = \text{Int}(\overline{F} \cup \sigma \overline{F} \cup g_1 \overline{F} \cup \sigma g_1 \overline{F}) = \{x : 0 < |x_2| < \frac{|x_1|}{\sqrt{3}}\}$.




FIGURE 2. The sets F , Φ , D and \mathcal{D} and their correspondence by an f -equivariant map in the the examples (i)-(iv).

If G is a discrete (infinite) group, then D has infinitely many connected components. As examples (iii) and (iv) above show, even when G is a finite group, D does not need to be connected. To characterize one of the connected components of D , let $G^a \subset f^{-1}(\Gamma_a)$ be the subgroup generated by $f^{-1}(\Gamma_a) \cap \{s_1, \dots, s_l\}$ and define

$$(3.4) \quad D_0 := \text{Int} \cup_{g \in G^a} g \overline{F}.$$

Since G^a is a reflection group generated by a subset of $\{s_1, \dots, s_l\}$, the set of reflections in the planes that bound F , the set D_0 is connected. To show that D_0 is one of the connected components of D we show that D_0 and $D \setminus D_0$ are disconnected. This is equivalent to prove that if s is the reflection in one of the planes that bound D_0 , then $s \notin f^{-1}(\Gamma_a)$. The definition of D_0 implies $s = s_i s^0 s_i$ for some $s_i \in G^a$ and some reflection $s^0 \in \{s_1, \dots, s_l\} \setminus f^{-1}(\Gamma_a)$ and therefore $s^0 = s_i s s_i \Rightarrow s^0 \in G^a$ if $s \in f^{-1}(\Gamma_a)$.

For examples (i)-(iv) we have: $F = D_0 = D$ in (i); $F \subsetneq D_0 = D$ in (ii); $F = D_0 \subsetneq D = D_0 \cup \sigma D_0$ in (iii); $F \subsetneq D_0 = \text{Int}(\overline{F} \cup g_1 \overline{F}) \subsetneq D = D_0 \cup \sigma D_0$ in (iv). Figure 2 illustrates these properties.

We are now in a position to state the main results.

Theorem 3.1. *Under Hypotheses 1–3, and assuming that G is a finite reflection group, there exists an f -equivariant classical solution u to system (1.1), and a positive decreasing map $q : [0, +\infty) \rightarrow \mathbb{R}$ with $\lim_{r \rightarrow +\infty} q(r) = 0$, such that*

- (i) $u(\overline{F}) \subset \overline{\Phi}$ and $u(\overline{D}) \subset \overline{\mathcal{D}}$,
- (ii) $|u(x) - a| \leq q(d(x, \partial D))$, for $x \in D$,
- (iii) *if the matrix $D^2W(a)$ is positive definite, then $q(r) \leq Ke^{-kr}$ for some constants $k, K > 0$ and therefore $|u(x) - a| \leq Ke^{-kd(x, \partial D)}$, for $x \in D$.*

Theorem 3.2. *Under Hypotheses 1–3, and assuming that G is a discrete reflection group, there exists for every $R > R_0$ (a positive constant), a nontrivial f -equivariant classical solution u_R to system*

$$(3.5) \quad \Delta u_R - R^2 W_u(u_R) = 0, \text{ for } x \in \mathbb{R}^n$$

such that

- (i) $u_R(\overline{F}) \subset \overline{\Phi}$ and $u_R(\overline{D}) \subset \overline{\mathcal{D}}$.
- (ii) $|u_R(x) - a| \leq q(Rd(x, \partial D))$, for $x \in D$ where q is as in Theorem 3.1,
- (iii) $|u_R(x) - a| \leq Ke^{-kRd(x, \partial D)}$, for $x \in D$,
for positive constants k, K , provided $D^2W(a)$ is positive definite.

The solution u_R of (3.5) given by Theorem 3.2 is periodic. We describe this periodic structure of u_R under the assumption that the positive homomorphism f satisfies ¹

$$(3.6) \quad f(t) = I_m, \text{ for } t \in T.$$

This covers the examples that we present below. On the other hand we are not aware of positive homomorphisms that do not satisfy (3.6). Assuming (3.6), if $G = TG_0$ with G_0 the point group of G and T its translation group we have

$$u_R(tx) = u_R(x), \text{ for } t \in T, x \in C$$

where

$$C = \text{Int} \cup_{g \in G_0} g\overline{F}$$

is the elementary cell. C is a convex polytope that satisfies

$$tC \cap C = \emptyset, \text{ for } t \in T \setminus \{I_n\}$$

and defines a tessellation of \mathbb{R}^n as the union of the translations tC , $t \in T$ of C : $\mathbb{R}^n = \overline{\cup_{t \in T} tC}$. In this sense we can say that u_R has a *crystalline* structure and that C is the elementary *crystal*.

Let us illustrate Theorem 3.2 with the help of the example described in Sec.2, where the discrete reflection group G' acts on the domain x -plane while the finite reflection group $\Gamma = D_3$ acts on the target u -plane. We have already verified that the homomorphism $f' : G' \rightarrow \Gamma$ is positive and therefore Hypothesis 1 is satisfied. When Hypotheses 2–3 also hold, Theorem 3.2 ensures the existence, for every R sufficiently large, of a nontrivial f' -equivariant solution u_R to (3.5), such that $u_R(\overline{F'}) \subset \overline{\Phi}$. By f' -equivariance, the other fundamental domains of G' are mapped into

¹ For an example of a homomorphism that does not satisfy (3.6) take $\Gamma = \{I_m, \gamma\}$ with γ the reflection in the plane $\{u_1 = 0\}$ and $G = \langle s_j \rangle_{j \in \mathbb{Z}} = TG_0$ where s_j is the reflection in the plane $\{x_1 = j\}$, T is the translation group generated by the translation t_0 by the vector $(2, 0, \dots, 0)$ and $G_0 = \{I_n, s_0\}$. Define $f : G \rightarrow \Gamma$ by

$$\begin{aligned} f(t_0) &= f(s_{2j}) = \gamma, \\ f(s_{2j+1}) &= I_m. \end{aligned}$$

fundamental domains of Γ as in Figure 3. Properties (ii) and (iii) state that for every $x \in D$, $u_R(x)$ approaches as R grows, the unique minimum a of W in $\bar{\Phi}$, with a speed that depends on $d(x, \partial D)$. If for instance the potential W has six minima (one in the interior of each fundamental domain of D_3), then the set D is the union of the fundamental domains of G' with the same colour (cf. Figure 3 left) and \mathcal{D} is the sector with the same color of D (cf. Figure 3 right). If a lies on the boundary of the fundamental domain of D_3 , for instance $a = (1, 0)$ then \mathcal{D} is the 120 degree sector that contains a and D is the union of all fundamental domains (triangles) with the same colors of the two sectors that compose \mathcal{D} . For this example condition (3.6) is satisfied and the elementary crystal C , is the hexagon determined by the fundamental domains (triangles) whose closure contains $0 \in \mathbb{R}^2$.




FIGURE 3. Fundamental domains for the actions on \mathbb{R}^2 of G' (left) and D_3 (right). The f' -equivariant solution u_R of (1.1) given by Theorem 3.2 maps fundamental domains into fundamental domains with the same color.

To give a first application of Theorem 3.1, consider the particular case where $G = D_6 = \{I_2, \sigma\} \times D_3$, $\Gamma = D_3$, and f the positive homomorphism defined by $f(\gamma) = \gamma$, $f(\sigma\gamma) = \gamma$, for every $\gamma \in D_3$. Figure 2 (iii) and (iv) shows the correspondence of the fundamental domains by f -equivariant solutions u of (1.1) when the potential W has respectively six and three minima. The case of W with three minima when u has a six-fold structure, see Figure 2 (iv) was first considered in [4]. Other examples of application of Theorem 3.1 and Theorem 3.2 are given in Sec.5 and Sec.6.

4. PROOFS OF THEOREM 3.1 AND THEOREM 3.2

Step 1 (Minimization).

In the case where G is a finite reflection group, we first construct the solution in a ball $B_R \subset \mathbb{R}^n$ of radius $R > 0$ centered at the origin. We set $F_R = F \cap B_R$ and define the class

$$A^R := \{u \in W^{1,2}(B_R, \mathbb{R}^m) : u \text{ is } f\text{-equivariant and } u(\bar{F}_R) \subset \bar{\Phi}\}$$

in which we have imposed the positivity constraint $u(\bar{F}_R) \subset \bar{\Phi}$. Then, we consider the minimization problem

$$\min_{A^R} J_{B_R}, \text{ where } J_{B_R}(u) = \int_{B_R} \left(\frac{1}{2} |\nabla u|^2 + W(u) \right) dx.$$

Since A^R is convex and hence weakly closed in $W^{1,2}(B_R, \mathbb{R}^m)$, a minimizer u_R exists, and because of Hypothesis 2 we can assume that

$$(4.1) \quad |u_R(x)| \leq M.$$

In the case where G is a discrete reflection group, we can work directly in the fundamental domain F . Suppose first that F is bounded. Then, we consider the class

$$A := \{u \in W_{\text{loc}}^{1,2}(\mathbb{R}^n, \mathbb{R}^m) : u \text{ is } f\text{-equivariant and such that } u(\overline{F}) \subset \overline{\Phi}\},$$

and choose a minimizer u_R of the problem

$$\min_A J_F^R, \text{ where } J_F^R(u) = \int_F \left(\frac{1}{2} |\nabla u|^2 + R^2 W(u) \right) dx,$$

satisfying the estimate (4.1).

Now, suppose that F is not bounded. This implies that all the reflection hyperplanes of G are parallel to a subspace $\{0\}^\nu \times \mathbb{R}^d \subset \mathbb{R}^n$ (with $\nu + d = n$, $d \geq 1$), and that G also acts in \mathbb{R}^ν . Since $F = F_\nu \times \mathbb{R}^d$, with $F_\nu \subset \mathbb{R}^\nu$ bounded, we have, according to the preceding argument, a minimizer $v_R : \mathbb{R}^\nu \rightarrow \mathbb{R}^m$ of

$$J_{F_\nu}^R(v) = \int_{F_\nu} \left(\frac{1}{2} |\nabla_{x_\nu} v|^2 + R^2 W(v) \right) dx_\nu,$$

in the analog of A with n replaced by ν , that is, the class of $W_{\text{loc}}^{1,2}(\mathbb{R}^\nu, \mathbb{R}^m)$ maps v , which are f -equivariant and satisfy $v(\overline{F_\nu}) \subset \overline{\Phi}$. Then, we set $u_R(x) := v_R(x_\nu)$, where $x = (x_\nu, x_d) \in \mathbb{R}^n$.

Step 2 (Removing the positivity constraint with the gradient flow).

To show that the positivity constraint built in A^R (or A) does not affect the Euler-Lagrange equation we will utilize the gradient flow associated to the elliptic system. In the case where G is a finite reflection group we consider

$$(4.2) \quad \begin{cases} \frac{\partial u}{\partial t} = \Delta u - W_u(u), & \text{in } B_R \times (0, \infty), \\ \frac{\partial u}{\partial \mathbf{n}} = 0, & \text{on } \partial B_R \times (0, \infty), \\ u(x, 0) = u_0(x), & \text{in } B_R, \end{cases}$$

where $\partial/\partial \mathbf{n}$ is the normal derivative.

In the case where G is a discrete reflection group, we consider

$$(4.3) \quad \begin{cases} \frac{\partial u}{\partial t} = \Delta u - R^2 W_u(u), & \text{in } \mathbb{R}^n \times (0, \infty), \\ u(x, 0) = u_0(x), & \text{in } \mathbb{R}^n. \end{cases}$$

Since W is C^3 , the results in [15] apply and provide a unique solution to (4.2) (or (4.3)) which is smooth if we assume that u_0 is globally Lipschitz. In the next two lemmas we will establish that the gradient flow preserves the f -equivariance and the positivity of a smooth initial condition.

Lemma 4.1. *Under Hypothesis 2, if the initial condition u_0 is a smooth, f -equivariant map, then for every $t > 0$, the solution $u(\cdot, t)$ of problem (4.2) (or (4.3)) is also f -equivariant.*

Proof. We only write the proof for (4.2), since it is identical for (4.3). Let $g \in G$ and $\gamma := f(g) \in \Gamma < O(\mathbb{R}^m)$. We are going to check that for every $x \in B_R$ and every $t > 0$, we have $u(gx, t) = \gamma u(x, t)$ or equivalently $u(x, t) = \gamma^\top u(gx, t)$. Let us set $v(x, t) := \gamma^\top u(gx, t)$. Since g is an isometry, $\Delta v(x, t) = \gamma^\top (\Delta u)(gx, t)$. On the other hand, we have $\frac{\partial v}{\partial t}(x, t) = \gamma^\top \frac{\partial u}{\partial t}(gx, t)$, and utilizing the symmetry of the potential: $W_u(v(x, t)) = \gamma^\top W_u(u(gx, t))$. Finally, we see that for $x \in \partial B_R$ and $t > 0$, $\frac{\partial v}{\partial \mathbf{n}}(x, t) = \gamma^\top \frac{\partial u}{\partial g \mathbf{n}}(gx, t) = 0$. Thus, v is also a smooth solution of (4.2) with initial condition $v_0(x) = \gamma^\top u_0(gx) = u_0(x)$, and by uniqueness $u \equiv v$. \square

Lemma 4.2. *Under Hypotheses 1–2, and assuming that the initial condition u_0 is a smooth map, we have:*

- $u_0 \in A^R \Rightarrow u(\cdot, t) \in A^R, \forall t > 0$, when G is a finite reflection group.
- $u_0 \in A \Rightarrow u(\cdot, t) \in A, \forall t > 0$, when G is a discrete reflection group.

Proof. We first present the proof in a specific case where the argument can be described with simpler notation and then consider the abstract situation and give the proof for the general case. The case we discuss first is the example of Sec.2 where we have the discrete reflection group G' acting on the domain, $\Gamma = D_3$ acting on the target, and the homomorphism $f' : G' \rightarrow \Gamma$. We conserve the notation of Sec.2 and refer to Figure 1 and to the comments following Theorem 3.2. In particular, we still denote by F' the fundamental domain of G' and by f' the homomorphism $G' \rightarrow \Gamma$ (which are denoted by F and f in the statement of Theorem 3.2 and in the first step of its proof). We also denote by ρ_1 and ρ_2 the outward unit normals to the lines Π_1 and Π_2 that bound the fundamental domain Φ of Γ . Let $\Pi_3 = \{u_2 = \sqrt{3}u_1\}$ be the third reflection line of Γ and let $\rho_3 := (-\sqrt{3}/2, 1/2) \perp \Pi_3$. From (4.3) and the symmetry of W given by (3.1), it follows that for every $j = 1, 2, 3$, the projection $h_j(x, t) := \langle u(x, t), \rho_j \rangle$ satisfies the linear scalar equation:

$$\Delta h_j + c_j^* h_j - \frac{\partial h_j}{\partial t} = 0 \text{ in } \mathbb{R}^2 \times (0, +\infty),$$

with $c_j^* = R^2 c_j$ and c_j (cf. (4.7) below) continuous and bounded on $\mathbb{R}^2 \times [0, T]$, for every $T > 0$.

Now, suppose that for some $t_0 > 0$, $u(\cdot, t_0)$ does not belong to the class A .

In order to have equations with nonpositive coefficients, we do the standard transformation and set $\tilde{h}_j(x, t) := e^{-\lambda t} h_j(x, t)$, where the constant λ is chosen larger than $\sup\{c_j^*(x, t) \mid x \in \mathbb{R}^2, t \in [0, t_0], j = 1, 2, 3\}$. Then, we have

$$(4.4) \quad \Delta \tilde{h}_j + \tilde{c}_j^* \tilde{h}_j - \frac{\partial \tilde{h}_j}{\partial t} = 0 \text{ in } \mathbb{R}^2 \times (0, t_0], \text{ with } \tilde{c}_j^* = c_j^* - \lambda \leq 0.$$

Let $\mu := \max\{d(e^{-\lambda t} u(x, t), \bar{\Phi}) \mid x \in \bar{F}', t \in [0, t_0]\} > 0$, and suppose that this is achieved at $\tilde{x} \in \bar{F}'$ at time $\tilde{t} \in (0, t_0]$ (since $u_0 \in A$). Define

$$\tilde{u} := e^{-\lambda \tilde{t}} u(\tilde{x}, \tilde{t}), \quad \rho := \frac{\tilde{u} - \tilde{v}}{|\tilde{u} - \tilde{v}|},$$

where \tilde{v} is the unique point of $\partial \Phi$ (since Φ is convex) such that $d(\tilde{u}, \tilde{v}) = \mu$. According to the direction of ρ , we distinguish the following cases:

- If $\rho = \rho_1$, then $\tilde{v} \in \Pi_1 \cap \bar{\Phi}$ and we define $\omega := \{(x, t) \in \mathbb{R}^2 \times (0, t_0] : \langle e^{-\lambda t} u(x, t), \rho_1 \rangle > 0\}$. Clearly, $(\tilde{x}, \tilde{t}) \in \omega$ which is relatively open in $\mathbb{R}^2 \times (0, t_0]$.
- If $\rho = \rho_2$, then $\tilde{v} \in \Pi_2 \cap \bar{\Phi}$. Similarly, define $\omega := \{(x, t) \in \mathbb{R}^2 \times (0, t_0] : \langle e^{-\lambda t} u(x, t), \rho_2 \rangle > 0\}$, and we have $(\tilde{x}, \tilde{t}) \in \omega$ which is relatively open in $\mathbb{R}^2 \times (0, t_0]$.

- (iii) If $\rho = \rho_3$, then $\tilde{v} = 0$. We check that $(\tilde{x}, \tilde{t}) \in \omega := \{(x, t) \in \mathbb{R}^2 \times (0, t_0] : \langle e^{-\lambda t} u(x, t), \rho_3 \rangle > 0\}$, which is relatively open in $\mathbb{R}^2 \times (0, t_0]$.
- (iv) If $\rho = \alpha_1 \rho_1 + \alpha_3 \rho_3$ with $\alpha_1, \alpha_3 > 0$, then $\tilde{v} = 0$ and we define $\omega := \{(x, t) \in \mathbb{R}^2 \times (0, t_0] : \langle e^{-\lambda t} u(x, t), \rho_j \rangle > 0 \text{ for } j = 1 \text{ and } j = 3\}$. Thanks to the fact that $\langle \rho_1, \rho_3 \rangle \geq 0$, we have again $(\tilde{x}, \tilde{t}) \in \omega$ which is relatively open in $\mathbb{R}^2 \times (0, t_0]$.
- (v) If $\rho = \alpha_2 \rho_2 + \alpha_3 \rho_3$ with $\alpha_2, \alpha_3 > 0$, then $\tilde{v} = 0$ and we define ω in a similar way.

We want to apply the maximum principle to $\tilde{h}(x, t) := \langle e^{-\lambda t} u(x, t), \rho \rangle$ in a neighborhood of (\tilde{x}, \tilde{t}) . In the cases (i), (ii) and (iii) above, the equation (4.4) trivially holds in ω . In the cases (iv) and (v), we have the inequality

$$\Delta \tilde{h} + \tilde{c}^* \tilde{h} - \frac{\partial \tilde{h}}{\partial t} \geq 0 \text{ in } \omega, \text{ with } \tilde{c}^* = \max\{\tilde{c}_j^* \mid j = 1, 2, 3\} \leq 0.$$

Indeed, we can check that for instance in the case (iv):

$$\Delta \tilde{h} + \tilde{c}^* \tilde{h} - \frac{\partial \tilde{h}}{\partial t} = \alpha_1 (\tilde{c}^* - \tilde{c}_1^*) \tilde{h}_1 + \alpha_3 (\tilde{c}^* - \tilde{c}_3^*) \tilde{h}_3 \geq 0, \quad \forall (x, t) \in \omega.$$

At this stage, the fact that Φ is an acute angle sector (i.e. $\langle \rho_1, \rho_2 \rangle \leq 0$) is essential to conclude the proof. This property implies that

$$(4.5) \quad \tilde{u} \in \Pi_j \text{ (with } j = 1, 2) \Rightarrow \tilde{v}, \rho \in \Pi_j,$$

and as a consequence

$$(4.6) \quad f'(s_i) \notin \Gamma_\rho \Rightarrow \tilde{x} \notin P_i,$$

where P_i ($i = 1, 2, 3$) is a line bounding F' , corresponding to the reflection $s_i \in G'$. To show (4.6), suppose that $\tilde{x} \in P_i$. Then, $s_i(\tilde{x}) = \tilde{x}$, and by f' -equivariance

$$u(\tilde{x}, \tilde{t}) = u(s_i(\tilde{x}), \tilde{t}) = f'(s_i)u(\tilde{x}, \tilde{t}) \Leftrightarrow u(\tilde{x}, \tilde{t}) \in \ker(f'(s_i) - I_2).$$

Since $\ker(f'(s_i) - I_2)$ is either the line Π_1 or the line Π_2 (cf. Hypothesis 1), we deduce thanks to (4.5) that

$$\rho \in \ker(f'(s_i) - I_2) \Leftrightarrow f'(s_i) \in \Gamma_\rho.$$

Property (4.6) will enable us to locate \tilde{x} in $\overline{F'}$. Let \tilde{G}' be the subgroup of G' generated by the reflections s_i ($i = 1, 2, 3$) such that $f'(s_i) \in \Gamma_\rho$. By f' -equivariance we have

$$\mu = \tilde{h}(\tilde{x}, \tilde{t}) = \max\{\tilde{h}(x, t) : x \in \cup_{g \in \tilde{G}'} g \overline{F'}, t \in [0, t_0]\}.$$

But now $\tilde{x} \in \text{Int}\left(\cup_{g \in \tilde{G}'} g \overline{F'}\right)$, thus thanks to the maximum principle for parabolic equations applied in ω , we can see that $\tilde{h}(x, \tilde{t}) \equiv \mu$, for $x \in B_\delta(\tilde{x}) \cap \overline{F'}$ (where $\delta > 0$). To finish the proof, we are going to show that the set $S := \{y \in \overline{F'} : \tilde{h}(y, \tilde{t}) = \mu\}$ is relatively open. Indeed, let $y \in S$ and let w be the projection of $e^{-\lambda \tilde{t}} u(y, \tilde{t})$ on $\overline{\Phi}$. We have $e^{-\lambda \tilde{t}} u(y, \tilde{t}) - w = \mu \rho$, and repeating the above argument we find $\tilde{h}(x, \tilde{t}) \equiv \mu$, for $x \in B_{\delta'}(y) \cap \overline{F'}$ (where $\delta' > 0$). Thus, by connectedness $\tilde{h}(\cdot, \tilde{t}) \equiv \mu > 0$ on $\overline{F'}$ which is a contradiction since $\tilde{h}(0, \tilde{t}) = 0$.

Let us now give the proof of the Lemma for arbitrary groups. We just present it when G is a finite reflection group since it is similar for discrete reflection groups. We will need to apply the maximum principle to some projections of the solution u . We denote by ρ_1, \dots, ρ_k the outward unit normals to the hyperplanes Π_1, \dots, Π_k that bound the fundamental domain Φ (see Definition 1). We also consider the collection Π_1, \dots, Π_q ($k \leq q$) of all the reflection hyperplanes of Γ , and

choose for $j > k$, a unit normal ρ_j to Π_j . Since the potential W is symmetric, we know that for every $j = 1, \dots, q$, the projection $h_j(x, t) := \langle u(x, t), \rho_j \rangle$ satisfies the linear scalar equation:

$$\Delta h_j + c_j h_j - \frac{\partial h_j}{\partial t} = 0 \text{ in } B_R \times (0, +\infty),$$

with

$$(4.7) \quad c_j = - \left\langle \int_0^1 W_{uu}(u + (s-1)\langle u, \rho_j \rangle \rho_j) \rho_j \, ds, \rho_j \right\rangle.$$

From this formula, one can see that c_j is continuous and bounded on $B_R \times [0, T]$, for every $T > 0$.

Now, suppose that for some $t_0 > 0$, $u(\cdot, t_0)$ does not belong to the class A^R .

In order to have an equation with a nonpositive coefficient, we again do the standard transformation and set $\tilde{h}_j(x, t) := e^{-\lambda t} h_j(x, t)$, where the constant λ is chosen bigger than $\sup\{c_j(x, t) : x \in B_R, t \in [0, t_0], j = 1, \dots, q\}$. Then, we have

$$\Delta \tilde{h}_j + \tilde{c}_j \tilde{h}_j - \frac{\partial \tilde{h}_j}{\partial t} = 0 \text{ in } B_R \times (0, t_0], \text{ with } \tilde{c}_j = c_j - \lambda \leq 0.$$

Let $\mu := \max\{d(e^{-\lambda t} u(x, t), \bar{\Phi}) \mid x \in \overline{F_R}, t \in [0, t_0]\} > 0$, and suppose that this is achieved at $\tilde{x} \in \overline{F_R}$ at time $\tilde{t} \in (0, t_0]$ (since $u_0 \in A_R$). Define

$$\tilde{u} := e^{-\lambda \tilde{t}} u(\tilde{x}, \tilde{t}), \quad \rho := \frac{\tilde{u} - \tilde{v}}{|\tilde{u} - \tilde{v}|},$$

where \tilde{v} is the unique point of $\partial\Phi$ (since Φ is convex) such that $d(\tilde{u}, \tilde{v}) = \mu$. We will apply the maximum principle to $\tilde{h}(x, t) := \langle e^{-\lambda t} u(x, t), \rho \rangle$ in a neighborhood of (\tilde{x}, \tilde{t}) in $\overline{B_R} \times (0, t_0]$. To do this, in analogy to what was done in the special case considered above we need to consider various possibilities for the unit vector ρ . If \tilde{v} belongs to the interior of a $m - p$ dimensional face $\Pi_1 \cap \dots \cap \Pi_p \cap \bar{\Phi}$ ($1 \leq p \leq k$) of Φ , then, using also that ρ_1, \dots, ρ_k are linearly independent, we have

$$\rho \perp E, \quad E := \Pi_1 \cap \dots \cap \Pi_p, \text{ that is, } \rho \in E^\perp = \mathbb{R}\rho_1 \oplus \dots \oplus \mathbb{R}\rho_p$$

where $\mathbb{R}\rho_j = \{x : x = t\rho_j, t \in \mathbb{R}\}$ and E^\perp is the orthogonal complement of E . Let $\tilde{\Gamma}$ be the subgroup of Γ generated by the reflections with respect to the hyperplanes Π_1, \dots, Π_p . The elements of $\tilde{\Gamma}$ leave invariant the subspace E , and actually $\tilde{\Gamma}$ acts in E^\perp . Let $N \supset \{\pm\rho_1, \dots, \pm\rho_p\}$, be the set of all the unit normals to the reflection hyperplanes of $\tilde{\Gamma}$. We claim that

$$(4.8) \quad \begin{cases} \rho = \alpha_1 \nu_1 + \dots + \alpha_{\tilde{p}} \nu_{\tilde{p}}, & \text{with } 1 \leq \tilde{p} \leq p, \alpha_1, \dots, \alpha_{\tilde{p}} > 0, \nu_1, \dots, \nu_{\tilde{p}} \in N, \\ \text{and } \langle \nu_i, \nu_j \rangle \geq 0, & \text{for } 1 \leq i, j \leq \tilde{p}. \end{cases}$$

Given p linearly independent vectors $\nu_1, \dots, \nu_p \in N$ we denote by $C(\nu_1, \dots, \nu_p)$ the cone

$$(4.9) \quad C(\nu_1, \dots, \nu_p) := \{\alpha_1 \nu_1 + \dots + \alpha_p \nu_p \mid \alpha_1, \dots, \alpha_p \geq 0\}.$$

To prove the claim we start by observing that (since ρ_1, \dots, ρ_p are linearly independent) we have $\rho \in \mathcal{C}$ with

$$\mathcal{C} = C(\rho'_1, \dots, \rho'_p), \quad \text{with } \rho'_j = \rho_j \text{ or } -\rho_j, j = 1, \dots, p.$$

To conclude the proof we show that \mathcal{C} can be partitioned in cones of the form (4.9) that satisfy the condition

$$(4.10) \quad C(\nu_1, \dots, \nu_p) \cap N = \{\nu_1, \dots, \nu_p\}.$$

Note that (4.10) and the fact that $\tilde{\Gamma}$ is a reflection group automatically imply

$$\langle \nu_i, \nu_j \rangle \geq 0, \quad \text{for } 1 \leq i, j \leq p.$$

Indeed N is invariant under $\tilde{\Gamma}$ (N is the root system of $\tilde{\Gamma}$) and therefore $\langle \nu_i, \nu_j \rangle < 0$ implies that the reflection $\nu \notin \{\nu_i, \nu_j\}$ of ν_j in the hyperplane orthogonal to ν_i belongs to both N and $C(\nu_1, \dots, \nu_p)$ in contradiction with (4.10). If \mathcal{C} does not satisfy (4.10) there exists $\nu \in N$ that (possibly after a renumbering of the vectors ρ'_1, \dots, ρ'_p) has the form $\nu = \alpha_1 \rho'_1 + \dots + \alpha_{\hat{p}} \rho'_{\hat{p}}$ with $2 \leq \hat{p} \leq p$, $\alpha_1, \dots, \alpha_{\hat{p}} > 0$ and we can partition \mathcal{C} into the \hat{p} cones $\mathcal{C}_i = C(\rho'_1, \dots, \rho'_{i-1}, \nu, \rho'_{i+1}, \dots, \rho'_p)$, $i = 1, \dots, \hat{p}$ defined by the linearly independent vectors $\rho'_1, \dots, \rho'_{i-1}, \nu, \rho'_{i+1}, \dots, \rho'_p$, $i = 1, \dots, \hat{p}$. If \mathcal{C}_i does not satisfy (4.10) we partition \mathcal{C}_i in the same fashion used for \mathcal{C} and continue in this way. Note that at each step (if some of the cones of the partition does not satisfies (4.10)) the number of vectors in N used to generate the cones of the partition increases by one. Therefore, since N is a finite set, the process terminates after a finite number of steps exactly when all the cones of the partition satisfy (4.10). This conclude the proof of the claim.

Since $\tilde{v} \in E$, it follows that (with $\nu_1, \dots, \nu_{\tilde{p}}$ the vectors in (4.8))

$$(\tilde{x}, \tilde{t}) \in \omega := \{(x, t) \in \overline{B_R} \times (0, t_0] \mid \langle e^{-\lambda t} u(x, t), \nu_j \rangle > 0, \forall j = 1, \dots, \tilde{p}\},$$

which is relatively open in $\overline{B_R} \times (0, t_0]$, and in addition we have

$$\Delta \tilde{h} + \tilde{c} \tilde{h} - \frac{\partial \tilde{h}}{\partial t} \geq 0 \text{ in } \omega, \text{ with } \tilde{c} = \max\{\tilde{c}_j \mid j = 1, \dots, q\} \leq 0.$$

At this stage, the fact that Φ has acute angles (i. e. $\langle \rho_i, \rho_j \rangle \leq 0$ for $1 \leq i < j \leq k$) is essential to conclude the proof. This property implies that

$$(4.11) \quad \tilde{u} \in \Pi_j \text{ (with } 1 \leq j \leq k) \Rightarrow \tilde{v}, \rho \in \Pi_j,$$

and as a consequence

$$(4.12) \quad f(s_i) \notin \Gamma_\rho \Rightarrow \tilde{x} \notin P_i,$$

where P_i is, as in Definition 1, a hyperplane bounding F corresponding to the reflection $s_i \in G$. To show (4.12), suppose that $\tilde{x} \in P_i$. Then

$$u(\tilde{x}, \tilde{t}) \in \ker(f(s_i) - I_m) \text{ by } f\text{-equivariance,}$$

and thanks to Hypothesis 1 and (4.11),

$$\rho \in \ker(f(s_i) - I_m) \Rightarrow f(s_i) \in \Gamma_\rho.$$

Property (4.12) will enable us to locate \tilde{x} in $\overline{F_R}$. Let \tilde{G} be the subgroup of G generated by the reflections s_i ($i = 1, \dots, l$) such that $f(s_i) \in \Gamma_\rho$. By f -equivariance we have

$$\mu = \tilde{h}(\tilde{x}, \tilde{t}) = \max\{\tilde{h}(x, t) \mid x \in \cup_{g \in \tilde{G}} g \overline{F_R}, t \in [0, t_0]\}.$$

But now, either $\tilde{x} \in \text{Int}\left(\cup_{g \in \tilde{G}} g \overline{F_R}\right)$ or $\tilde{x} \in \text{Int}\left(\cup_{g \in \tilde{G}} g \overline{F}\right) \cap \partial B_R$. In both cases, we can see, thanks to the maximum principle for parabolic equations applied in ω and thanks to Hopf's Lemma, that $\tilde{h}(x, \tilde{t}) \equiv \mu$, for $x \in B_\delta(\tilde{x}) \cap \overline{F_R}$ (where $\delta > 0$). To finish the proof, we are going to show that the set $S := \{y \in \overline{F_R} \mid \tilde{h}(y, \tilde{t}) = \mu\}$ is relatively open. Indeed, let $y \in S$ and let w be the projection of $e^{-\lambda \tilde{t}} u(y, \tilde{t})$ on $\overline{\Phi}$. We have $e^{-\lambda \tilde{t}} u(y, \tilde{t}) - w = \mu \rho$ and according to the preceding argument $\tilde{h}(x, \tilde{t}) \equiv \mu$ for $x \in B_{\delta'}(y) \cap \overline{F_R}$ (where $\delta' > 0$). Thus, by connectedness $\tilde{h}(\cdot, \tilde{t}) \equiv \mu > 0$ on $\overline{F_R}$ which is a contradiction since $\tilde{h}(0, \tilde{t}) = 0$. \square

If the initial condition in (4.2) (respectively (4.3)) is a $W^{1,2}(B_R, \mathbb{R}^m)$ (respectively, $W_{\text{loc}}^{1,2}(\mathbb{R}^n, \mathbb{R}^m)$), bounded map, then the solution to (4.2) (respectively, (4.3)) belongs to $C([0, \infty), W^{1,2}(B_R, \mathbb{R}^m))$ (respectively $C([0, \infty), W_{\text{loc}}^{1,2}(\mathbb{R}^n, \mathbb{R}^m))$), and is smooth for $t > 0$. We are now going to take the minimizer u_R constructed in Step 1, as the initial condition in (4.2) (respectively (4.3)). Thanks to Lemma 4.3 below (and to its analog for discrete reflection groups), we can construct a sequence of smooth, f -equivariant, and positive maps (u_k) which converges to u_R for the $W^{1,2}$ norm, as $k \rightarrow \infty$. Applying then Lemmas 4.1 and 4.2 to u_k , and utilizing the continuous dependence for the flow of the initial condition, we obtain that the solution to (4.2) (respectively (4.3)), with initial condition u_R , is f -equivariant and positive, that is, $u(\cdot, t; u_R) \in A^R$, for $t \geq 0$ (respectively $u(\cdot, t; u_R) \in A$, for $t \geq 0$).

Lemma 4.3. *Let $u \in W^{1,2}(B_R, \mathbb{R}^m) \cap L^\infty(B_R, \mathbb{R}^m)$ be a f -equivariant map such that $u(\overline{F_R}) \subset \overline{\Phi}$. Then, there exists a sequence $(u_k) \subset C(\overline{B_R}, \mathbb{R}^m)$ of globally Lipschitz maps with the following properties:*

- (i) u_k is f -equivariant,
- (ii) $\|u_k\|_{L^\infty(B_R, \mathbb{R}^m)} \leq \|u\|_{L^\infty(B_R, \mathbb{R}^m)}$,
- (iii) $u_k(\overline{F_R}) \subset \overline{\Phi}$ (positivity),
- (iv) u_k converges to u in $W^{1,2}(B_R, \mathbb{R}^m)$, as $k \rightarrow \infty$.

Proof. See Proposition 5.2 in [3]. □

Thanks to the fact that u_R is a global minimizer of J_{B_R} (respectively J_F^R) in A^R (respectively A), and since $u(\cdot, t; u_R)$ is a classical solution to (4.2) (respectively (4.3)) for $t > 0$, we conclude from the computation

$$(4.13) \quad \frac{d}{dt} J_{B_R}(u(\cdot, t)) = - \int_{B_R} |u_t|^2 dx \quad \left(\text{respectively } \frac{d}{dt} J_F^R(u(\cdot, t)) = - \int_F |u_t|^2 dx \right)$$

that $|u_t(x, t)| = 0$, for all $x \in B_R$ (respectively $x \in F$) and $t > 0$. Hence, for $t > 0$, $u(\cdot, t)$ satisfies

$$(4.14) \quad \Delta u(x, t) - W_u(u(x, t)) = 0 \quad (\text{respectively } \Delta u(x, t) - R^2 W_u(u(x, t)) = 0).$$

By taking $t \rightarrow 0+$ and utilizing the continuity of the flow in $W^{1,2}(B_R, \mathbb{R}^m)$ (respectively $W_{\text{loc}}^{1,2}(\mathbb{R}^n, \mathbb{R}^m)$) at $t = 0$, we obtain that u_R is a f -equivariant, classical solution to system (1.1) (respectively (3.5)) satisfying also $u_R(\overline{F_R}) \subset \overline{\Phi}$ (respectively $u_R(\overline{F}) \subset \overline{\Phi}$).

If G is a finite reflection group, from the family of solutions $u_R \in C^3(B_R, \mathbb{R}^m)$, $R \geq 1$ we can deduce the existence of an entire f -equivariant classical solution $u \in C^3(\mathbb{R}^n, \mathbb{R}^m)$ to system (1.1) defined by

$$(4.15) \quad u(x) = \lim_{j \rightarrow +\infty} u_{R_j}(x),$$

where $R_j \rightarrow +\infty$ is a suitable subsequence and the convergence is in the C^2 sense in compact subsets of \mathbb{R}^n . This follows from the fact that u_R satisfies the bound

$$(4.16) \quad \|u_R\|_{C^{2+\alpha}(\overline{B_R}, \mathbb{R}^m)} \leq M',$$

for some $\alpha \in (0, 1)$ and $M' > 0$ independent of $R \geq 1$. The estimate (4.16) follows by elliptic regularity from (4.1), from the assumed smoothness of W and from the fact that ∂B_R is uniformly smooth for $R \geq 1$. The solution u satisfies also: $u(\overline{F}) \subset \overline{\Phi}$.

Step 3 (Pointwise estimates).

From Step.2 to complete the proof of Theorem 3.1 and Theorem 3.2 it remains to prove that the entire solution u to system (1.1) and the solution u_R to system (3.5) satisfy the pointwise estimates stated in Theorem 3.1 and Theorem 3.2 respectively. For establishing these estimates we utilize the following theorem that we quote from [12]. We need the concept of *local minimizer*

Definition 2. Let $\Omega \subset \mathbb{R}^n$ an open set. A map $u \in C^2(\Omega; \mathbb{R}^m) \cap L^\infty(\Omega, \mathbb{R}^m)$ is a local minimizer if

$$(4.17) \quad J_{\Omega'}(u) \leq J_{\Omega'}(u + v),$$

for all $v \in W_0^{1,2}(\Omega', \mathbb{R}^m) \cap L^\infty(\Omega', \mathbb{R}^m)$ and for all smooth bounded open subset $\Omega' \subset \Omega$.

Theorem 4.4. Assume $W : \mathbb{R}^m \rightarrow \mathbb{R}$ is a nonnegative C^3 potential and that $W(a) = 0$ for some $a \in \mathbb{R}^m$ satisfying condition (ii) in Hypothesis 2. Let $Z \subset \mathbb{R}^m$ be the set $Z = \{z \neq a : W(z) = 0\}$ and let $Z_\delta = \cup_{z \in Z} B_{z,\delta}$ a δ -neighborhood of Z .

Assume that $u : \Omega \rightarrow \mathbb{R}^m$ is a local minimizer and that there is $M > 0$ such that

$$(4.18) \quad |u| + |\nabla u| \leq M, \quad x \in \Omega$$

Then there is a $\bar{q} \in (0, q^*]$ and a strictly decreasing continuous function $r : (0, \bar{q}] \rightarrow (0, \infty)$, $\lim_{q \rightarrow 0^+} r(q) = +\infty$, such that the condition

$$(4.19) \quad B_{r(q)}(x_0) \subset \Omega \quad \text{and} \quad u(B_{r(q)}(x_0)) \cap Z_\delta = \emptyset,$$

implies

$$|u(x_0) - a| < q.$$

The map r depends only on W and M if $Z = \emptyset$ and also on δ otherwise.

Moreover, if a is nondegenerate in the sense that $D^2W(a)$ is positive definite, then there exists a constant $k_0 > 0$ such that

$$(4.20) \quad r(q) \leq r(\bar{q}) + \frac{1}{k_0} \left| \log \frac{q}{\bar{q}} \right|, \quad q \in (0, \bar{q}].$$

Remark 4.5. The map r has a strictly decreasing inverse $q : [r(\bar{q}), +\infty) \rightarrow (0, \bar{q}]$. If a is nondegenerate (4.20) implies

$$(4.21) \quad q(r) < K_0 e^{-k_0 r}, \quad r \in [r(\bar{q}), +\infty),$$

for some $K_0 > 0$.

For the proof of Theorem 4.4 we refer to [12], [11]. In [12] the proof is for the case of a generic potential and covers all cases where $D_0 = F$. To treat the general case $F \subset D_0$ one needs to show that Theorem 4.4 holds true in the case of symmetric potentials and f -equivariant local minimizers. In [11] the validity of Theorem 4.4 is established for symmetric potential and $f = I$. The arguments in [11] extend naturally to cover the general case of f -equivariance.

If G is a finite reflection group we apply Theorem 4.4 to u_R with $\Omega = D_{0,R} := D_0 \cap B_R$ and $Z = \{\gamma a\}_{\gamma \in \Gamma} \setminus \{a\}$. From Step.2 we have $u_R(\bar{F}_R) \subset \bar{\Phi}$ and therefore by f -equivariance

$$(4.22) \quad u_R(\bar{D}_{0,R}) \subset u_R(\bar{D}_R) \subset \cup_{\gamma \in \Gamma_a} \gamma \bar{\Phi} = \bar{\mathcal{D}}.$$

Since by Hypothesis 3, as we have seen, a is the unique minimizer of W in \mathcal{D} it results

$$(4.23) \quad u_R(D_{0,R}) \cap Z_\delta = \emptyset, \quad \text{for } \delta = d(a, \partial \mathcal{D}) > 0.$$

This, the bound (4.16) and Theorem 4.4 imply

$$(4.24) \quad |u_R(x) - a| < q(d(x, \partial D_{0,R})), \quad x \in D_{0,R},$$

for some positive function $r \rightarrow q(r)$ that does not depend on R . Therefore the inequality (4.24) passes to the limit along the sequence u_{R_j} that defines the entire solution u . This and the f -equivariance of u establish (ii). The exponential estimate (iii) follows from Remark 4.5. The proof of Theorem 3.1 is complete.

Assume now that G is a discrete group. In this case the fundamental domain can be bounded as for the group G' considered in Section 2 or unbounded as for the group G generated by the reflections in the plane $\{x_1 = 0\}$ and $\{x_1 = 1\}$ of \mathbb{R}^n , $n > 1$ where $F = \{x \in \mathbb{R}^n : 0 < x_1 < 1\}$. For establishing the estimates (ii) and (iii) in Theorem 3.2 it suffices to consider the case where F is bounded. For $R > 0$ define

$$(4.25) \quad \begin{aligned} v_R(y) &= u_R\left(\frac{y}{R}\right), \quad y \in \mathbb{R}^n, \\ F^R &= \{y \in \mathbb{R}^n : \frac{y}{R} \in F\}, \end{aligned}$$

and let G^R the discrete reflection group generated by the reflections in the planes P_1^R, \dots, P_l^R that bound F^R . There is an obvious group isomorphism $\eta^R : G^R \rightarrow G$ between G^R and G and the minimality of u_R implies that $v_R \in W_{\text{loc}}^{1,2}(\mathbb{R}^n, \mathbb{R}^m)$ is a local minimizers in the class of f^R -equivariant maps where $f^R := f \circ \eta^R$. Therefore v_R is a solution of

$$(4.26) \quad \Delta v - W_u(v) = 0, \quad \text{in } \mathbb{R}^n.$$

This, (4.1) and elliptic regularity implies

$$(4.27) \quad |\nabla v_R| \leq M', \quad \text{in } \mathbb{R}^n,$$

for some $M' > 0$ independent of R . As before we have

$$u_R(\overline{D}_0) \subset \cup_{\gamma \in \Gamma_a} \gamma \overline{\Phi} = \overline{\mathcal{D}},$$

or equivalently

$$(4.28) \quad v_R(\overline{D}_0^R) \subset \overline{\mathcal{D}},$$

where $\overline{D}_0^R = \cup_{g \in G^a} (\eta^R)^{-1}(g) \overline{F}^R$. From (4.28) it follows

$$(4.29) \quad v_R(D_0^R) \cap Z_\delta = \emptyset, \quad \text{for } \delta = d(a, \partial \mathcal{D}) > 0.$$

Therefore we can apply Theorem 4.4 to v_R with $\Omega = D_0^R$ and deduce, for $R \geq R_0 := r(\bar{q})$

$$(4.30) \quad |v_R(y) - a| \leq q(d(y, \partial D_0^R)), \quad y \in D_0^R,$$

which is equivalent to

$$(4.31) \quad |u_R(x) - a| \leq q(Rd(x, \partial D_0)), \quad x \in D_0.$$

The rest of the proof is as in the case of G finite discussed before. The proof of Theorem 3.2 is complete.

5. THREE DETAILED EXAMPLES INVOLVING THE REFLECTION GROUP OF THE TETRAHEDRON

5.1. Preliminaries. In this subsection we recall some properties of the group of symmetry of a regular tetrahedron and the group of symmetry of a cube.

Let \mathcal{T} be the group of symmetry of a regular tetrahedron $A_1 A_2 A_3 A_4$ (with $A_1 = (1, 1, 1)$, $A_2 = (-1, -1, 1)$, $A_3 = (1, -1, -1)$, $A_4 = (-1, 1, -1)$) which can be inscribed in a cube centered at the origin O (see Figure 4). The order of \mathcal{T} is $|\mathcal{T}| = 24$ and \mathcal{T} is isomorphic to the permutation group S_4 . The 24 elements of \mathcal{T} are associated to the following elements of S_4 :

- I_3 the identity map of \mathbb{R}^3 , corresponds to the unit of S_4 (I_3 fixes the 4 vertices),

- the 8 rotations of angle $\pm 2\pi/3$ with respect to the axes OA_i correspond to the 3-cycles $(j\ k\ l)$ (only the vertex A_i is fixed),
- the 6 reflections with respect to the planes OA_iA_j correspond to the transpositions $(k\ l)$ (the vertex A_i and A_j are fixed),
- the 3 symmetries with respect to the coordinate axes correspond to the permutations $(i\ j)(k\ l)$ (no vertex fixed),
- a reflection with respect to the plane OA_kA_l composed with a rotation with respect to the axis OA_i corresponds to one of the six 4-cycles $(i\ j\ k\ l)$ (no vertex fixed).

Since there exists a homomorphism between the permutation groups S_4 and S_3 , and since S_3 is isomorphic to the dihedral group D_3 which is the group of symmetry of an equilateral triangle in the plane, we also have a homomorphism $\phi : \mathcal{T} \rightarrow D_3$. ϕ associates:

- I_3 and the symmetries with respect to the coordinate axes to I_2 (i.e. $\ker \phi = \{I_3 \text{ and the three symmetries with respect to the coordinate axes}\}$),
- the rotation of angle $2\pi/3$ with respect to the axis OA_1 to the rotation ρ of angle $2\pi/3$ in the plane,
- the reflections with respect to the planes OA_1A_2 and OA_3A_4 to the reflection $\sigma_0 : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ in the u_1 axis (cf. Figure 4),
- the reflections with respect to the planes OA_1A_4 and OA_2A_3 to the reflection $\sigma_1 = \rho\sigma_0$,
- the reflections with respect to the planes OA_1A_3 and OA_2A_4 to the reflection $\sigma_2 = \rho^2\sigma_0$.

Let \mathcal{K} be the group of symmetry of a cube centered at the origin O with vertices at the points $(\pm 1, \pm 1, \pm 1)$. The order of \mathcal{K} is $|\mathcal{K}| = 48$, and \mathcal{K} contains the group of symmetry of the regular tetrahedron as a subgroup (i.e. $\mathcal{T} < \mathcal{K}$). Let $\sigma : \mathbb{R}^3 \rightarrow \mathbb{R}^3$, $\sigma(x) = -x$, be the antipodal map. Clearly, σ is an element of \mathcal{K} of order 2 which does not belong to \mathcal{T} , and $\mathcal{K} = \mathcal{T} \cup \sigma\mathcal{T}$. Furthermore, σ commutes with the reflections with respect to the planes OA_iA_j , thus σ commutes with every element of \mathcal{K} . As a consequence, the correspondence

$$\{I_3, \sigma\} \times \mathcal{T} \ni (\alpha, \beta) \rightarrow \alpha\beta \in \mathcal{K},$$

defines an isomorphism of the group product $\{I_3, \sigma\} \times \mathcal{T}$ onto \mathcal{K} , and we can define a homomorphism $\psi : \mathcal{K} \rightarrow \mathcal{T}$ by setting $\psi(\beta) = \beta$, and $\psi(\sigma\beta) = \beta$, for every $\beta \in \mathcal{T}$. By definition, ψ leaves invariant the elements of \mathcal{T} . We also mention that \mathcal{K} contains the 3 reflections with respect to the coordinate planes $x_i = 0$ (which are the symmetries with respect to the coordinate axes Ox_i composed with σ).

5.2. A solution $u : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ to (1.1) with the reflection group of the tetrahedron acting on the domain and the reflection group of the equilateral triangle acting on the target.

In this example we consider the aforementioned homomorphism $\phi : \mathcal{T} \rightarrow D_3$. Let F be the fundamental domain of \mathcal{T} bounded by the planes OA_1A_2 , OA_3A_4 and OA_1A_4 , and Φ be the fundamental domain of D_3 bounded by the lines $u_2 = 0$ and $u_2 = \sqrt{3}u_1$ corresponding to the reflections σ_0 and σ_1 . According to what precedes, the image by ϕ of the reflections with respect to the planes OA_1A_2 and OA_3A_4 is σ_0 , while the image by ϕ of the reflection with respect to the plane OA_3A_4 is σ_1 . Thus ϕ is a positive homomorphism that associates F to Φ , and Hypothesis 1 is satisfied. If Hypotheses 2–3 also hold, Theorem 3.1 ensures the existence of a ϕ -equivariant solution u to (1.1). In particular (see Figure 4) ϕ -equivariance implies that u maps

- the coordinate axes into the reflection lines (with the same colour),
- the planes OA_1A_2 and OA_3A_4 into the reflection line $u_2 = 0$,
- the planes OA_1A_4 and OA_2A_3 into the reflection line $u_2 = \sqrt{3}u_1$,

- the planes OA_1A_3 and OA_2A_4 into the reflection line $u_2 = -\sqrt{3}u_1$,
- the diagonals of the cube at the origin O .

In addition Theorem 3.1 implies that the solution u is positive (i.e. $u(\overline{F}) \subset \overline{\Phi}$), and therefore u maps each fundamental domain of \mathcal{T} into a fundamental domain of D_3 as in Figure 4. If for instance the potential W has 6 minima (one in the interior of each fundamental domain of D_3), then the domain space \mathbb{R}^3 is also split into six regions as in Figure 4. Properties (ii) and (iii) of Theorem 3.1 state that for every x in such a region D , $u(x)$ converges as $d(x, \partial D) \rightarrow \infty$, to the corresponding minimum a of W .




FIGURE 4. Fundamental domains for the action of \mathcal{T} on \mathbb{R}^3 (left) and for the action of D_3 on \mathbb{R}^2 (right). The ϕ -equivariant solution $u : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ of (1.1) given by Theorem 3.1 maps fundamental domains into fundamental domains with the same color. In particular u maps the infinite double cone (union of four fundamental domains) generated by O and by the two yellow triangle into the sector Φ .

5.3. A solution $u : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ to (1.1) with the reflection group of the cube acting on the domain and the reflection group of the tetrahedron acting on the target.

In this example we consider the homomorphism $\psi : \mathcal{K} \rightarrow \mathcal{T}$ defined previously. Now, we denote by F be the fundamental domain of \mathcal{K} bounded by the planes OA_1A_2 , OA_1A_4 and $x_2 = 0$, and by Φ be the fundamental domain of \mathcal{T} bounded by the planes OA_1A_2 , OA_1A_4 and OA_2A_3 (cf. Figure 5). According to what precedes, ψ leaves invariant the reflections with respect to the planes OA_1A_2 and OA_1A_4 , while the image by ψ of the reflection with respect to the plane $x_2 = 0$ is the coordinate axis x_2 , that is, the intersection of the planes OA_1A_4 and OA_2A_3 . Thus ψ is a positive homomorphism that associates F to Φ , and Hypothesis 1 is satisfied. If Hypotheses 2–3 also hold, Theorem 3.1 ensures the existence of a ψ -equivariant solution u to (1.1). More precisely, ψ -equivariance implies that u maps

- every plane OA_iA_j into itself,
- every diagonal of the cube into itself,
- the coordinate planes into the perpendicular coordinate axes,

- the coordinate axes at the origin O .

In addition, the solution u is positive (i.e. $u(\overline{F}) \subset \overline{\Phi}$), and in fact u maps each fundamental domain of \mathcal{K} into a fundamental domain of \mathcal{T} as in Figure 5.




FIGURE 5. Fundamental domains for the action on \mathbb{R}^3 of \mathcal{K} (left) and \mathcal{T} (right). The ψ -equivariant solution $u : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ of (1.1) given by Theorem 3.1 maps fundamental domains into fundamental domains with the same color. Note in particular that u maps $\overline{F} \cup \sigma \overline{F}$ into $\overline{\Phi}$.

If the potential W has for instance 4 minima (located at the vertices of the tetrahedron A_1, A_2, A_3 and A_4), then the stabilizer Γ_a of $a = A_1$ in $\Gamma = \mathcal{T}$ has six elements: I_3 , the reflections with respect to the planes OA_1A_2 , OA_1A_3 and OA_1A_4 , and the rotations of angle $\pm 2\pi/3$ with respect to the axis OA_1 . Thus \mathcal{D} is the (interior of the closure) of the union of the six fundamental domains that have A_1 on their boundary; the group $\psi^{-1}(\Gamma_a) = \Gamma_a \cup \sigma \Gamma_a$ has 12 elements, and the set D has two connected components: the solid right angle $D_0 = \{x_i > 0, \forall i = 1, 2, 3\}$ and $\sigma D_0 = -D_0$ (cf. Figure 6 and also note that the group G^a is in this particular case the group Γ_a). According to Theorem 3.1, if $x \in D_0$ and $d(x, \partial D_0) \rightarrow \infty$ (that is, if $x_i \rightarrow +\infty$ for every $i = 1, 2, 3$) then $u(x) \rightarrow a$. Of course, the same result is true when $x \in -D_0$ and $d(x, \partial(-D_0)) \rightarrow \infty$, and the solution also converges in the other solid right angle cones to the corresponding minima of W as in Figure 6.

5.4. A crystalline structure in \mathbb{R}^3 .

Now, let us consider the discrete reflection group \mathcal{K}' acting in \mathbb{R}^3 which is generated by the reflections s_1, s_2, s_3 and s_4 with respect to the corresponding planes $P_1 := OA_1A_2$, $P_2 := OA_1A_4$, $P_3 := \{x_2 = 0\}$ and $P_4 := \{x_1 + x_3 = 2\}$. These planes bound the fundamental domain F' of \mathcal{K}' with vertices at the points $O, A_1, I := (1, 0, 1)$ and $B := (0, 0, 2)$ (cf. Figure 7). The point group of \mathcal{K}' , that is the stabilizer of the origin, is the group \mathcal{K} , and we have $\mathcal{K}' = T\mathcal{K}$, where T is the




FIGURE 6. The sets $\mathcal{D} = \{u = \alpha_1(1, 1, -1) + \alpha_2(-1, 1, 1) + \alpha_3(1, -1, 1), \alpha_i > 0, i = 1, 2, 3\}$, $D_0 = \{x_1 > 0, \forall i = 1, 2, 3\}$ and $D = D_0 \cup \sigma D_0$ when W has four minima on the vertices of the tetrahedron. In this case the solution u of (1.1) given by Theorem 3.1 satisfies: $u(x) \rightarrow A_1$, for $\min_i |x_i| \rightarrow +\infty, x \in D$.

translation group of \mathcal{K}' . T is generated by the translations with respect to the vectors $t_1 := (2, 0, 2)$, $t_2 := (0, 2, 2)$ and $t_3 := (0, -2, 2)$. By composing the canonical homomorphism $p : \mathcal{K}' \rightarrow \mathcal{K}$ such that $p(tg) = g$ for every $t \in T$ and $g \in \mathcal{K}$, with the homomorphism $\psi : \mathcal{K} \rightarrow \mathcal{T}$ defined previously, we obtain a homomorphism $\psi' : \mathcal{K}' \rightarrow \mathcal{T}$. We note that $\psi'(s_4)$ is the image by ψ of the reflection with respect to the plane OA_2A_3 , which is the reflection with respect to OA_2A_3 . Thus, ψ' is a positive homomorphism which associates F' to the fundamental domain Φ of \mathcal{T} bounded by the planes OA_1A_2 , OA_1A_4 and OA_2A_3 . In this case the elementary crystal $C = \cup_{g \in \mathcal{K}} g\overline{F'}$ is a *rhombic dodecahedron* (cf. Figure 7) that tiles the three dimensional space when translated by the elements of T ². Several structures are possible for the solution u_R given by Theorem 3.2 depending on the position of $a \in \overline{\Phi}$. For instance if $a \in \Phi$ we have $\mathcal{D} = \Phi$ and $D_0 = F'$. If $a = (0, 1, 0)$ (cf. Figure 8) then $\mathcal{D} = \{u : \max\{|u_1|, |u_3|\} < u_2\}$ and D_0 is the pyramid with basis the rhombus defined by the points $A_1, B, (1, -1, 1), (2, 0, 0)$, and vertex in O . Finally, if $a = A_1$, \mathcal{D} is the cone which has vertex in O and is generated by the triangle with vertices at the points $(1, 1, -1), (1, -1, 1), (-1, 1, 1)$, while D_0 is the polyhedron (union of two pyramids) with vertices at the points $O, (2, 0, 0), (0, 2, 2), B, A_1$ and $D = \cup_{t \in T} (D_0 \cup \sigma D_0)$.

6. APPENDIX: OTHER EXAMPLES IN LOWER DIMENSION

6.1. Positive homomorphisms between finite reflection groups of the plane.

²Space filling tessellation with rhombic dodecahedra is the crystal structure in which often are found garnets and other minerals like pyrite and magnetite.




FIGURE 7. Fundamental domains for the action on \mathbb{R}^3 of \mathcal{K}' (left) and \mathcal{T} (right). The fundamental domain F' of \mathcal{K}' is a pyramid with basis the triangle A_1BI and vertex in O . Under the action of the point group \mathcal{K} , F' generates the rhombic dodecahedron C (left) which tiles the domain space \mathbb{R}^3 when translated by the elements of T . The ψ' -equivariant solution $u : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ of (1.1) given by Theorem 3.2 maps fundamental domains into fundamental domains with the same color. Note in particular that u maps $\cup_{t \in T} (\overline{F'} \cup \sigma \overline{F'})$ into $\overline{\Phi}$.

The finite reflection groups of the plane are the dihedral groups D_n with $n \geq 1$. D_n contains $2n$ elements: the rotations $r_n^0 = I_2, r_n^1, \dots, r_n^{n-1}$ (where I_2 is the identity map of the plane, and r_n is the rotation of angle $2\pi/n$), and the reflections $r_n^0 s = s, r_n^1 s, \dots, r_n^{n-1} s$ (where s is the reflection with respect to the x_1 coordinate axis). Similarly, the elements of D_{nk} (with $k \geq 1$) are the rotations: $r_{nk}^0 = I_2, r_{nk}^1, \dots, r_{nk}^{nk-1}$ (where r_{nk} is the rotation of angle $2\pi/nk$) and the reflections $r_{nk}^0 s = s, r_{nk}^1 s, \dots, r_{nk}^{nk-1} s$. In the two Propositions below we have determined all the positive homomorphisms between finite reflection groups of the plane (up to an isomorphism). From the list of homomorphisms between dihedral groups established in [17], we have extracted the positive ones.

For $m = \pm 1$ we define the homomorphism $f_m : D_{nk} \rightarrow D_n$, by setting $f_m(r_{nk}^p) = r_n^{mp}$ and $f_m(r_{nk}^p s) = r_n^{mp} s$, for every integer p . Thanks to the property $sr = r^{-1}s$ (which holds for every reflection s and every rotation r), it is easy to check that f_m is a homomorphism from D_{nk} onto D_n . We can also define the homomorphism $g : D_{2k} \rightarrow D_2$, by setting $g(r_{2k}^p) = s^p$ and $g(r_{2k}^p s) = s^p \sigma$ for every integer p , where σ denotes the antipodal map $\sigma u = -u$.

Proposition 6.1. *If $n \geq 2$, $k \geq 1$, $G = D_{nk}$ acts on the domain plane \mathbb{R}^2 , and $\Gamma = D_n$ on the target plane \mathbb{R}^2 , then for every $m = \pm 1$, f_m is a positive homomorphism which associates the fundamental domain $F := \{re^{it} \mid 0 < r, 0 < t < \pi/nk\}$ of G to the fundamental domain $\Phi := \{re^{it} \mid 0 < r, 0 < mt < \pi/n\}$ of Γ . In addition, the homomorphism f_m leaves invariant the elements of $D_n < D_{nk}$ if and only if $mk = 1 \pmod n$.*




FIGURE 8. The sets D_0 and \mathcal{D} when W has six minima (one in the middle of each side of the tetrahedron). In this case the ψ' -equivariant solution $u_R : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ given by Theorem 3.2 satisfies $\lim_{R \rightarrow +\infty} u_R(x+t) = (0, 1, 0)$ for $x \in D_0 \cup \sigma D_0$, $t \in T$.

Proof. By definition, the lines that bound the fundamental domain F correspond to the reflections s and $r_{nk}s$. Since for $m = \pm 1$, the fixed points of the reflections $f_m(s) = s$ and $f_m(r_{nk}s) = r_n^m s$ are the lines that bound the fundamental domain Φ , the homomorphism f_m can associate F to Φ . f_m leaves invariant the elements of $D_n < D_{nk}$ if and only if $f_m(r_n) = r_n$, that is, if and only if

$$f_m(r_n) = f_m(r_{nk}^k) = r_n^{mk} = r_n \Leftrightarrow r_n^{mk-1} = I_2 \Leftrightarrow mk = 1 \pmod{n}.$$

□

Proposition 6.2. *If $k \geq 1$, $G = D_{2k}$ acts on the domain plane \mathbb{R}^2 , and $\Gamma = D_2$ on the target plane \mathbb{R}^2 , then g is a positive homomorphism which associates the fundamental domain $F = \{re^{it} \mid 0 < r, 0 < t < \pi/2k\}$ of G to the fundamental domain $\Phi = \{re^{it} \mid 0 < r, 0 < t < \pi/2\}$ of Γ .*

Proof. As previously, we see that $g(s) = \sigma$ fixes only the origin, while $g(r_{2k}s) = \sigma s$ fixes the u_2 coordinate axis. Thus, the homomorphism g can associate F to Φ (and in fact it can associate F to any of the 4 fundamental domains of D_2). □

To illustrate the Propositions above, let us give some examples.

- The homomorphism $f : D_6 \rightarrow D_3$ that was mentioned at the beginning of Sec.2 (cf. also [4]) coincides with the homomorphism f_{-1} of Proposition 6.1 with $n = 3$, $k = 2$ and $m = -1$. Since $mk = -2 = 1 \pmod{3}$, we see again that it leaves invariant the elements of D_3 .
- Taking $n = 3$, $k = 5$ and $m = -1$, we check that $mk = 1 \pmod{n}$, and we obtain a new homomorphism $f_{-1} : D_{15} \rightarrow D_3$ that leaves invariant the elements of D_3 . The kernel of this homomorphism is the cyclic group generated by the rotation r_5 .

- Taking $n = 2$, $k = 2$ and $m = 1$, we obtain the positive homomorphism $f_1 : D_4 \rightarrow D_2$. When Hypotheses 2–3 also hold, Theorem 3.1 ensure the existence of a f_1 -equivariant solution to (1.1) which maps \bar{F} into $\bar{\Phi}$, and the other fundamental domains of $G = D_4$ as in Figure 9.
- Considering the homomorphism $g : D_4 \rightarrow D_2$ of Proposition 6.2 (with $k=2$) we can also construct a g -equivariant solution u to (1.1). This solution has the particularity that the coordinate axes are mapped at the origin. Indeed, by g -equivariance, if $x, y \in \mathbb{R}^2$ are symmetric with respect to one of the coordinate axes, then $u(x) = -u(y)$ (cf. Figure 10 for the correspondence of the fundamental domains).




FIGURE 9. The correspondence of the fundamental domains for a solution to (1.1) equivariant with respect to the homomorphism $f_1 : D_4 \rightarrow D_2$.




FIGURE 10. The correspondence of the fundamental domains for a solution to (1.1) equivariant with respect to the homomorphism $g : D_4 \rightarrow D_2$.

6.2. Saddle solutions.

In this subsection we are going to construct as an application of Theorems 3.1 and 3.2, scalar solutions to (1.1) and (3.5) having particular symmetries. The only finite reflection group that acts on the target space \mathbb{R} is the dihedral group $\Gamma = D_1$ with two elements: the identity I_1 and the antipodal map $\sigma u = -u$. We assume that a finite or a discrete reflection group G acts on the domain space \mathbb{R}^n , and that $W : \mathbb{R} \rightarrow \mathbb{R}$ satisfies Hypotheses 2–3, that is,

- W is a nonnegative and even function,
- there exists $M > 0$ such that $W(u) \geq W(M)$, for $u \geq M$,
- $W(u) = 0 \Leftrightarrow u = \pm a$, with in addition $a > 0$ and $W''(a) > 0$.

Clearly, the map ϵ which sends the orientation-preserving motions to I_1 , and the orientation-reversing motions to σ , is a positive homomorphism from G onto Γ . Thus, Theorems 3.1 and 3.2 ensure the existence of classical solutions $u : \mathbb{R}^n \rightarrow \mathbb{R}$ to (1.1) and (3.5) with the following properties:

- (i) ϵ -equivariance implies that if $x, y \in \mathbb{R}^n$ are symmetric with respect to a reflection plane of G , then $u(x) = -u(y)$. In particular, u vanishes on the reflection planes of G . If G is a discrete reflection group, then u is periodic in the sense that $u(x+t) = u(x)$, for every $x \in \mathbb{R}^n$, and every translation t in the translation group $T < G$.
- (ii) Positivity means that either $u \geq 0$ or $u \leq 0$ in each fundamental domain F of G .
- (iii) In each fundamental domain F , $u(x)$ approaches either a or $-a$, as $x \in F$ and $d(x, \partial F)$ increases.

We also give another example, more elaborated, when $G = D_{2k}$ (with $k \geq 1$) acts on \mathbb{R}^2 . Let us consider the homomorphism $h : D_{2k} \rightarrow D_1$ such that $h(r_{2k}^p) = \sigma^p$ and $h(r_{2k}^p s) = \sigma^p$, for every integer p (see the previous subsection for the notation). In this particular set-up, we can again construct a h -equivariant solution $u : \mathbb{R}^2 \rightarrow \mathbb{R}$ to (1.1) which has in each fundamental domain of D_{2k} alternatively even and odd symmetries. Figure 11 represents the symmetries of such a solution for $k = 2$.




FIGURE 11. The symmetries of a solution $u : \mathbb{R}^2 \rightarrow \mathbb{R}$ to (1.1) equivariant with respect to the homomorphism $h : D_4 \rightarrow D_1$.

6.3. Other examples involving discrete reflection groups.

To finish, we give some more examples illustrating Theorem 3.2. Let us assume again that the discrete reflection group G' acts on the domain x -plane as in Sec.2 (cf. also the end of Sec.3), but let us consider now a new reflection group acting on the target u -plane: the group $\Gamma = D_2$. We construct a homomorphism $f'' : G' \rightarrow D_2$ by composing the canonical projection $p : G' \rightarrow D_6$ with the homomorphism $g : D_6 \rightarrow D_2$ defined in subsection 6.1 (that is, $f'' = g \circ p$). As we did before for the homomorphism f' , we can check that f'' is a positive homomorphism. Thus, once again, Theorem 3.2 allows us to construct f'' -equivariant solutions u_R to (3.5). Figure 12 represents the correspondence of the fundamental domains of G' with the fundamental domains of D_2 for such solutions (compare with Figure 3). The f'' -equivariant solutions have the particularity that some reflection lines of the group G' are mapped at the origin.

Let us also mention a last example involving the discrete reflection group of the plane H , generated by the reflections with respect to the lines $x_2 = 0$, $x_2 = x_1$ and $x_1 = 1$. The point group associated to H is the group D_4 , and we can compose the canonical projection $H \rightarrow D_4$ either with




FIGURE 12. The correspondence of the fundamental domains by the f'' -equivariant solution u_R : on the right the fundamental domains of the discrete reflection group G' and on the left the fundamental domains of the finite reflection group D_2 .

the homomorphism $f_1 : D_4 \rightarrow D_2$ or with the homomorphism $g : D_4 \rightarrow D_2$ defined in subsection 6.1, to construct positive homomorphisms from H onto D_2 .

REFERENCES

- [1] N. D. Alikakos and G. Fusco. Entire solutions to equivariant elliptic systems with variational structure. *Arch. Rat. Mech. Anal.* **202** No. 2 (2011), pp. 567-597.
- [2] N. D. Alikakos. A new proof for the existence of an equivariant entire solution connecting the minima of the potential for the system $\Delta u - W_u(u) = 0$. *Comm. Partial Diff. Eqs.* **37** No. 12 (2012), pp. 2093-2115.
- [3] N. D. Alikakos and P. Smyrnelis. Existence of lattice solutions to semilinear elliptic systems with periodic potential. *Electron. J. Diff. Equ.* **15** (2012), pp. 1-15.
- [4] P. Bates, G. Fusco and P. Smyrnelis. Entire solutions with six-fold junctions to elliptic gradient systems with triangle symmetry. *Advanced Nonlinear Studies* **13** No. 1 (2013), pp. 1-12.
- [5] L. Bronsard, C. Gui, and M. Schatzman. A three-layered minimizer in \mathbb{R}^2 for a variational problem with a symmetric three-well potential. *Comm. Pure. Appl. Math.* **49** No. 7 (1996), pp. 677-715.
- [6] L. Bronsard and F. Reitich. On three-phase boundary motion and the singular limit of a vector-valued Ginzburg-Landau equation. *Arch. Rat. Mech. Anal.* **124** No. 4 (1993), pp. 355-379.
- [7] R. J. Braun, J. W. Cahn, G. B. MacFadden and A. A. Wheeler. Anisotropy of interfaces in an ordered alloy: a multiple-order-parameter model. *Trans. Roy. Soc. London, A.* **355** (1997), pp. 1787-
- [8] E. N. Dancer. New solutions for equations on \mathbb{R}^n . *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **30** No.3 4 (2002), pp. 535-563.
- [9] M. Del Pino, M. Kowalczyk, F. Pacard and J. Wei. Multiple-end solutions to the Allen-Cahn equation in \mathbb{R}^2 . *Jour. Funct. Anal.* **258** No. 2 (2010), pp. 458-503.
- [10] A. Farina and E. Valdinoci. The state of the art for a conjecture of De Giorgi and related problems. *Recent progress on reaction-diffusion systems and viscosity solutions*, Taichung, Taiwan, January, 2007. (Eds. Y. Du, H. Ishii and W.-Y. Lin) World Scientific, (2008) Hackensack, NJ, pp.74-96.
- [11] G. Fusco. *Equivariant entire solutions to the elliptic system $\Delta u = W_u(u)$ for general G -invariant potentials*, Calc. Var. Part. Diff. Eqs. February (2013), pp. 1-23.
- [12] G. Fusco. On some elementary properties of vector minimizers of the Allen-Cahn energy. *Comm. Pure Appl. Analysis*, **13** no. 3, (2014), 10451060.

- [13] L. C. Grove and C. T. Benson. *Finite reflection groups*. Graduate Texts in Mathematics **99**, Springer-Verlag, Berlin, second edition, 1985.
- [14] C. Gui and M. Schatzman. Symmetric quadruple phase transitions. *Ind. Univ. Math. J.* **57** No. 2 (2008), pp. 781–836.
- [15] D. Henry. *Geometric theory of semilinear parabolic equations*. Lecture Notes in Mathematics **840**, Springer-Verlag, Berlin-Heidelberg-New York, 1981.
- [16] J. E. Humphreys *Reflection groups and Coxeter groups*. Cambridge Studies in Advanced Mathematics **29**, Cambridge University Press, Cambridge, 1992.
- [17] J. Johnson The Number of Group Homomorphisms from D_m into D_n . Preprint. arXiv:1201.2363v3, (2012).
- [18] A. Malchiodi. Some new entire solutions of semilinear elliptic equations in \mathbb{R}^n . *Adv. Math.* **221** No. 6 (2009), pp. 1843–1909.
- [19] P. Smyrnelis. *Solutions to elliptic systems with mixed boundary conditions*. Phd thesis, University of Thessaloniki, 2012.
- [20] J. E. Taylor. The structure of singularities in soap-bubble-like and soap-film-like minimal surfaces. *Ann. Math.* **103** (1976), pp. 489–539.

(P. Bates) DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING, MICHIGAN 48824
E-mail address, P. Bates: bates@math.msu.edu

(G. Fusco) DIPARTIMENTO DI MATEMATICA PURA ED APPLICATA, UNIVERSITÀ DEGLI STUDI DELL'AQUILA, VIA VETOIO, 67010 COPPITO, L'AQUILA, ITALY
E-mail address, G. Fusco: fusco@univaq.it

(P. Smyrnelis) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ATHENS, PANEPISTEMIOPOLIS, 15784 ATHENS, GREECE
E-mail address, P. Smyrnelis: smpanos@math.uoa.gr